

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 September 2001 (27.09.2001)

PCT

(10) International Publication Number
WO 01/70977 A2

(51) International Patent Classification⁷: C12N 15/12,
15/62, 15/86, 5/16, C07K 14/71, 16/28, A01K 67/027,
G01N 33/68, A61K 38/17, 31/711, 49/00

Thomas, Charles [US/US]; 3010 Deer Valley Avenue,
Newbury Park, CA 91320 (US). COVEY, Todd [US/US];
12060 Alderbrook Street, Moorpark, CA 93021 (US).

(21) International Application Number: PCT/US01/09073

(74) Agent: NOONAN, Kevin, E.; McDonnell Boehnen Hul-
bert & Berghoff, Suite 3200, 300 South Wacker Drive,
Chicago, IL 60606 (US).

(22) International Filing Date: 22 March 2001 (22.03.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/191,379 22 March 2000 (22.03.2000) US

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ,
BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE,
ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP,
KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD,
MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD,
SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ,
VN, YU, ZA, ZW.

(63) Related by continuation (CON) or continuation-in-part
(CIP) to earlier application:
US 60/191,379 (CON)
Filed on 22 March 2000 (22.03.2000)

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): AMGEN,
INC. [US/US]; One Amgen Center Drive, Thousand Oaks,
CA 91320-1799 (US).

Published:

— *without international search report and to be republished
upon receipt of that report*

(71) Applicants and

(72) Inventors: SARIS, Christiaan, M. [NE/US]; 4027
Colonett Place, Newbury Park, CA 91320 (US). MU,
Sharon, X. [US/US]; 2954 Rikkard Drive, Thousand
Oaks, CA 91362 (US). XIA, Min [CA/US]; 3946 Calle
Buena Vista, Newbury Park, CA 91320 (US). BOONE,

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

WO 01/70977 A2

(54) Title: FIBROBLAST GROWTH FACTOR RECEPTOR-LIKE MOLECULES AND USES THEREOF

(57) Abstract: The present invention provides Fibroblast Growth Factor Receptor-Like (FGFR-L) polypeptides and nucleic acid molecules encoding the same. The invention also provides selective binding agents, vectors, host cells, and methods for producing FGFR-L polypeptides. The invention further provides pharmaceutical compositions and methods for the diagnosis, treatment, amelioration, and/or prevention of diseases, disorders, and conditions associated with FGFR-L polypeptides.

FIBROBLAST GROWTH FACTOR RECEPTOR-LIKE MOLECULES AND USES THEREOF

5 This application is a continuation of U.S. Provisional Patent Application No. 60/191,379, filed on March 22, 2000, the disclosure of which is explicitly incorporated by reference herein.

Field of the Invention

10 The present invention relates to Fibroblast Growth Factor Receptor-Like (FGFR-L) polypeptides and nucleic acid molecules encoding the same. The invention also relates to selective binding agents, vectors, host cells, and methods for producing FGFR-L polypeptides. The invention further relates to pharmaceutical compositions and methods for the diagnosis, treatment, amelioration, and/or prevention of diseases, disorders, and conditions associated with FGFR-L polypeptides.

15

Background of the Invention

Technical advances in the identification, cloning, expression, and manipulation of nucleic acid molecules and the deciphering of the human genome have greatly accelerated the discovery of novel therapeutics. Rapid nucleic acid sequencing techniques can now generate sequence information at unprecedented rates and, coupled with computational analyses, allow the assembly of overlapping sequences into partial and entire genomes and the identification of polypeptide-encoding regions. A comparison of a predicted amino acid sequence against a database compilation of known amino acid sequences allows one to determine the extent of homology to previously identified sequences and/or structural landmarks. The cloning and expression of a polypeptide-encoding region of a nucleic acid molecule provides a polypeptide product for structural and functional analyses. The manipulation of nucleic acid molecules and encoded polypeptides may confer advantageous properties on a product for use as a therapeutic.

30 In spite of the significant technical advances in genome research over the past decade, the potential for the development of novel therapeutics based on the human genome is still largely unrealized. Many genes encoding potentially beneficial polypeptide therapeutics or those encoding polypeptides, which may act as "targets"

for therapeutic molecules, have still not been identified. Accordingly, it is an object of the invention to identify novel polypeptides, and nucleic acid molecules encoding the same, which have diagnostic or therapeutic benefit.

5 Summary of the Invention

The present invention relates to novel FGFR-L nucleic acid molecules and encoded polypeptides.

The invention provides for an isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- 10 (a) the nucleotide sequence as set forth in either SEQ ID NO: 1 or SEQ ID NO: 4;
- (b) the nucleotide sequence of the DNA insert in ATCC Deposit No. _____;
- (c) a nucleotide sequence encoding the polypeptide as set forth in either
- 15 SEQ ID NO: 2 or SEQ ID NO: 5;
- (d) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a) - (c); and
- (e) a nucleotide sequence complementary to any of (a) - (c).

20 The invention also provides for an isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- (a) a nucleotide sequence encoding a polypeptide which is at least about 70 percent identical to the polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5, wherein the encoded polypeptide has an activity of the polypeptide set forth in
- 25 either SEQ ID NO: 2 or SEQ ID NO: 5;
- (b) a nucleotide sequence encoding an allelic variant or splice variant of the nucleotide sequence as set forth in either SEQ ID NO: 1 or SEQ ID NO: 4, the nucleotide sequence of the DNA insert in ATCC Deposit No. _____, or (a);
- (c) a region of the nucleotide sequence of either SEQ ID NO: 1 or SEQ ID
- 30 NO: 4, the DNA insert in ATCC Deposit No. _____, (a), or (b) encoding a polypeptide fragment of at least about 25 amino acid residues, wherein the polypeptide fragment has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5, or is antigenic;

(d) a region of the nucleotide sequence of either SEQ ID NO: 1 or SEQ ID NO: 4, the DNA insert in ATCC Deposit No. _____, or any of (a) - (c) comprising a fragment of at least about 16 nucleotides;

5 (e) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a) - (d); and

(f) a nucleotide sequence complementary to any of (a) - (d).

The invention further provides for an isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

10 (a) a nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one conservative amino acid substitution, wherein the encoded polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

15 (b) a nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one amino acid insertion, wherein the encoded polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

20 (c) a nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one amino acid deletion, wherein the encoded polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

25 (d) a nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 which has a C- and/or N- terminal truncation, wherein the encoded polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

30 (e) a nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one modification selected from the group consisting of amino acid substitutions, amino acid insertions, amino acid deletions, C-terminal truncation, and N-terminal truncation, wherein the encoded polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

(f) a nucleotide sequence of any of (a) - (e) comprising a fragment of at least about 16 nucleotides;

(g) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a) - (f); and

(h) a nucleotide sequence complementary to any of (a) - (e).

5 The present invention provides for an isolated polypeptide comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5; and

10 (b) the amino acid sequence encoded by the DNA insert in ATCC Deposit No. _____.

The invention also provides for an isolated polypeptide comprising the amino acid sequence selected from the group consisting of

15 (a) the amino acid sequence as set forth in SEQ ID NO: 3 or SEQ ID NO: 6, optionally further comprising an amino-terminal methionine;

(b) an amino acid sequence for an ortholog of either SEQ ID NO: 2 or SEQ ID NO: 5;

20 (c) an amino acid sequence which is at least about 70 percent identical to the amino acid sequence of either SEQ ID NO: 2 or SEQ ID NO: 5, wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

25 (d) a fragment of the amino acid sequence set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 comprising at least about 25 amino acid residues, wherein the fragment has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5, or is antigenic; and

(e) an amino acid sequence for an allelic variant or splice variant of the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5, the amino acid sequence encoded by the DNA insert in ATCC Deposit No. _____, or any of (a) - (c).

30

The invention further provides for an isolated polypeptide comprising the amino acid sequence selected from the group consisting of:

(a) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one conservative amino acid substitution, wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

5 (b) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one amino acid insertion, wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

(c) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one amino acid deletion, wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

10 (d) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 which has a C- and/or N-terminal truncation, wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5; and

(e) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one modification selected from the group consisting of amino acid substitutions, amino acid insertions, amino acid deletions, C-terminal truncation, and N-terminal truncation, wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5.

20 Also provided are fusion polypeptides comprising FGFR-L amino acid sequences.

The present invention also provides for an expression vector comprising the isolated nucleic acid molecules as set forth herein, recombinant host cells comprising the recombinant nucleic acid molecules as set forth herein, and a method of producing an FGFR-L polypeptide comprising culturing the host cells and optionally isolating the polypeptide so produced.

25 A transgenic non-human animal comprising a nucleic acid molecule encoding an FGFR-L polypeptide is also encompassed by the invention. The FGFR-L nucleic acid molecules are introduced into the animal in a manner that allows expression and increased levels of an FGFR-L polypeptide, which may include increased circulating levels. Alternatively, the FGFR-L nucleic acid molecules are introduced into the animal in a manner that prevents expression of endogenous FGFR-L polypeptide (*i.e.*, generates a transgenic animal possessing an FGFR-L polypeptide gene knockout).

The transgenic non-human animal is preferably a mammal, and more preferably a rodent, such as a rat or a mouse.

Also provided are derivatives of the FGFR-L polypeptides of the present invention.

5 Additionally provided are selective binding agents such as antibodies and peptides capable of specifically binding the FGFR-L polypeptides of the invention. Such antibodies and peptides may be agonistic or antagonistic.

10 Pharmaceutical compositions comprising the nucleotides, polypeptides, or selective binding agents of the invention and one or more pharmaceutically acceptable formulation agents are also encompassed by the invention. The pharmaceutical compositions are used to provide therapeutically effective amounts of the nucleotides or polypeptides of the present invention. The invention is also directed to methods of using the polypeptides, nucleic acid molecules, and selective binding agents.

15 The FGFR-L polypeptides and nucleic acid molecules of the present invention may be used to treat, prevent, ameliorate, and/or detect diseases and disorders, including those recited herein.

20 The present invention also provides a method of assaying test molecules to identify a test molecule that binds to an FGFR-L polypeptide. The method comprises contacting an FGFR-L polypeptide with a test molecule to determine the extent of binding of the test molecule to the polypeptide. The method further comprises determining whether such test molecules are agonists or antagonists of an FGFR-L polypeptide. The present invention further provides a method of testing the impact of molecules on the expression of FGFR-L polypeptide or on the activity of FGFR-L polypeptide.

25 Methods of regulating expression and modulating (*i.e.*, increasing or decreasing) levels of an FGFR-L polypeptide are also encompassed by the invention. One method comprises administering to an animal a nucleic acid molecule encoding an FGFR-L polypeptide. In another method, a nucleic acid molecule comprising elements that regulate or modulate the expression of an FGFR-L polypeptide may be administered. Examples of these methods include gene therapy, cell therapy, and anti-sense therapy as further described herein.

30

The FGFR-L polypeptide can be used for identifying ligands thereof. Various forms of "expression cloning" have been used for cloning ligands for receptors (*e.g.*,

Davis *et al.*, 1996, *Cell*, 87:1161-69). These and other FGFR-L polypeptide ligand cloning experiments are described in greater detail herein. Isolation of an FGFR-L polypeptide ligand allows for the identification or development of novel agonists or antagonists of the FGFR-L polypeptide signaling pathway. Such agonists and antagonists include FGFR-L polypeptide ligands, anti-FGFR-L polypeptide ligand antibodies and derivatives thereof, small molecules, or antisense oligonucleotides, any of which can be used for potentially treating one or more diseases or disorders, including those recited herein.

10 Brief Description of the Figures

Figures 1A-1C illustrate the nucleotide sequence of the murine FGFR-L gene (SEQ ID NO: 1) and the deduced amino acid sequence of murine FGFR-L polypeptide (SEQ ID NO: 2). The predicted signal peptide (underline) and transmembrane domain (double-underline) are indicated;

15

Figures 2A-2B illustrate the amino acid sequence alignment of murine FGFR-L polypeptide (Smaf2-00017-f4; SEQ ID NO: 2) and Iberian ribbed newt (*Pleurodeles waltlii*) Fibroblast Growth Factor Receptor-4 (PIR:B49151; SEQ ID NO: 7);

20

Figures 3A-3B illustrate the nucleotide sequence of a cDNA clone encoding the N-terminal portion of the human FGFR-L gene (SEQ ID NO: 4) and the deduced amino acid sequence of the N-terminal portion of the human FGFR-L polypeptide (SEQ ID NO: 5). The predicted signal peptide (underline) and transmembrane domain (double-underline) are indicated;

25

Figure 4 illustrates the amino acid sequence alignment of murine FGFR-L polypeptide (SEQ ID NO: 2) and a virtual human FGFR-L polypeptide sequence (SEQ ID NO: 8) constructed from residues 1-472 of SEQ ID NO: 5 and residues 473-504 of GenBank Accession No. AJ277437. The predicted signal peptide (underline), transmembrane domain (double-underline), and N-linked glycosylation sites (**bold**) are indicated;

30

Figure 5 illustrates the expression of FGFR-L mRNA as detected by Northern blot analysis in day 7, 11, 15, and 17 mouse embryos;

5 Figure 6 illustrates the expression of FGFR-L mRNA as detected by Northern blot analysis in murine heart, brain, spleen, lung, liver, skeletal muscle, kidney, and testis;

Figure 7 illustrates the expression of FGFR-L mRNA as detected by Northern blot analysis in NIH 3T3 cells and F10, F4, and D3 mouse bone marrow-derived stromal cell lines;

10 Figure 8 illustrates the expression of FGFR-L mRNA as detected by Northern blot analysis in human brain, heart, skeletal muscle, colon, thymus, spleen, kidney, liver, small intestine, placenta, lung, and peripheral blood leukocytes;

15 Figure 9 illustrates the expression of FGFR-L mRNA as detected by Northern blot analysis in promyelocytic leukemia HL-60 cells, HeLa S3 cells, chronic myelogenous leukemia L-562 cells, lymphoblastic leukemia MOLT-4 cells, Burkitt's lymphoma Raji cells, colorectal adenocarcinoma SW480 cells, lung carcinoma A549 cells, and melanoma G361 cells;

20 Figure 10 illustrates the expression of FGFR-L mRNA as detected by Northern blot analysis in human heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas;

25 Figure 11 illustrates the expression of FGFR-L mRNA as detected by Northern blot analysis in 266-6 cells, AR42J cells, CaPan I cells, HIG-82 cells, OHS4 cells, SW 1353 cells, SW 872 cells, K562 (old, *i.e.*, later passage) cells, K562 (new, *i.e.*, earlier passage) cells, Jurkat cells, and F4 cells;

30 Figures 12A-12B illustrate the expression of FGFR-L mRNA as detected by Northern blot analysis in human adipose tissue (using a human FGFR-L-derived probe) and murine adipose tissue (using a murine FGFR-L-derived probe);

Figure 13 illustrates the expression of FGFR-L mRNA in a number of murine tissues as detected in an RNase protection assay. The absence of the cyclophilin band in the pancreas RNA sample suggests that thi sample was degraded;

5 Figure 14 illustrates the expression of FGFR-L mRNA as detected by *in situ* hybridization in the peri-renal, white, and brown adipose tissue of a normal adult mouse (H&E = hematoxylin and eosin counterstaining; ISH = *in situ* hybridization);

10 Figure 15 illustrates the expression of FGFR-L mRNA as detected by *in situ* hybridization in the duodenum, ileum, colon, and pancreas of a normal adult mouse (H&E = hematoxylin and eosin counterstaining; ISH = *in situ* hybridization);

15 Figure 16 illustrates the expression of FGFR-L mRNA as detected by *in situ* hybridization in the trachea, articular cartilage of the knee joint, spleen, and uterus of a normal adult mouse (H&E = hematoxylin and eosin counterstaining; ISH = *in situ* hybridization);

20 Figure 17 illustrates the induction of FGFR-L mRNA in osteoblastic ST2 cells under conditions of osteoclastogenesis (*i.e.*, 5-day exposure to vitamin D3 and dexamethasone);

25 Figure 18 illustrates the results of Western blot analysis of *E.coli*-derived Des7-FGFR-L/ECD and CHO-derived FGFR-L/ECD-Fc proteins using FGFR-L polypeptide antiserum;

Figure 19 illustrates the results of Western blot analysis of murine eye (lane 1) and adipose tissue (lane 2) using FGFR-L polypeptide antiserum;

30 Figures 20A-20B illustrate the results of FACS analysis on F4 and D3 bone marrow stromal cells using FGFR-L polypeptide antiserum;

Figures 21A-21D illustrate the results of proliferation assays using D3 bone marrow stromal cells (either untransduced or transduced with a construct encoding FGFR-L

polypeptide) following 72 hour exposure to rhuPDGF (panel A), rhuFGF-2 (panel B), rhuFGF-4 (panel C), or rhuFGF-6 (panel D);

5 Figure 22 illustrates the results of proliferation assays using A5-F bone marrow stromal cells following exposure to *E. coli*-derived Des7-FGFR-L/ECD protein and serum, PDGF, FGF-2, FGF-4, or FGF-6;

10 Figure 23 illustrates the results of proliferation assays using A5-F bone marrow stromal cells following exposure to CHO-derived FGFR-L/ECD-Fc protein and serum, PDGF, FGF-4, or FGF-6;

15 Figure 24 illustrates the expression of the neomycin resistance gene as detected by Northern blot analysis of peripheral blood mononuclear cell (PBMN) RNA from two FGFR-L/neo-transduced mice (lanes 1 and 2) and two neo-transduced control mice (lanes 3 and 4).

Detailed Description of the Invention

20 The section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described. All references cited in this application are expressly incorporated by reference herein.

Definitions

25 The terms "FGFR-L gene" or "FGFR-L nucleic acid molecule" or "FGFR-L polynucleotide" refer to a nucleic acid molecule comprising or consisting of a nucleotide sequence as set forth in either SEQ ID NO: 1 or SEQ ID NO: 4, a nucleotide sequence encoding the polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5, a nucleotide sequence of the DNA insert in ATCC Deposit No. _____, and nucleic acid molecules as defined herein.

30 The term "FGFR-L polypeptide allelic variant" refers to one of several possible naturally occurring alternate forms of a gene occupying a given locus on a chromosome of an organism or a population of organisms.

The term "FGFR-L polypeptide splice variant" refers to a nucleic acid molecule, usually RNA, which is generated by alternative processing of intron

sequences in an RNA transcript of FGFR-L polypeptide amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5.

The term "isolated nucleic acid molecule" refers to a nucleic acid molecule of the invention that (1) has been separated from at least about 50 percent of proteins, lipids, carbohydrates, or other materials with which it is naturally found when total nucleic acid is isolated from the source cells, (2) is not linked to all or a portion of a polynucleotide to which the "isolated nucleic acid molecule" is linked in nature, (3) is operably linked to a polynucleotide which it is not linked to in nature, or (4) does not occur in nature as part of a larger polynucleotide sequence. Preferably, the isolated nucleic acid molecule of the present invention is substantially free from any other contaminating nucleic acid molecule(s) or other contaminants that are found in its natural environment that would interfere with its use in polypeptide production or its therapeutic, diagnostic, prophylactic or research use.

The term "nucleic acid sequence" or "nucleic acid molecule" refers to a DNA or RNA sequence. The term encompasses molecules formed from any of the known base analogs of DNA and RNA such as, but not limited to 4-acetylcytosine, 8-hydroxy-N6-methyladenosine, aziridinyl-cytosine, pseudoisocytosine, 5-(carboxyhydroxymethyl) uracil, 5-fluorouracil, 5-bromouracil, 5-carboxymethylaminomethyl-2-thiouracil, 5-carboxy-methylaminomethyluracil, dihydrouracil, inosine, N6-iso-pentenyladenine, 1-methyladenine, 1-methylpseudouracil, 1-methylguanine, 1-methylinosine, 2,2-dimethyl-guanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-methyladenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyamino-methyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarbonyl-methyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, oxybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, N-uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, pseudouracil, queosine, 2-thiocytosine, and 2,6-diaminopurine.

The term "vector" is used to refer to any molecule (e.g., nucleic acid, plasmid, or virus) used to transfer coding information to a host cell.

The term "expression vector" refers to a vector that is suitable for transformation of a host cell and contains nucleic acid sequences that direct and/or

control the expression of inserted heterologous nucleic acid sequences. Expression includes, but is not limited to, processes such as transcription, translation, and RNA splicing, if introns are present.

5 The term "operably linked" is used herein to refer to an arrangement of flanking sequences wherein the flanking sequences so described are configured or assembled so as to perform their usual function. Thus, a flanking sequence operably linked to a coding sequence may be capable of effecting the replication, transcription and/or translation of the coding sequence. For example, a coding sequence is operably linked to a promoter when the promoter is capable of directing transcription
10 of that coding sequence. A flanking sequence need not be contiguous with the coding sequence, so long as it functions correctly. Thus, for example, intervening untranslated yet transcribed sequences can be present between a promoter sequence and the coding sequence and the promoter sequence can still be considered "operably linked" to the coding sequence.

15 The term "host cell" is used to refer to a cell which has been transformed, or is capable of being transformed with a nucleic acid sequence and then of expressing a selected gene of interest. The term includes the progeny of the parent cell, whether or not the progeny is identical in morphology or in genetic make-up to the original parent, so long as the selected gene is present.

20 The term "FGFR-L polypeptide" refers to a polypeptide comprising the amino acid sequence of either SEQ ID NO: 2 or SEQ ID NO: 5 and related polypeptides. Related polypeptides include FGFR-L polypeptide fragments, FGFR-L polypeptide orthologs, FGFR-L polypeptide variants, and FGFR-L polypeptide derivatives, which possess at least one activity of the polypeptide as set forth in either SEQ ID NO: 2 or
25 SEQ ID NO: 5. FGFR-L polypeptides may be mature polypeptides, as defined herein, and may or may not have an amino-terminal methionine residue, depending on the method by which they are prepared.

The term "FGFR-L polypeptide fragment" refers to a polypeptide that comprises a truncation at the amino-terminus (with or without a leader sequence)
30 and/or a truncation at the carboxyl-terminus of the polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5. The term "FGFR-L polypeptide fragment" also refers to amino-terminal and/or carboxyl-terminal truncations of FGFR-L polypeptide orthologs, FGFR-L polypeptide derivatives, or FGFR-L polypeptide variants, or to

amino-terminal and/or carboxyl-terminal truncations of the polypeptides encoded by FGFR-L polypeptide allelic variants or FGFR-L polypeptide splice variants. FGFR-L polypeptide fragments may result from alternative RNA splicing or from *in vivo* protease activity. Membrane-bound forms of an FGFR-L polypeptide are also contemplated by the present invention. In preferred embodiments, truncations and/or deletions comprise about 10 amino acids, or about 20 amino acids, or about 50 amino acids, or about 75 amino acids, or about 100 amino acids, or more than about 100 amino acids. The polypeptide fragments so produced will comprise about 25 contiguous amino acids, or about 50 amino acids, or about 75 amino acids, or about 100 amino acids, or about 150 amino acids, or about 200 amino acids, or more than about 200 amino acids. Such FGFR-L polypeptide fragments may optionally comprise an amino-terminal methionine residue. It will be appreciated that such fragments can be used, for example, to generate antibodies to FGFR-L polypeptides.

The term "FGFR-L polypeptide ortholog" refers to a polypeptide from another species that corresponds to FGFR-L polypeptide amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5. For example, mouse and human FGFR-L polypeptides are considered orthologs of each other.

The term "FGFR-L polypeptide variants" refers to FGFR-L polypeptides comprising amino acid sequences having one or more amino acid sequence substitutions, deletions (such as internal deletions and/or FGFR-L polypeptide fragments), and/or additions (such as internal additions and/or FGFR-L fusion polypeptides) as compared to the FGFR-L polypeptide amino acid sequence set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 (with or without a leader sequence). Variants may be naturally occurring (*e.g.*, FGFR-L polypeptide allelic variants, FGFR-L polypeptide orthologs, and FGFR-L polypeptide splice variants) or artificially constructed. Such FGFR-L polypeptide variants may be prepared from the corresponding nucleic acid molecules having a DNA sequence that varies accordingly from the DNA sequence as set forth in either SEQ ID NO: 1 or SEQ ID NO: 4. In preferred embodiments, the variants have from 1 to 3, or from 1 to 5, or from 1 to 10, or from 1 to 15, or from 1 to 20, or from 1 to 25, or from 1 to 50, or from 1 to 75, or from 1 to 100, or more than 100 amino acid substitutions, insertions, additions and/or deletions, wherein the substitutions may be conservative, or non-conservative, or any combination thereof.

The term "FGFR-L polypeptide derivatives" refers to the polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5, FGFR-L polypeptide fragments, FGFR-L polypeptide orthologs, or FGFR-L polypeptide variants, as defined herein, that have been chemically modified. The term "FGFR-L polypeptide derivatives" also refers to the polypeptides encoded by FGFR-L polypeptide allelic variants or FGFR-L polypeptide splice variants, as defined herein, that have been chemically modified.

The term "mature FGFR-L polypeptide" refers to an FGFR-L polypeptide lacking a leader sequence. A mature FGFR-L polypeptide may also include other modifications such as proteolytic processing of the amino-terminus (with or without a leader sequence) and/or the carboxyl-terminus, cleavage of a smaller polypeptide from a larger precursor, N-linked and/or O-linked glycosylation, and the like. An exemplary mature FGFR-L polypeptide is depicted by the amino acid sequence of either SEQ ID NO: 3 or SEQ ID NO: 6.

The term "FGFR-L fusion polypeptide" refers to a fusion of one or more amino acids (such as a heterologous protein or peptide) at the amino- or carboxyl-terminus of the polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5, FGFR-L polypeptide fragments, FGFR-L polypeptide orthologs, FGFR-L polypeptide variants, or FGFR-L derivatives, as defined herein. The term "FGFR-L fusion polypeptide" also refers to a fusion of one or more amino acids at the amino- or carboxyl-terminus of the polypeptide encoded by FGFR-L polypeptide allelic variants or FGFR-L polypeptide splice variants, as defined herein.

The term "biologically active FGFR-L polypeptides" refers to FGFR-L polypeptides having at least one activity characteristic of the polypeptide comprising the amino acid sequence of either SEQ ID NO: 2 or SEQ ID NO: 5. In addition, an FGFR-L polypeptide may be active as an immunogen; that is, the FGFR-L polypeptide contains at least one epitope to which antibodies may be raised.

The term "isolated polypeptide" refers to a polypeptide of the present invention that (1) has been separated from at least about 50 percent of polynucleotides, lipids, carbohydrates, or other materials with which it is naturally found when isolated from the source cell, (2) is not linked (by covalent or noncovalent interaction) to all or a portion of a polypeptide to which the "isolated polypeptide" is linked in nature, (3) is operably linked (by covalent or noncovalent

interaction) to a polypeptide with which it is not linked in nature, or (4) does not occur in nature. Preferably, the isolated polypeptide is substantially free from any other contaminating polypeptides or other contaminants that are found in its natural environment that would interfere with its therapeutic, diagnostic, prophylactic or research use.

The term "identity," as known in the art, refers to a relationship between the sequences of two or more polypeptide molecules or two or more nucleic acid molecules, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between nucleic acid molecules or polypeptides, as the case may be, as determined by the match between strings of two or more nucleotide or two or more amino acid sequences. "Identity" measures the percent of identical matches between the smaller of two or more sequences with gap alignments (if any) addressed by a particular mathematical model or computer program (*i.e.*, "algorithms").

The term "similarity" is a related concept, but in contrast to "identity," "similarity" refers to a measure of relatedness which includes both identical matches and conservative substitution matches. If two polypeptide sequences have, for example, 10/20 identical amino acids, and the remainder are all non-conservative substitutions, then the percent identity and similarity would both be 50%. If in the same example, there are five more positions where there are conservative substitutions, then the percent identity remains 50%, but the percent similarity would be 75% (15/20). Therefore, in cases where there are conservative substitutions, the percent similarity between two polypeptides will be higher than the percent identity between those two polypeptides.

The term "naturally occurring" or "native" when used in connection with biological materials such as nucleic acid molecules, polypeptides, host cells, and the like, refers to materials which are found in nature and are not manipulated by man. Similarly, "non-naturally occurring" or "non-native" as used herein refers to a material that is not found in nature or that has been structurally modified or synthesized by man.

The terms "effective amount" and "therapeutically effective amount" each refer to the amount of an FGFR-L polypeptide or FGFR-L nucleic acid molecule used

to support an observable level of one or more biological activities of the FGFR-L polypeptides as set forth herein.

The term "pharmaceutically acceptable carrier" or "physiologically acceptable carrier" as used herein refers to one or more formulation materials suitable for accomplishing or enhancing the delivery of the FGFR-L polypeptide, FGFR-L nucleic acid molecule, or FGFR-L selective binding agent as a pharmaceutical composition.

The term "antigen" refers to a molecule or a portion of a molecule capable of being bound by a selective binding agent, such as an antibody, and additionally capable of being used in an animal to produce antibodies capable of binding to an epitope of that antigen. An antigen may have one or more epitopes.

The term "selective binding agent" refers to a molecule or molecules having specificity for an FGFR-L polypeptide. As used herein, the terms, "specific" and "specificity" refer to the ability of the selective binding agents to bind to human FGFR-L polypeptides and not to bind to human non-FGFR-L polypeptides. It will be appreciated, however, that the selective binding agents may also bind orthologs of the polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5, that is, interspecies versions thereof, such as mouse and rat FGFR-L polypeptides.

The term "transduction" is used to refer to the transfer of genes from one bacterium to another, usually by a phage. "Transduction" also refers to the acquisition and transfer of eukaryotic cellular sequences by retroviruses.

The term "transfection" is used to refer to the uptake of foreign or exogenous DNA by a cell, and a cell has been "transfected" when the exogenous DNA has been introduced inside the cell membrane. A number of transfection techniques are well known in the art and are disclosed herein. See, e.g., Graham *et al.*, 1973, *Virology* 52:456; Sambrook *et al.*, *Molecular Cloning, A Laboratory Manual* (Cold Spring Harbor Laboratories, 1989); Davis *et al.*, *Basic Methods in Molecular Biology* (Elsevier, 1986); and Chu *et al.*, 1981, *Gene* 13:197. Such techniques can be used to introduce one or more exogenous DNA moieties into suitable host cells.

The term "transformation" as used herein refers to a change in a cell's genetic characteristics, and a cell has been transformed when it has been modified to contain a new DNA. For example, a cell is transformed where it is genetically modified from its native state. Following transfection or transduction, the transforming DNA may recombine with that of the cell by physically integrating into a chromosome of the

cell, may be maintained transiently as an episomal element without being replicated, or may replicate independently as a plasmid. A cell is considered to have been stably transformed when the DNA is replicated with the division of the cell.

5 Relatedness of Nucleic Acid Molecules and/or Polypeptides

It is understood that related nucleic acid molecules include allelic or splice variants of the nucleic acid molecule of either SEQ ID NO: 1 or SEQ ID NO: 4, and include sequences which are complementary to any of the above nucleotide sequences. Related nucleic acid molecules also include a nucleotide sequence
10 encoding a polypeptide comprising or consisting essentially of a substitution, modification, addition and/or deletion of one or more amino acid residues compared to the polypeptide in either SEQ ID NO: 2 or SEQ ID NO: 5. Such related FGFR-L polypeptides may comprise, for example, an addition and/or a deletion of one or more N-linked or O-linked glycosylation sites or an addition and/or a deletion of one or
15 more cysteine residues.

Related nucleic acid molecules also include fragments of FGFR-L nucleic acid molecules which encode a polypeptide of at least about 25 contiguous amino acids, or about 50 amino acids, or about 75 amino acids, or about 100 amino acids, or about 150 amino acids, or about 200 amino acids, or more than about 200 amino acid
20 residues of the FGFR-L polypeptide of either SEQ ID NO: 2 or SEQ ID NO: 5.

In addition, related FGFR-L nucleic acid molecules also include those molecules which comprise nucleotide sequences which hybridize under moderately or highly stringent conditions as defined herein with the fully complementary sequence of the FGFR-L nucleic acid molecule of either SEQ ID NO: 1 or SEQ ID NO: 4, or of
25 a molecule encoding a polypeptide, which polypeptide comprises the amino acid sequence as shown in either SEQ ID NO: 2 or SEQ ID NO: 5, or of a nucleic acid fragment as defined herein, or of a nucleic acid fragment encoding a polypeptide as defined herein. Hybridization probes may be prepared using the FGFR-L sequences provided herein to screen cDNA, genomic or synthetic DNA libraries for related
30 sequences. Regions of the DNA and/or amino acid sequence of FGFR-L polypeptide that exhibit significant identity to known sequences are readily determined using sequence alignment algorithms as described herein and those regions may be used to design probes for screening.

The term "highly stringent conditions" refers to those conditions that are designed to permit hybridization of DNA strands whose sequences are highly complementary, and to exclude hybridization of significantly mismatched DNAs. Hybridization stringency is principally determined by temperature, ionic strength, and the concentration of denaturing agents such as formamide. Examples of "highly stringent conditions" for hybridization and washing are 0.015 M sodium FGFR-Loride, 0.0015 M sodium citrate at 65-68°C or 0.015 M sodium FGFR-Loride, 0.0015 M sodium citrate, and 50% formamide at 42°C. See Sambrook, Fritsch & Maniatis, *Molecular Cloning: A Laboratory Manual* (2nd ed., Cold Spring Harbor Laboratory, 1989); Anderson *et al.*, *Nucleic Acid Hybridisation: A Practical Approach* Ch. 4 (IRL Press Limited).

More stringent conditions (such as higher temperature, lower ionic strength, higher formamide, or other denaturing agent) may also be used – however, the rate of hybridization will be affected. Other agents may be included in the hybridization and washing buffers for the purpose of reducing non-specific and/or background hybridization. Examples are 0.1% bovine serum albumin, 0.1% polyvinylpyrrolidone, 0.1% sodium pyrophosphate, 0.1% sodium dodecylsulfate, NaDodSO₄, (SDS), ficoll, Denhardt's solution, sonicated salmon sperm DNA (or another non-complementary DNA), and dextran sulfate, although other suitable agents can also be used. The concentration and types of these additives can be changed without substantially affecting the stringency of the hybridization conditions. Hybridization experiments are usually carried out at pH 6.8-7.4; however, at typical ionic strength conditions, the rate of hybridization is nearly independent of pH. See Anderson *et al.*, *Nucleic Acid Hybridisation: A Practical Approach* Ch. 4 (IRL Press Limited).

Factors affecting the stability of DNA duplex include base composition, length, and degree of base pair mismatch. Hybridization conditions can be adjusted by one skilled in the art in order to accommodate these variables and allow DNAs of different sequence relatedness to form hybrids. The melting temperature of a perfectly matched DNA duplex can be estimated by the following equation:

$$T_m(^{\circ}\text{C}) = 81.5 + 16.6(\log[\text{Na}^+]) + 0.41(\%G+C) - 600/N - 0.72(\%\text{formamide})$$

where N is the length of the duplex formed, [Na⁺] is the molar concentration of the sodium ion in the hybridization or washing solution, %G+C is the percentage of

(guanine+cytosine) bases in the hybrid. For imperfectly matched hybrids, the melting temperature is reduced by approximately 1°C for each 1% mismatch.

The term "moderately stringent conditions" refers to conditions under which a DNA duplex with a greater degree of base pair mismatching than could occur under
5 "highly stringent conditions" is able to form. Examples of typical "moderately stringent conditions" are 0.015 M sodium FGFR-Loride, 0.0015 M sodium citrate at 50-65°C or 0.015 M sodium FGFR-Loride, 0.0015 M sodium citrate, and 20% formamide at 37-50°C. By way of example, "moderately stringent conditions" of 50°C in 0.015 M sodium ion will allow about a 21% mismatch.

10 It will be appreciated by those skilled in the art that there is no absolute distinction between "highly stringent conditions" and "moderately stringent conditions." For example, at 0.015 M sodium ion (no formamide), the melting temperature of perfectly matched long DNA is about 71°C. With a wash at 65°C (at the same ionic strength), this would allow for approximately a 6% mismatch. To
15 capture more distantly related sequences, one skilled in the art can simply lower the temperature or raise the ionic strength.

A good estimate of the melting temperature in 1M NaCl* for oligonucleotide probes up to about 20nt is given by:

$$T_m = 2^{\circ}\text{C per A-T base pair} + 4^{\circ}\text{C per G-C base pair}$$

20 *The sodium ion concentration in 6X salt sodium citrate (SSC) is 1M. See Suggs *et al.*, *Developmental Biology Using Purified Genes* 683 (Brown and Fox, eds., 1981).

High stringency washing conditions for oligonucleotides are usually at a temperature of 0-5°C below the T_m of the oligonucleotide in 6X SSC, 0.1% SDS.

In another embodiment, related nucleic acid molecules comprise or consist of
25 a nucleotide sequence that is at least about 70 percent identical to the nucleotide sequence as shown in either SEQ ID NO: 1 or SEQ ID NO: 4, or comprise or consist essentially of a nucleotide sequence encoding a polypeptide that is at least about 70 percent identical to the polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5. In preferred embodiments, the nucleotide sequences are about 75 percent, or
30 about 80 percent, or about 85 percent, or about 90 percent, or about 95, 96, 97, 98, or 99 percent identical to the nucleotide sequence as shown in either SEQ ID NO: 1 or SEQ ID NO: 4, or the nucleotide sequences encode a polypeptide that is about 75 percent, or about 80 percent, or about 85 percent, or about 90 percent, or about 95, 96,

97, 98, or 99 percent identical to the polypeptide sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5. Related nucleic acid molecules encode polypeptides possessing at least one activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5.

5 Differences in the nucleic acid sequence may result in conservative and/or non-conservative modifications of the amino acid sequence relative to the amino acid sequence of either SEQ ID NO: 2 or SEQ ID NO: 5.

Conservative modifications to the amino acid sequence of either SEQ ID NO: 2 or SEQ ID NO: 5 (and the corresponding modifications to the encoding nucleotides) will produce a polypeptide having functional and chemical characteristics similar to those of FGFR-L polypeptides. In contrast, substantial modifications in the functional and/or chemical characteristics of FGFR-L polypeptides may be accomplished by selecting substitutions in the amino acid sequence of either SEQ ID NO: 2 or SEQ ID NO: 5 that differ significantly in their effect on maintaining (a) the structure of the molecular backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain.

For example, a "conservative amino acid substitution" may involve a substitution of a native amino acid residue with a nonnative residue such that there is little or no effect on the polarity or charge of the amino acid residue at that position. Furthermore, any native residue in the polypeptide may also be substituted with alanine, as has been previously described for "alanine scanning mutagenesis."

Conservative amino acid substitutions also encompass non-naturally occurring amino acid residues that are typically incorporated by chemical peptide synthesis rather than by synthesis in biological systems. These include peptidomimetics, and other reversed or inverted forms of amino acid moieties.

Naturally occurring residues may be divided into classes based on common side chain properties:

- 1) hydrophobic: norleucine, Met, Ala, Val, Leu, Ile;
- 2) neutral hydrophilic: Cys, Ser, Thr;
- 3) acidic: Asp, Glu;
- 4) basic: Asn, Gln, His, Lys, Arg;
- 5) residues that influence chain orientation: Gly, Pro; and

6) aromatic: Trp, Tyr, Phe.

For example, non-conservative substitutions may involve the exchange of a member of one of these classes for a member from another class. Such substituted residues may be introduced into regions of the human FGFR-L polypeptide that are homologous with non-human FGFR-L polypeptides, or into the non-homologous regions of the molecule.

In making such changes, the hydropathic index of amino acids may be considered. Each amino acid has been assigned a hydropathic index on the basis of its hydrophobicity and charge characteristics. The hydropathic indices are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (-0.4); threonine (-0.7); serine (-0.8); tryptophan (-0.9); tyrosine (-1.3); proline (-1.6); histidine (-3.2); glutamate (-3.5); glutamine (-3.5); aspartate (-3.5); asparagine (-3.5); lysine (-3.9); and arginine (-4.5).

The importance of the hydropathic amino acid index in conferring interactive biological function on a protein is generally understood in the art (Kyte *et al.*, 1982, *J. Mol. Biol.* 157:105-31). It is known that certain amino acids may be substituted for other amino acids having a similar hydropathic index or score and still retain a similar biological activity. In making changes based upon the hydropathic index, the substitution of amino acids whose hydropathic indices are within ± 2 is preferred, those which are within ± 1 are particularly preferred, and those within ± 0.5 are even more particularly preferred.

It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity, particularly where the biologically functionally equivalent protein or peptide thereby created is intended for use in immunological embodiments, as in the present case. The greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with its immunogenicity and antigenicity, *i.e.*, with a biological property of the protein.

The following hydrophilicity values have been assigned to these amino acid residues: arginine (+3.0); lysine (+3.0); aspartate (+3.0 \pm 1); glutamate (+3.0 \pm 1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (-0.4); proline (-0.5 \pm 1); alanine (-0.5); histidine (-0.5); cysteine (-1.0); methionine (-1.3); valine (-1.5); leucine (-1.8); isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5);

and tryptophan (-3.4). In making changes based upon similar hydrophilicity values, the substitution of amino acids whose hydrophilicity values are within ± 2 is preferred, those which are within ± 1 are particularly preferred, and those within ± 0.5 are even more particularly preferred. One may also identify epitopes from primary amino acid sequences on the basis of hydrophilicity. These regions are also referred to as "epitopic core regions."

Desired amino acid substitutions (whether conservative or non-conservative) can be determined by those skilled in the art at the time such substitutions are desired. For example, amino acid substitutions can be used to identify important residues of the FGFR-L polypeptide, or to increase or decrease the affinity of the FGFR-L polypeptides described herein. Exemplary amino acid substitutions are set forth in Table I.

Table I

Amino Acid Substitutions

Original Residues	Exemplary Substitutions	Preferred Substitutions
Ala	Val, Leu, Ile	Val
Arg	Lys, Gln, Asn	Lys
Asn	Gln	Gln
Asp	Glu	Glu
Cys	Ser, Ala	Ser
Gln	Asn	Asn
Glu	Asp	Asp
Gly	Pro, Ala	Ala
His	Asn, Gln, Lys, Arg	Arg
Ile	Leu, Val, Met, Ala, Phe, Norleucine	Leu
Leu	Norleucine, Ile, Val, Met, Ala, Phe	Ile
Lys	Arg, 1,4 Diamino-butyric Acid, Gln, Asn	Arg
Met	Leu, Phe, Ile	Leu

Phe	Leu, Val, Ile, Ala, Tyr	Leu
Pro	Ala	Gly
Ser	Thr, Ala, Cys	Thr
Thr	Ser	Ser
Trp	Tyr, Phe	Tyr
Tyr	Trp, Phe, Thr, Ser	Phe
Val	Ile, Met, Leu, Phe, Ala, Norleucine	Leu

A skilled artisan will be able to determine suitable variants of the polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 using well-known techniques. For identifying suitable areas of the molecule that may be changed without destroying biological activity, one skilled in the art may target areas not believed to be important for activity. For example, when similar polypeptides with similar activities from the same species or from other species are known, one skilled in the art may compare the amino acid sequence of an FGFR-L polypeptide to such similar polypeptides. With such a comparison, one can identify residues and portions of the molecules that are conserved among similar polypeptides. It will be appreciated that changes in areas of the FGFR-L molecule that are not conserved relative to such similar polypeptides would be less likely to adversely affect the biological activity and/or structure of an FGFR-L polypeptide. One skilled in the art would also know that, even in relatively conserved regions, one may substitute chemically similar amino acids for the naturally occurring residues while retaining activity (conservative amino acid residue substitutions). Therefore, even areas that may be important for biological activity or for structure may be subject to conservative amino acid substitutions without destroying the biological activity or without adversely affecting the polypeptide structure.

Additionally, one skilled in the art can review structure-function studies identifying residues in similar polypeptides that are important for activity or structure. In view of such a comparison, one can predict the importance of amino acid residues in an FGFR-L polypeptide that correspond to amino acid residues that are important for activity or structure in similar polypeptides. One skilled in the art may opt for

chemically similar amino acid substitutions for such predicted important amino acid residues of FGFR-L polypeptides.

One skilled in the art can also analyze the three-dimensional structure and amino acid sequence in relation to that structure in similar polypeptides. In view of such information, one skilled in the art may predict the alignment of amino acid residues of FGFR-L polypeptide with respect to its three dimensional structure. One skilled in the art may choose not to make radical changes to amino acid residues predicted to be on the surface of the protein, since such residues may be involved in important interactions with other molecules. Moreover, one skilled in the art may generate test variants containing a single amino acid substitution at each amino acid residue. The variants could be screened using activity assays known to those with skill in the art. Such variants could be used to gather information about suitable variants. For example, if one discovered that a change to a particular amino acid residue resulted in destroyed, undesirably reduced, or unsuitable activity, variants with such a change would be avoided. In other words, based on information gathered from such routine experiments, one skilled in the art can readily determine the amino acids where further substitutions should be avoided either alone or in combination with other mutations.

A number of scientific publications have been devoted to the prediction of secondary structure. See Moulton, 1996, *Curr. Opin. Biotechnol.* 7:422-27; Chou *et al.*, 1974, *Biochemistry* 13:222-45; Chou *et al.*, 1974, *Biochemistry* 113:211-22; Chou *et al.*, 1978, *Adv. Enzymol. Relat. Areas Mol. Biol.* 47:45-48; Chou *et al.*, 1978, *Ann. Rev. Biochem.* 47:251-276; and Chou *et al.*, 1979, *Biophys. J.* 26:367-84. Moreover, computer programs are currently available to assist with predicting secondary structure. One method of predicting secondary structure is based upon homology modeling. For example, two polypeptides or proteins which have a sequence identity of greater than 30%, or similarity greater than 40%, often have similar structural topologies. The recent growth of the protein structural database (PDB) has provided enhanced predictability of secondary structure, including the potential number of folds within the structure of a polypeptide or protein. See Holm *et al.*, 1999, *Nucleic Acids Res.* 27:244-47. It has been suggested that there are a limited number of folds in a given polypeptide or protein and that once a critical number of structures have

been resolved, structural prediction will become dramatically more accurate (Brenner *et al.*, 1997, *Curr. Opin. Struct. Biol.* 7:369-76).

Additional methods of predicting secondary structure include "threading" (Jones, 1997, *Curr. Opin. Struct. Biol.* 7:377-87; Sippl *et al.*, 1996, *Structure* 4:15-19), "profile analysis" (Bowie *et al.*, 1991, *Science*, 253:164-70; Gribskov *et al.*, 1990, *Methods Enzymol.* 183:146-59; Gribskov *et al.*, 1987, *Proc. Nat. Acad. Sci. U.S.A.* 84:4355-58), and "evolutionary linkage" (See Holm *et al.*, *supra*, and Brenner *et al.*, *supra*).

Preferred FGFR-L polypeptide variants include glycosylation variants wherein the number and/or type of glycosylation sites have been altered compared to the amino acid sequence set forth in either SEQ ID NO: 2 or SEQ ID NO: 5. In one embodiment, FGFR-L polypeptide variants comprise a greater or a lesser number of N-linked glycosylation sites than the amino acid sequence set forth in either SEQ ID NO: 2 or SEQ ID NO: 5. An N-linked glycosylation site is characterized by the sequence: Asn-X-Ser or Asn-X-Thr, wherein the amino acid residue designated as X may be any amino acid residue except proline. The substitution of amino acid residues to create this sequence provides a potential new site for the addition of an N-linked carbohydrate chain. Alternatively, substitutions that eliminate this sequence will remove an existing N-linked carbohydrate chain. Also provided is a rearrangement of N-linked carbohydrate chains wherein one or more N-linked glycosylation sites (typically those that are naturally occurring) are eliminated and one or more new N-linked sites are created. Additional preferred FGFR-L variants include cysteine variants, wherein one or more cysteine residues are deleted or substituted with another amino acid (*e.g.*, serine) as compared to the amino acid sequence set forth in either SEQ ID NO: 2 or SEQ ID NO: 5. Cysteine variants are useful when FGFR-L polypeptides must be refolded into a biologically active conformation such as after the isolation of insoluble inclusion bodies. Cysteine variants generally have fewer cysteine residues than the native protein, and typically have an even number to minimize interactions resulting from unpaired cysteines.

In other embodiments, related nucleic acid molecules comprise or consist of a nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one amino acid insertion and wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5, or a

nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one amino acid deletion and wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5. Related nucleic acid molecules also comprise or consist of a nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 wherein the polypeptide has a carboxyl- and/or amino-terminal truncation and further wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5. Related nucleic acid molecules also comprise or consist of a nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one modification selected from the group consisting of amino acid substitutions, amino acid insertions, amino acid deletions, carboxyl-terminal truncations, and amino-terminal truncations and wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5.

In addition, the polypeptide comprising the amino acid sequence of either SEQ ID NO: 2 or SEQ ID NO: 5, or other FGFR-L polypeptide, may be fused to a homologous polypeptide to form a homodimer or to a heterologous polypeptide to form a heterodimer. Heterologous peptides and polypeptides include, but are not limited to: an epitope to allow for the detection and/or isolation of an FGFR-L fusion polypeptide; a transmembrane receptor protein or a portion thereof, such as an extracellular domain or a transmembrane and intracellular domain; a ligand or a portion thereof which binds to a transmembrane receptor protein; an enzyme or portion thereof which is catalytically active; a polypeptide or peptide which promotes oligomerization, such as a leucine zipper domain; a polypeptide or peptide which increases stability, such as an immunoglobulin constant region; and a polypeptide which has a therapeutic activity different from the polypeptide comprising the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5, or other FGFR-L polypeptide.

Fusions can be made either at the amino-terminus or at the carboxyl-terminus of the polypeptide comprising the amino acid sequence set forth in either SEQ ID NO: 2 or SEQ ID NO: 5, or other FGFR-L polypeptide. Fusions may be direct with no linker or adapter molecule or may be through a linker or adapter molecule. A linker or adapter molecule may be one or more amino acid residues, typically from about 20 to about 50 amino acid residues. A linker or adapter molecule may also be designed

with a cleavage site for a DNA restriction endonuclease or for a protease to allow for the separation of the fused moieties. It will be appreciated that once constructed, the fusion polypeptides can be derivatized according to the methods described herein.

In a further embodiment of the invention, the polypeptide comprising the amino acid sequence of either SEQ ID NO: 2 or SEQ ID NO: 5, or other FGFR-L polypeptide, is fused to one or more domains of an Fc region of human IgG. Antibodies comprise two functionally independent parts, a variable domain known as "Fab," that binds an antigen, and a constant domain known as "Fc," that is involved in effector functions such as complement activation and attack by phagocytic cells. An Fc has a long serum half-life, whereas an Fab is short-lived. Capon *et al.*, 1989, *Nature* 337:525-31. When constructed together with a therapeutic protein, an Fc domain can provide longer half-life or incorporate such functions as Fc receptor binding, protein A binding, complement fixation, and perhaps even placental transfer. *Id.* Table II summarizes the use of certain Fc fusions known in the art.

Table II

Fc Fusion with Therapeutic Proteins

Form of Fc	Fusion partner	Therapeutic implications	Reference
IgG1	N-terminus of CD30-L	Hodgkin's disease; anaplastic lymphoma; T-cell leukemia	U.S. Patent No. 5,480,981
Murine Fcγ2a	IL-10	anti-inflammatory; transplant rejection	Zheng <i>et al.</i> , 1995, <i>J. Immunol.</i> 154:5590-600
IgG1	TNF receptor	septic shock	Fisher <i>et al.</i> , 1996, <i>N. Engl. J. Med.</i> 334:1697-1702; Van Zee <i>et al.</i> , 1996, <i>J. Immunol.</i> 156:2221-30
IgG, IgA, IgM, or IgE (excluding the first domain)	TNF receptor	inflammation, autoimmune disorders	U.S. Patent No. 5,808,029
IgG1	CD4 receptor	AIDS	Capon <i>et al.</i> , 1989, <i>Nature</i> 337: 525-31
IgG1, IgG3	N-terminus of IL-2	anti-cancer, antiviral	Harvill <i>et al.</i> , 1995, <i>Immunotech.</i> 1:95-105
IgG1	C-terminus of OPG	osteoarthritis; bone density	WO 97/23614
IgG1	N-terminus of leptin	anti-obesity	PCT/US 97/23183, filed December 11, 1997

Human Ig Cy1	CTLA-4	autoimmune disorders	Linsley, 1991, <i>J. Exp. Med.</i> , 174:561-69
--------------	--------	----------------------	---

In one example, a human IgG hinge, CH2, and CH3 region may be fused at either the amino-terminus or carboxyl-terminus of the FGFR-L polypeptides using methods known to the skilled artisan. In another example, a human IgG hinge, CH2, and CH3 region may be fused at either the amino-terminus or carboxyl-terminus of an FGFR-L polypeptide fragment (e.g., the predicted extracellular portion of FGFR-L polypeptide).

The resulting FGFR-L fusion polypeptide may be purified by use of a Protein A affinity column. Peptides and proteins fused to an Fc region have been found to exhibit a substantially greater half-life *in vivo* than the unfused counterpart. Also, a fusion to an Fc region allows for dimerization/multimerization of the fusion polypeptide. The Fc region may be a naturally occurring Fc region, or may be altered to improve certain qualities, such as therapeutic qualities, circulation time, or reduced aggregation.

Identity and similarity of related nucleic acid molecules and polypeptides are readily calculated by known methods. Such methods include, but are not limited to those described in *Computational Molecular Biology* (A.M. Lesk, ed., Oxford University Press 1988); *Biocomputing: Informatics and Genome Projects* (D.W. Smith, ed., Academic Press 1993); *Computer Analysis of Sequence Data* (Part 1, A.M. Griffin and H.G. Griffin, eds., Humana Press 1994); G. von Heinle, *Sequence Analysis in Molecular Biology* (Academic Press 1987); *Sequence Analysis Primer* (M. Gribskov and J. Devereux, eds., M. Stockton Press 1991); and Carillo *et al.*, 1988, *SIAM J. Applied Math.*, 48:1073.

Preferred methods to determine identity and/or similarity are designed to give the largest match between the sequences tested. Methods to determine identity and similarity are described in publicly available computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, the GCG program package, including GAP (Devereux *et al.*, 1984, *Nucleic Acids Res.* 12:387; Genetics Computer Group, University of Wisconsin, Madison, WI), BLASTP, BLASTN, and FASTA (Altschul *et al.*, 1990, *J. Mol. Biol.* 215:403-10). The BLASTX program is publicly available from the

National Center for Biotechnology Information (NCBI) and other sources (Altschul *et al.*, *BLAST Manual* (NCB NLM NIH, Bethesda, MD); Altschul *et al.*, 1990, *supra*). The well-known Smith Waterman algorithm may also be used to determine identity.

5 Certain alignment schemes for aligning two amino acid sequences may result in the matching of only a short region of the two sequences, and this small aligned region may have very high sequence identity even though there is no significant relationship between the two full-length sequences. Accordingly, in a preferred embodiment, the selected alignment method (GAP program) will result in an alignment that spans at least 50 contiguous amino acids of the claimed polypeptide.

10 For example, using the computer algorithm GAP (Genetics Computer Group, University of Wisconsin, Madison, WI), two polypeptides for which the percent sequence identity is to be determined are aligned for optimal matching of their respective amino acids (the "matched span," as determined by the algorithm). A gap opening penalty (which is calculated as 3X the average diagonal; the "average diagonal" is the average of the diagonal of the comparison matrix being used; the
15 "diagonal" is the score or number assigned to each perfect amino acid match by the particular comparison matrix) and a gap extension penalty (which is usually 0.1X the gap opening penalty), as well as a comparison matrix such as PAM 250 or BLOSUM 62 are used in conjunction with the algorithm. A standard comparison matrix is also
20 used by the algorithm (*see Dayhoff et al.*, 5 *Atlas of Protein Sequence and Structure* (Supp. 3 1978)(PAM250 comparison matrix); Henikoff *et al.*, 1992, *Proc. Natl. Acad. Sci USA* 89:10915-19 (BLOSUM 62 comparison matrix)).

Preferred parameters for polypeptide sequence comparison include the following:

25

Algorithm: Needleman and Wunsch, 1970, *J. Mol. Biol.* 48:443-53;

Comparison matrix: BLOSUM 62 (Henikoff *et al.*, *supra*);

Gap Penalty: 12

Gap Length Penalty: 4

30

Threshold of Similarity: 0

The GAP program is useful with the above parameters. The aforementioned parameters are the default parameters for polypeptide comparisons (along with no penalty for end gaps) using the GAP algorithm.

5 Preferred parameters for nucleic acid molecule sequence comparison include the following:

Algorithm: Needleman and Wunsch, *supra*;
Comparison matrix: matches = +10, mismatch = 0
Gap Penalty: 50
10 Gap Length Penalty: 3

The GAP program is also useful with the above parameters. The aforementioned parameters are the default parameters for nucleic acid molecule comparisons.

15 Other exemplary algorithms, gap opening penalties, gap extension penalties, comparison matrices, and thresholds of similarity may be used, including those set forth in the Program Manual, Wisconsin Package, Version 9, September, 1997. The particular choices to be made will be apparent to those of skill in the art and will depend on the specific comparison to be made, such as DNA-to-DNA, protein-to-protein, protein-to-DNA; and additionally, whether the comparison is between given
20 pairs of sequences (in which case GAP or BestFit are generally preferred) or between one sequence and a large database of sequences (in which case FASTA or BLASTA are preferred).

Nucleic Acid Molecules

25 The nucleic acid molecules encoding a polypeptide comprising the amino acid sequence of an FGFR-L polypeptide can readily be obtained in a variety of ways including, without limitation, chemical synthesis, cDNA or genomic library screening, expression library screening, and/or PCR amplification of cDNA.

30 Recombinant DNA methods used herein are generally those set forth in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1989) and/or *Current Protocols in Molecular Biology* (Ausubel *et al.*, eds., Green Publishers Inc. and Wiley and Sons 1994). The invention provides for nucleic acid molecules as described herein and methods for obtaining such molecules.

Where a gene encoding the amino acid sequence of an FGFR-L polypeptide has been identified from one species, all or a portion of that gene may be used as a probe to identify orthologs or related genes from the same species. The probes or primers may be used to screen cDNA libraries from various tissue sources believed to express the FGFR-L polypeptide. In addition, part or all of a nucleic acid molecule having the sequence as set forth in either SEQ ID NO: 1 or SEQ ID NO: 4 may be used to screen a genomic library to identify and isolate a gene encoding the amino acid sequence of an FGFR-L polypeptide. Typically, conditions of moderate or high stringency will be employed for screening to minimize the number of false positives obtained from the screening.

Nucleic acid molecules encoding the amino acid sequence of FGFR-L polypeptides may also be identified by expression cloning which employs the detection of positive clones based upon a property of the expressed protein. Typically, nucleic acid libraries are screened by the binding an antibody or other binding partner (*e.g.*, receptor or ligand) to cloned proteins that are expressed and displayed on a host cell surface. The antibody or binding partner is modified with a detectable label to identify those cells expressing the desired clone.

Recombinant expression techniques conducted in accordance with the descriptions set forth below may be followed to produce these polynucleotides and to express the encoded polypeptides. For example, by inserting a nucleic acid sequence that encodes the amino acid sequence of an FGFR-L polypeptide into an appropriate vector, one skilled in the art can readily produce large quantities of the desired nucleotide sequence. The sequences can then be used to generate detection probes or amplification primers. Alternatively, a polynucleotide encoding the amino acid sequence of an FGFR-L polypeptide can be inserted into an expression vector. By introducing the expression vector into an appropriate host, the encoded FGFR-L polypeptide may be produced in large amounts.

Another method for obtaining a suitable nucleic acid sequence is the polymerase chain reaction (PCR). In this method, cDNA is prepared from poly(A)+RNA or total RNA using the enzyme reverse transcriptase. Two primers, typically complementary to two separate regions of cDNA encoding the amino acid sequence of an FGFR-L polypeptide, are then added to the cDNA along with a

polymerase such as *Taq* polymerase, and the polymerase amplifies the cDNA region between the two primers.

Another means of preparing a nucleic acid molecule encoding the amino acid sequence of an FGFR-L polypeptide is chemical synthesis using methods well known to the skilled artisan such as those described by Engels *et al.*, 1989, *Angew. Chem. Intl. Ed.* 28:716-34. These methods include, *inter alia*, the phosphotriester, phosphoramidite, and H-phosphonate methods for nucleic acid synthesis. A preferred method for such chemical synthesis is polymer-supported synthesis using standard phosphoramidite chemistry. Typically, the DNA encoding the amino acid sequence of an FGFR-L polypeptide will be several hundred nucleotides in length. Nucleic acids larger than about 100 nucleotides can be synthesized as several fragments using these methods. The fragments can then be ligated together to form the full-length nucleotide sequence of an FGFR-L gene. Usually, the DNA fragment encoding the amino-terminus of the polypeptide will have an ATG, which encodes a methionine residue. This methionine may or may not be present on the mature form of the FGFR-L polypeptide, depending on whether the polypeptide produced in the host cell is designed to be secreted from that cell. Other methods known to the skilled artisan may be used as well.

In certain embodiments, nucleic acid variants contain codons which have been altered for optimal expression of an FGFR-L polypeptide in a given host cell. Particular codon alterations will depend upon the FGFR-L polypeptide and host cell selected for expression. Such "codon optimization" can be carried out by a variety of methods, for example, by selecting codons which are preferred for use in highly expressed genes in a given host cell. Computer algorithms which incorporate codon frequency tables such as "Eco_high.Cod" for codon preference of highly expressed bacterial genes may be used and are provided by the University of Wisconsin Package Version 9.0 (Genetics Computer Group, Madison, WI). Other useful codon frequency tables include "Celegans_high.cod," "Celegans_low.cod," "Drosophila_high.cod," "Human_high.cod," "Maize_high.cod," and "Yeast_high.cod."

In some cases, it may be desirable to prepare nucleic acid molecules encoding FGFR-L polypeptide variants. Nucleic acid molecules encoding variants may be produced using site directed mutagenesis, PCR amplification, or other appropriate

methods, where the primer(s) have the desired point mutations (*see* Sambrook *et al.*,
supra, and Ausubel *et al.*, *supra*, for descriptions of mutagenesis techniques).
Chemical synthesis using methods described by Engels *et al.*, *supra*, may also be used
to prepare such variants. Other methods known to the skilled artisan may be used as
5 well.

Vectors and Host Cells

A nucleic acid molecule encoding the amino acid sequence of an FGFR-L
polypeptide is inserted into an appropriate expression vector using standard ligation
10 techniques. The vector is typically selected to be functional in the particular host cell
employed (*i.e.*, the vector is compatible with the host cell machinery such that
amplification of the gene and/or expression of the gene can occur). A nucleic acid
molecule encoding the amino acid sequence of an FGFR-L polypeptide may be
amplified/expressed in prokaryotic, yeast, insect (baculovirus systems) and/or
15 eukaryotic host cells. Selection of the host cell will depend in part on whether an
FGFR-L polypeptide is to be post-translationally modified (*e.g.*, glycosylated and/or
phosphorylated). If so, yeast, insect, or mammalian host cells are preferable. For a
review of expression vectors, *see Meth. Enz.*, vol. 185 (D.V. Goeddel, ed., Academic
Press 1990).

20 Typically, expression vectors used in any of the host cells will contain
sequences for plasmid maintenance and for cloning and expression of exogenous
nucleotide sequences. Such sequences, collectively referred to as "flanking
sequences" in certain embodiments will typically include one or more of the
following nucleotide sequences: a promoter, one or more enhancer sequences, an
25 origin of replication, a transcriptional termination sequence, a complete intron
sequence containing a donor and acceptor splice site, a sequence encoding a leader
sequence for polypeptide secretion, a ribosome binding site, a polyadenylation
sequence, a polylinker region for inserting the nucleic acid encoding the polypeptide
to be expressed, and a selectable marker element. Each of these sequences is
30 discussed below.

Optionally, the vector may contain a "tag"-encoding sequence, *i.e.*, an
oligonucleotide molecule located at the 5' or 3' end of the FGFR-L polypeptide
coding sequence; the oligonucleotide sequence encodes polyHis (such as hexaHis), or

another "tag" such as FLAG, HA (hemagglutinin influenza virus), or *myc* for which commercially available antibodies exist. This tag is typically fused to the polypeptide upon expression of the polypeptide, and can serve as a means for affinity purification of the FGFR-L polypeptide from the host cell. Affinity purification can be accomplished, for example, by column chromatography using antibodies against the tag as an affinity matrix. Optionally, the tag can subsequently be removed from the purified FGFR-L polypeptide by various means such as using certain peptidases for cleavage.

Flanking sequences may be homologous (*i.e.*, from the same species and/or strain as the host cell), heterologous (*i.e.*, from a species other than the host cell species or strain), hybrid (*i.e.*, a combination of flanking sequences from more than one source), or synthetic, or the flanking sequences may be native sequences which normally function to regulate FGFR-L polypeptide expression. As such, the source of a flanking sequence may be any prokaryotic or eukaryotic organism, any vertebrate or invertebrate organism, or any plant, provided that the flanking sequence is functional in, and can be activated by, the host cell machinery.

Flanking sequences useful in the vectors of this invention may be obtained by any of several methods well known in the art. Typically, flanking sequences useful herein – other than the FGFR-L gene flanking sequences – will have been previously identified by mapping and/or by restriction endonuclease digestion and can thus be isolated from the proper tissue source using the appropriate restriction endonucleases. In some cases, the full nucleotide sequence of a flanking sequence may be known. Here, the flanking sequence may be synthesized using the methods described herein for nucleic acid synthesis or cloning.

Where all or only a portion of the flanking sequence is known, it may be obtained using PCR and/or by screening a genomic library with a suitable oligonucleotide and/or flanking sequence fragment from the same or another species. Where the flanking sequence is not known, a fragment of DNA containing a flanking sequence may be isolated from a larger piece of DNA that may contain, for example, a coding sequence or even another gene or genes. Isolation may be accomplished by restriction endonuclease digestion to produce the proper DNA fragment followed by isolation using agarose gel purification, Qiagen® column chromatography (Chatsworth, CA), or other methods known to the skilled artisan. The selection of

suitable enzymes to accomplish this purpose will be readily apparent to one of ordinary skill in the art.

5 An origin of replication is typically a part of those prokaryotic expression vectors purchased commercially, and the origin aids in the amplification of the vector in a host cell. Amplification of the vector to a certain copy number can, in some cases, be important for the optimal expression of an FGFR-L polypeptide. If the vector of choice does not contain an origin of replication site, one may be chemically synthesized based on a known sequence, and ligated into the vector. For example, the origin of replication from the plasmid pBR322 (New England Biolabs, Beverly, MA)
10 is suitable for most gram-negative bacteria and various origins (*e.g.*, SV40, polyoma, adenovirus, vesicular stomatitis virus (VSV), or papillomaviruses such as HPV or BPV) are useful for cloning vectors in mammalian cells. Generally, the origin of replication component is not needed for mammalian expression vectors (for example, the SV40 origin is often used only because it contains the early promoter).

15 A transcription termination sequence is typically located 3' of the end of a polypeptide coding region and serves to terminate transcription. Usually, a transcription termination sequence in prokaryotic cells is a G-C rich fragment followed by a poly-T sequence. While the sequence is easily cloned from a library or even purchased commercially as part of a vector, it can also be readily synthesized
20 using methods for nucleic acid synthesis such as those described herein.

A selectable marker gene element encodes a protein necessary for the survival and growth of a host cell grown in a selective culture medium. Typical selection marker genes encode proteins that (a) confer resistance to antibiotics or other toxins, *e.g.*, ampicillin, tetracycline, or kanamycin for prokaryotic host cells; (b) complement
25 auxotrophic deficiencies of the cell; or (c) supply critical nutrients not available from complex media. Preferred selectable markers are the kanamycin resistance gene, the ampicillin resistance gene, and the tetracycline resistance gene. A neomycin resistance gene may also be used for selection in prokaryotic and eukaryotic host cells.

30 Other selection genes may be used to amplify the gene that will be expressed. Amplification is the process wherein genes that are in greater demand for the production of a protein critical for growth are reiterated in tandem within the chromosomes of successive generations of recombinant cells. Examples of suitable

selectable markers for mammalian cells include dihydrofolate reductase (DHFR) and thymidine kinase. The mammalian cell transformants are placed under selection pressure wherein only the transformants are uniquely adapted to survive by virtue of the selection gene present in the vector. Selection pressure is imposed by culturing
5 the transformed cells under conditions in which the concentration of selection agent in the medium is successively changed, thereby leading to the amplification of both the selection gene and the DNA that encodes an FGFR-L polypeptide. As a result, increased quantities of FGFR-L polypeptide are synthesized from the amplified DNA.

A ribosome binding site is usually necessary for translation initiation of
10 mRNA and is characterized by a Shine-Dalgarno sequence (prokaryotes) or a Kozak sequence (eukaryotes). The element is typically located 3' to the promoter and 5' to the coding sequence of an FGFR-L polypeptide to be expressed. The Shine-Dalgarno sequence is varied but is typically a polypurine (*i.e.*, having a high A-G content). Many Shine-Dalgarno sequences have been identified, each of which can be readily
15 synthesized using methods set forth herein and used in a prokaryotic vector.

A leader, or signal, sequence may be used to direct an FGFR-L polypeptide out of the host cell. Typically, a nucleotide sequence encoding the signal sequence is positioned in the coding region of an FGFR-L nucleic acid molecule, or directly at the 5' end of an FGFR-L polypeptide coding region. Many signal sequences have been
20 identified, and any of those that are functional in the selected host cell may be used in conjunction with an FGFR-L nucleic acid molecule. Therefore, a signal sequence may be homologous (naturally occurring) or heterologous to the FGFR-L nucleic acid molecule. Additionally, a signal sequence may be chemically synthesized using methods described herein. In most cases, the secretion of an FGFR-L polypeptide
25 from the host cell via the presence of a signal peptide will result in the removal of the signal peptide from the secreted FGFR-L polypeptide. The signal sequence may be a component of the vector, or it may be a part of an FGFR-L nucleic acid molecule that is inserted into the vector.

Included within the scope of this invention is the use of either a nucleotide
30 sequence encoding a native FGFR-L polypeptide signal sequence joined to an FGFR-L polypeptide coding region or a nucleotide sequence encoding a heterologous signal sequence joined to an FGFR-L polypeptide coding region. The heterologous signal sequence selected should be one that is recognized and processed, *i.e.*, cleaved by a

signal peptidase, by the host cell. For prokaryotic host cells that do not recognize and process the native FGFR-L polypeptide signal sequence, the signal sequence is substituted by a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, or heat-stable enterotoxin II leaders. For yeast secretion, the native FGFR-L polypeptide signal sequence may be substituted by the yeast invertase, alpha factor, or acid phosphatase leaders. In mammalian cell expression the native signal sequence is satisfactory, although other mammalian signal sequences may be suitable.

In some cases, such as where glycosylation is desired in a eukaryotic host cell expression system, one may manipulate the various presequences to improve glycosylation or yield. For example, one may alter the peptidase cleavage site of a particular signal peptide, or add pro-sequences, which also may affect glycosylation. The final protein product may have, in the -1 position (relative to the first amino acid of the mature protein) one or more additional amino acids incident to expression, which may not have been totally removed. For example, the final protein product may have one or two amino acid residues found in the peptidase cleavage site, attached to the amino-terminus. Alternatively, use of some enzyme cleavage sites may result in a slightly truncated form of the desired FGFR-L polypeptide, if the enzyme cuts at such area within the mature polypeptide.

In many cases, transcription of a nucleic acid molecule is increased by the presence of one or more introns in the vector; this is particularly true where a polypeptide is produced in eukaryotic host cells, especially mammalian host cells. The introns used may be naturally occurring within the FGFR-L gene especially where the gene used is a full-length genomic sequence or a fragment thereof. Where the intron is not naturally occurring within the gene (as for most cDNAs), the intron may be obtained from another source. The position of the intron with respect to flanking sequences and the FGFR-L gene is generally important, as the intron must be transcribed to be effective. Thus, when an FGFR-L cDNA molecule is being transcribed, the preferred position for the intron is 3' to the transcription start site and 5' to the poly-A transcription termination sequence. Preferably, the intron or introns will be located on one side or the other (*i.e.*, 5' or 3') of the cDNA such that it does not interrupt the coding sequence. Any intron from any source, including viral, prokaryotic and eukaryotic (plant or animal) organisms, may be used to practice this

invention, provided that it is compatible with the host cell into which it is inserted. Also included herein are synthetic introns. Optionally, more than one intron may be used in the vector.

5 The expression and cloning vectors of the present invention will typically contain a promoter that is recognized by the host organism and operably linked to the molecule encoding the FGFR-L polypeptide. Promoters are untranscribed sequences located upstream (*i.e.*, 5') to the start codon of a structural gene (generally within about 100 to 1000 bp) that control the transcription of the structural gene. Promoters are conventionally grouped into one of two classes: inducible promoters and
10 constitutive promoters. Inducible promoters initiate increased levels of transcription from DNA under their control in response to some change in culture conditions, such as the presence or absence of a nutrient or a change in temperature. Constitutive promoters, on the other hand, initiate continual gene product production; that is, there is little or no control over gene expression. A large number of promoters, recognized
15 by a variety of potential host cells, are well known. A suitable promoter is operably linked to the DNA encoding FGFR-L polypeptide by removing the promoter from the source DNA by restriction enzyme digestion and inserting the desired promoter sequence into the vector. The native FGFR-L promoter sequence may be used to direct amplification and/or expression of an FGFR-L nucleic acid molecule. A
20 heterologous promoter is preferred, however, if it permits greater transcription and higher yields of the expressed protein as compared to the native promoter, and if it is compatible with the host cell system that has been selected for use.

Promoters suitable for use with prokaryotic hosts include the beta-lactamase and lactose promoter systems; alkaline phosphatase; a tryptophan (trp) promoter
25 system; and hybrid promoters such as the tac promoter. Other known bacterial promoters are also suitable. Their sequences have been published, thereby enabling one skilled in the art to ligate them to the desired DNA sequence, using linkers or adapters as needed to supply any useful restriction sites.

Suitable promoters for use with yeast hosts are also well known in the art.
30 Yeast enhancers are advantageously used with yeast promoters. Suitable promoters for use with mammalian host cells are well known and include, but are not limited to, those obtained from the genomes of viruses such as polyoma virus, fowlpox virus, adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus,

cytomegalovirus, retroviruses, hepatitis-B virus and most preferably Simian Virus 40 (SV40). Other suitable mammalian promoters include heterologous mammalian promoters, for example, heat-shock promoters and the actin promoter.

Additional promoters which may be of interest in controlling FGFR-L gene
5 expression include, but are not limited to: the SV40 early promoter region (Bernoist and Chambon, 1981, *Nature* 290:304-10); the CMV promoter; the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto, *et al.*, 1980, *Cell* 22:787-97); the herpes thymidine kinase promoter (Wagner *et al.*, 1981, *Proc. Natl. Acad. Sci. U.S.A.* 78:1444-45); the regulatory sequences of the metallothionine gene
10 (Brinster *et al.*, 1982, *Nature* 296:39-42); prokaryotic expression vectors such as the beta-lactamase promoter (Villa-Kamaroff *et al.*, 1978, *Proc. Natl. Acad. Sci. U.S.A.*, 75:3727-31); or the tac promoter (DeBoer *et al.*, 1983, *Proc. Natl. Acad. Sci. U.S.A.*, 80:21-25). Also of interest are the following animal transcriptional control regions, which exhibit tissue specificity and have been utilized in transgenic animals: the
15 elastase I gene control region which is active in pancreatic acinar cells (Swift *et al.*, 1984, *Cell* 38:639-46; Ornitz *et al.*, 1986, *Cold Spring Harbor Symp. Quant. Biol.* 50:399-409 (1986); MacDonald, 1987, *Hepatology* 7:425-515); the insulin gene control region which is active in pancreatic beta cells (Hanahan, 1985, *Nature* 315:115-22); the immunoglobulin gene control region which is active in lymphoid
20 cells (Grosschedl *et al.*, 1984, *Cell* 38:647-58; Adames *et al.*, 1985, *Nature* 318:533-38; Alexander *et al.*, 1987, *Mol. Cell. Biol.*, 7:1436-44); the mouse mammary tumor virus control region which is active in testicular, breast, lymphoid and mast cells (Leder *et al.*, 1986, *Cell* 45:485-95); the albumin gene control region which is active in liver (Pinkert *et al.*, 1987, *Genes and Devel.* 1:268-76); the alpha-feto-protein gene
25 control region which is active in liver (Krumlauf *et al.*, 1985, *Mol. Cell. Biol.*, 5:1639-48; Hammer *et al.*, 1987, *Science* 235:53-58); the alpha 1-antitrypsin gene control region which is active in the liver (Kelsey *et al.*, 1987, *Genes and Devel.* 1:161-71); the beta-globin gene control region which is active in myeloid cells (Mogam *et al.*, 1985, *Nature* 315:338-40; Kollias *et al.*, 1986, *Cell* 46:89-94); the myelin basic
30 protein gene control region which is active in oligodendrocyte cells in the brain (Readhead *et al.*, 1987, *Cell* 48:703-12); the myosin light chain-2 gene control region which is active in skeletal muscle (Sani, 1985, *Nature* 314:283-86); and the

gonadotropic releasing hormone gene control region which is active in the hypothalamus (Mason *et al.*, 1986, *Science* 234:1372-78).

5 An enhancer sequence may be inserted into the vector to increase the transcription of a DNA encoding an FGFR-L polypeptide of the present invention by higher eukaryotes. Enhancers are cis-acting elements of DNA, usually about 10-300 bp in length, that act on the promoter to increase transcription. Enhancers are relatively orientation and position independent. They have been found 5' and 3' to the transcription unit. Several enhancer sequences available from mammalian genes are known (*e.g.*, globin, elastase, albumin, alpha-feto-protein and insulin). Typically, 10 however, an enhancer from a virus will be used. The SV40 enhancer, the cytomegalovirus early promoter enhancer, the polyoma enhancer, and adenovirus enhancers are exemplary enhancing elements for the activation of eukaryotic promoters. While an enhancer may be spliced into the vector at a position 5' or 3' to an FGFR-L nucleic acid molecule, it is typically located at a site 5' from the 15 promoter.

Expression vectors of the invention may be constructed from a starting vector such as a commercially available vector. Such vectors may or may not contain all of the desired flanking sequences. Where one or more of the flanking sequences described herein are not already present in the vector, they may be individually 20 obtained and ligated into the vector. Methods used for obtaining each of the flanking sequences are well known to one skilled in the art.

Preferred vectors for practicing this invention are those which are compatible with bacterial, insect, and mammalian host cells. Such vectors include, *inter alia*, pCRII, pCR3, and pcDNA3.1 (Invitrogen, San Diego, CA), pBSII (Stratagene, La Jolla, CA), pET15 (Novagen, Madison, WI), pGEX (Pharmacia Biotech, Piscataway, 25 NJ), pEGFP-N2 (Clontech, Palo Alto, CA), pETL (BlueBacII, Invitrogen), pDSR-alpha (PCT Pub. No. WO 90/14363) and pFastBacDual (Gibco-BRL, Grand Island, NY).

Additional suitable vectors include, but are not limited to, cosmids, plasmids, 30 or modified viruses, but it will be appreciated that the vector system must be compatible with the selected host cell. Such vectors include, but are not limited to plasmids such as Bluescript[®] plasmid derivatives (a high copy number ColE1-based phagemid, Stratagene Cloning Systems, La Jolla CA), PCR cloning plasmids

designed for cloning Taq-amplified PCR products (*e.g.*, TOPO™ TA Cloning® Kit, PCR2.1® plasmid derivatives, Invitrogen, Carlsbad, CA), and mammalian, yeast or virus vectors such as a baculovirus expression system (pBacPAK plasmid derivatives, Clontech, Palo Alto, CA).

5 After the vector has been constructed and a nucleic acid molecule encoding an FGFR-L polypeptide has been inserted into the proper site of the vector, the completed vector may be inserted into a suitable host cell for amplification and/or polypeptide expression. The transformation of an expression vector for an FGFR-L polypeptide into a selected host cell may be accomplished by well known methods
10 including methods such as transfection, infection, calcium FGFR-Loride, electroporation, microinjection, lipofection, DEAE-dextran method, or other known techniques. The method selected will in part be a function of the type of host cell to be used. These methods and other suitable methods are well known to the skilled artisan, and are set forth, for example, in Sambrook *et al.*, *supra*.

15 Host cells may be prokaryotic host cells (such as *E. coli*) or eukaryotic host cells (such as a yeast, insect, or vertebrate cell). The host cell, when cultured under appropriate conditions, synthesizes an FGFR-L polypeptide which can subsequently be collected from the culture medium (if the host cell secretes it into the medium) or directly from the host cell producing it (if it is not secreted). The selection of an
20 appropriate host cell will depend upon various factors, such as desired expression levels, polypeptide modifications that are desirable or necessary for activity (such as glycosylation or phosphorylation) and ease of folding into a biologically active molecule.

 A number of suitable host cells are known in the art and many are available
25 from the American Type Culture Collection (ATCC), Manassas, VA. Examples include, but are not limited to, mammalian cells, such as Chinese hamster ovary cells (CHO), CHO DHFR(-) cells (Urlaub *et al.*, 1980, *Proc. Natl. Acad. Sci. U.S.A.* 97:4216-20), human embryonic kidney (HEK) 293 or 293T cells, or 3T3 cells. The selection of suitable mammalian host cells and methods for transformation, culture,
30 amplification, screening, product production, and purification are known in the art. Other suitable mammalian cell lines, are the monkey COS-1 and COS-7 cell lines, and the CV-1 cell line. Further exemplary mammalian host cells include primate cell lines and rodent cell lines, including transformed cell lines. Normal diploid cells, cell

strains derived from *in vitro* culture of primary tissue, as well as primary explants, are also suitable. Candidate cells may be genotypically deficient in the selection gene, or may contain a dominantly acting selection gene. Other suitable mammalian cell lines include but are not limited to, mouse neuroblastoma N2A cells, HeLa, mouse L-929
5 cells, 3T3 lines derived from Swiss, Balb-c or NIH mice, BHK or HaK hamster cell lines. Each of these cell lines is known by and available to those skilled in the art of protein expression.

Similarly useful as host cells suitable for the present invention are bacterial cells. For example, the various strains of *E. coli* (e.g., HB101, DH5 α , DH10, and
10 MC1061) are well-known as host cells in the field of biotechnology. Various strains of *B. subtilis*, *Pseudomonas spp.*, other *Bacillus spp.*, *Streptomyces spp.*, and the like may also be employed in this method.

Many strains of yeast cells known to those skilled in the art are also available as host cells for the expression of the polypeptides of the present invention. Preferred
15 yeast cells include, for example, *Saccharomyces cerevisiae* and *Pichia pastoris*.

Additionally, where desired, insect cell systems may be utilized in the methods of the present invention. Such systems are described, for example, in Kitts
20 *et al.*, 1993, *Biotechniques*, 14:810-17; Lucklow, 1993, *Curr. Opin. Biotechnol.* 4:564-72; and Lucklow *et al.*, 1993, *J. Virol.*, 67:4566-79. Preferred insect cells are Sf-9 and Hi5 (Invitrogen).

One may also use transgenic animals to express glycosylated FGFR-L polypeptides. For example, one may use a transgenic milk-producing animal (a cow or goat, for example) and obtain the present glycosylated polypeptide in the animal milk. One may also use plants to produce FGFR-L polypeptides, however, in general,
25 the glycosylation occurring in plants is different from that produced in mammalian cells, and may result in a glycosylated product which is not suitable for human therapeutic use.

Polypeptide Production

30 Host cells comprising an FGFR-L polypeptide expression vector may be cultured using standard media well known to the skilled artisan. The media will usually contain all nutrients necessary for the growth and survival of the cells. Suitable media for culturing *E. coli* cells include, for example, Luria Broth (LB)

and/or Terrific Broth (TB). Suitable media for culturing eukaryotic cells include Roswell Park Memorial Institute medium 1640 (RPMI 1640), Minimal Essential Medium (MEM) and/or Dulbecco's Modified Eagle Medium (DMEM), all of which may be supplemented with serum and/or growth factors as necessary for the particular
5 cell line being cultured. A suitable medium for insect cultures is Grace's medium supplemented with yeastolate, lactalbumin hydrolysate, and/or fetal calf serum as necessary.

Typically, an antibiotic or other compound useful for selective growth of transfected or transformed cells is added as a supplement to the media. The
10 compound to be used will be dictated by the selectable marker element present on the plasmid with which the host cell was transformed. For example, where the selectable marker element is kanamycin resistance, the compound added to the culture medium will be kanamycin. Other compounds for selective growth include ampicillin, tetracycline, and neomycin.

15 The amount of an FGFR-L polypeptide produced by a host cell can be evaluated using standard methods known in the art. Such methods include, without limitation, Western blot analysis, SDS-polyacrylamide gel electrophoresis, non-denaturing gel electrophoresis, High Performance Liquid Chromatography (HPLC) separation, immunoprecipitation, and/or activity assays such as DNA binding gel shift
20 assays.

If an FGFR-L polypeptide has been designed to be secreted from the host cells, the majority of polypeptide may be found in the cell culture medium. If however, the FGFR-L polypeptide is not secreted from the host cells, it will be present in the cytoplasm and/or the nucleus (for eukaryotic host cells) or in the
25 cytosol (for gram-negative bacteria host cells).

For an FGFR-L polypeptide situated in the host cell cytoplasm and/or nucleus (for eukaryotic host cells) or in the cytosol (for bacterial host cells), the intracellular material (including inclusion bodies for gram-negative bacteria) can be extracted from the host cell using any standard technique known to the skilled artisan. For example,
30 the host cells can be lysed to release the contents of the periplasm/cytoplasm by French press, homogenization, and/or sonication followed by centrifugation.

If an FGFR-L polypeptide has formed inclusion bodies in the cytosol, the inclusion bodies can often bind to the inner and/or outer cellular membranes and thus

will be found primarily in the pellet material after centrifugation. The pellet material can then be treated at pH extremes or with a chaotropic agent such as a detergent, guanidine, guanidine derivatives, urea, or urea derivatives in the presence of a reducing agent such as dithiothreitol at alkaline pH or tris carboxyethyl phosphine at acid pH to release, break apart, and solubilize the inclusion bodies. The solubilized FGFR-L polypeptide can then be analyzed using gel electrophoresis, immunoprecipitation, or the like. If it is desired to isolate the FGFR-L polypeptide, isolation may be accomplished using standard methods such as those described herein and in Marston *et al.*, 1990, *Meth. Enz.*, 182:264-75.

10 In some cases, an FGFR-L polypeptide may not be biologically active upon isolation. Various methods for "refolding" or converting the polypeptide to its tertiary structure and generating disulfide linkages can be used to restore biological activity. Such methods include exposing the solubilized polypeptide to a pH usually above 7 and in the presence of a particular concentration of a chaotrope. The selection of chaotrope is very similar to the choices used for inclusion body solubilization, but usually the chaotrope is used at a lower concentration and is not necessarily the same as chaotropes used for the solubilization. In most cases the refolding/oxidation solution will also contain a reducing agent or the reducing agent plus its oxidized form in a specific ratio to generate a particular redox potential allowing for disulfide shuffling to occur in the formation of the protein's cysteine bridges. Some of the commonly used redox couples include cysteine/cystamine, glutathione (GSH)/dithiobis GSH, cupric FGFR-Loride, dithiothreitol(DTT)/dithiane DTT, and 2-2-mercaptoethanol(bME)/dithio-b(ME). In many instances, a cosolvent may be used or may be needed to increase the efficiency of the refolding, and the more common reagents used for this purpose include glycerol, polyethylene glycol of various molecular weights, arginine and the like.

If inclusion bodies are not formed to a significant degree upon expression of an FGFR-L polypeptide, then the polypeptide will be found primarily in the supernatant after centrifugation of the cell homogenate. The polypeptide may be further isolated from the supernatant using methods such as those described herein.

The purification of an FGFR-L polypeptide from solution can be accomplished using a variety of techniques. If the polypeptide has been synthesized such that it contains a tag such as Hexahistidine (FGFR-L polypeptide/hexaHis) or

other small peptide such as FLAG (Eastman Kodak Co., New Haven, CT) or *myc* (Invitrogen, Carlsbad, CA) at either its carboxyl- or amino-terminus, it may be purified in a one-step process by passing the solution through an affinity column where the column matrix has a high affinity for the tag.

5 For example, polyhistidine binds with great affinity and specificity to nickel. Thus, an affinity column of nickel (such as the Qiagen[®] nickel columns) can be used for purification of FGFR-L polypeptide/polyHis. *See, e.g., Current Protocols in Molecular Biology* § 10.11.8 (Ausubel *et al.*, eds., Green Publishers Inc. and Wiley and Sons 1993).

10 Additionally, FGFR-L polypeptides may be purified through the use of a monoclonal antibody that is capable of specifically recognizing and binding to an FGFR-L polypeptide.

 Other suitable procedures for purification include, without limitation, affinity chromatography, immunoaffinity chromatography, ion exchange chromatography,
15 molecular sieve chromatography, HPLC, electrophoresis (including native gel electrophoresis) followed by gel elution, and preparative isoelectric focusing ("Isoprime" machine/technique, Hoefer Scientific, San Francisco, CA). In some cases, two or more purification techniques may be combined to achieve increased purity.

20 FGFR-L polypeptides may also be prepared by chemical synthesis methods (such as solid phase peptide synthesis) using techniques known in the art such as those set forth by Merrifield *et al.*, 1963, *J. Am. Chem. Soc.* 85:2149; Houghten *et al.*, 1985, *Proc Natl Acad. Sci. USA* 82:5132; and Stewart and Young, *Solid Phase Peptide Synthesis* (Pierce Chemical Co. 1984). Such polypeptides may be
25 synthesized with or without a methionine on the amino-terminus. Chemically synthesized FGFR-L polypeptides may be oxidized using methods set forth in these references to form disulfide bridges. Chemically synthesized FGFR-L polypeptides are expected to have comparable biological activity to the corresponding FGFR-L polypeptides produced recombinantly or purified from natural sources, and thus may
30 be used interchangeably with a recombinant or natural FGFR-L polypeptide.

 Another means of obtaining FGFR-L polypeptide is via purification from biological samples such as source tissues and/or fluids in which the FGFR-L polypeptide is naturally found. Such purification can be conducted using methods for

protein purification as described herein. The presence of the FGFR-L polypeptide during purification may be monitored, for example, using an antibody prepared against recombinantly produced FGFR-L polypeptide or peptide fragments thereof.

5 A number of additional methods for producing nucleic acids and polypeptides are known in the art, and the methods can be used to produce polypeptides having specificity for FGFR-L polypeptide. *See, e.g., Roberts et al., 1997, Proc. Natl. Acad. Sci. U.S.A. 94:12297-303*, which describes the production of fusion proteins between an mRNA and its encoded peptide. *See also, Roberts, 1999, Curr. Opin. Chem. Biol. 3:268-73*. Additionally, U.S. Patent No. 5,824,469 describes methods for obtaining
10 oligonucleotides capable of carrying out a specific biological function. The procedure involves generating a heterogeneous pool of oligonucleotides, each having a 5' randomized sequence, a central preselected sequence, and a 3' randomized sequence. The resulting heterogeneous pool is introduced into a population of cells that do not exhibit the desired biological function. Subpopulations of the cells are then screened
15 for those that exhibit a predetermined biological function. From that subpopulation, oligonucleotides capable of carrying out the desired biological function are isolated.

U.S. Patent Nos. 5,763,192; 5,814,476; 5,723,323; and 5,817,483 describe processes for producing peptides or polypeptides. This is done by producing stochastic genes or fragments thereof, and then introducing these genes into host cells
20 which produce one or more proteins encoded by the stochastic genes. The host cells are then screened to identify those clones producing peptides or polypeptides having the desired activity.

Another method for producing peptides or polypeptides is described in PCT/US98/20094 (WO99/15650) filed by Athersys, Inc. Known as "Random
25 Activation of Gene Expression for Gene Discovery" (RAGE-GD), the process involves the activation of endogenous gene expression or over-expression of a gene by *in situ* recombination methods. For example, expression of an endogenous gene is activated or increased by integrating a regulatory sequence into the target cell which is capable of activating expression of the gene by non-homologous or illegitimate
30 recombination. The target DNA is first subjected to radiation, and a genetic promoter inserted. The promoter eventually locates a break at the front of a gene, initiating transcription of the gene. This results in expression of the desired peptide or polypeptide.

5 It will be appreciated that these methods can also be used to create comprehensive FGFR-L polypeptide expression libraries, which can subsequently be used for high throughput phenotypic screening in a variety of assays, such as biochemical assays, cellular assays, and whole organism assays (*e.g.*, plant, mouse, etc.).

Synthesis

10 It will be appreciated by those skilled in the art that the nucleic acid and polypeptide molecules described herein may be produced by recombinant and other means.

Selective Binding Agents

15 The term "selective binding agent" refers to a molecule that has specificity for one or more FGFR-L polypeptides. Suitable selective binding agents include, but are not limited to, antibodies and derivatives thereof, polypeptides, and small molecules. Suitable selective binding agents may be prepared using methods known in the art. An exemplary FGFR-L polypeptide selective binding agent of the present invention is capable of binding a certain portion of the FGFR-L polypeptide thereby inhibiting the binding of the polypeptide to an FGFR-L polypeptide receptor.

20 Selective binding agents such as antibodies and antibody fragments that bind FGFR-L polypeptides are within the scope of the present invention. The antibodies may be polyclonal including monospecific polyclonal; monoclonal (MAbs); recombinant; chimeric; humanized, such as CDR-grafted; human; single chain; and/or bispecific; as well as fragments; variants; or derivatives thereof. Antibody fragments
25 include those portions of the antibody that bind to an epitope on the FGFR-L polypeptide. Examples of such fragments include Fab and F(ab') fragments generated by enzymatic cleavage of full-length antibodies. Other binding fragments include those generated by recombinant DNA techniques, such as the expression of recombinant plasmids containing nucleic acid sequences encoding antibody variable
30 regions.

Polyclonal antibodies directed toward an FGFR-L polypeptide generally are produced in animals (*e.g.*, rabbits or mice) by means of multiple subcutaneous or intraperitoneal injections of FGFR-L polypeptide and an adjuvant. It may be useful to

conjugate an FGFR-L polypeptide to a carrier protein that is immunogenic in the species to be immunized, such as keyhole limpet hemocyanin, serum, albumin, bovine thyroglobulin, or soybean trypsin inhibitor. Also, aggregating agents such as alum are used to enhance the immune response. After immunization, the animals are
5 bled and the serum is assayed for anti-FGFR-L antibody titer.

Monoclonal antibodies directed toward FGFR-L polypeptides are produced using any method that provides for the production of antibody molecules by continuous cell lines in culture. Examples of suitable methods for preparing monoclonal antibodies include the hybridoma methods of Kohler *et al.*, 1975, *Nature*
10 256:495-97 and the human B-cell hybridoma method (Kozbor, 1984, *J. Immunol.* 133:3001; Brodeur *et al.*, *Monoclonal Antibody Production Techniques and Applications* 51-63 (Marcel Dekker, Inc., 1987). Also provided by the invention are hybridoma cell lines that produce monoclonal antibodies reactive with FGFR-L polypeptides.

15 Monoclonal antibodies of the invention may be modified for use as therapeutics. One embodiment is a "chimeric" antibody in which a portion of the heavy (H) and/or light (L) chain is identical with or homologous to a corresponding sequence in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is/are identical with or
20 homologous to a corresponding sequence in antibodies derived from another species or belonging to another antibody class or subclass. Also included are fragments of such antibodies, so long as they exhibit the desired biological activity. See U.S. Patent No. 4,816,567; Morrison *et al.*, 1985, *Proc. Natl. Acad. Sci.* 81:6851-55.

In another embodiment, a monoclonal antibody of the invention is a
25 "humanized" antibody. Methods for humanizing non-human antibodies are well known in the art. See U.S. Patent Nos. 5,585,089 and 5,693,762. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source that is non-human. Humanization can be performed, for example, using methods described in the art (Jones *et al.*, 1986, *Nature* 321:522-25; Riechmann *et al.*,
30 1998, *Nature* 332:323-27; Verhoeyen *et al.*, 1988, *Science* 239:1534-36), by substituting at least a portion of a rodent complementarity-determining region (CDR) for the corresponding regions of a human antibody.

Also encompassed by the invention are human antibodies that bind FGFR-L polypeptides. Using transgenic animals (*e.g.*, mice) that are capable of producing a repertoire of human antibodies in the absence of endogenous immunoglobulin production such antibodies are produced by immunization with an FGFR-L polypeptide antigen (*i.e.*, having at least 6 contiguous amino acids), optionally conjugated to a carrier. *See, e.g.*, Jakobovits *et al.*, 1993, *Proc. Natl. Acad. Sci.* 90:2551-55; Jakobovits *et al.*, 1993, *Nature* 362:255-58; Bruggermann *et al.*, 1993, *Year in Immuno.* 7:33. In one method, such transgenic animals are produced by incapacitating the endogenous loci encoding the heavy and light immunoglobulin chains therein, and inserting loci encoding human heavy and light chain proteins into the genome thereof. Partially modified animals, that is those having less than the full complement of modifications, are then cross-bred to obtain an animal having all of the desired immune system modifications. When administered an immunogen, these transgenic animals produce antibodies with human (rather than, *e.g.*, murine) amino acid sequences, including variable regions which are immunospecific for these antigens. *See* PCT App. Nos. PCT/US96/05928 and PCT/US93/06926. Additional methods are described in U.S. Patent No. 5,545,807, PCT App. Nos. PCT/US91/245 and PCT/GB89/01207, and in European Patent Nos. 546073B1 and 546073A1. Human antibodies can also be produced by the expression of recombinant DNA in host cells or by expression in hybridoma cells as described herein.

In an alternative embodiment, human antibodies can also be produced from phage-display libraries (Hoogenboom *et al.*, 1991, *J. Mol. Biol.* 227:381; Marks *et al.*, 1991, *J. Mol. Biol.* 222:581). These processes mimic immune selection through the display of antibody repertoires on the surface of filamentous bacteriophage, and subsequent selection of phage by their binding to an antigen of choice. One such technique is described in PCT App. No. PCT/US98/17364, which describes the isolation of high affinity and functional agonistic antibodies for MPL- and msk-receptors using such an approach.

Chimeric, CDR grafted, and humanized antibodies are typically produced by recombinant methods. Nucleic acids encoding the antibodies are introduced into host cells and expressed using materials and procedures described herein. In a preferred embodiment, the antibodies are produced in mammalian host cells, such as CHO cells. Monoclonal (*e.g.*, human) antibodies may be produced by the expression of

recombinant DNA in host cells or by expression in hybridoma cells as described herein.

5 The anti-FGFR-L antibodies of the invention may be employed in any known assay method, such as competitive binding assays, direct and indirect sandwich assays, and immunoprecipitation assays (Sola, *Monoclonal Antibodies: A Manual of Techniques* 147-158 (CRC Press, Inc., 1987)) for the detection and quantitation of FGFR-L polypeptides. The antibodies will bind FGFR-L polypeptides with an affinity that is appropriate for the assay method being employed.

10 For diagnostic applications, in certain embodiments, anti-FGFR-L antibodies may be labeled with a detectable moiety. The detectable moiety can be any one that is capable of producing, either directly or indirectly, a detectable signal. For example, the detectable moiety may be a radioisotope, such as ^3H , ^{14}C , ^{32}P , ^{35}S , ^{125}I , ^{99}Tc , ^{111}In , or ^{67}Ga ; a fluorescent or chemiluminescent compound, such as fluorescein isothiocyanate, rhodamine, or luciferin; or an enzyme, such as alkaline phosphatase, β -galactosidase, or horseradish peroxidase (Bayer, *et al.*, 1990, *Meth. Enz.* 184:138-15 63).

Competitive binding assays rely on the ability of a labeled standard (*e.g.*, an FGFR-L polypeptide, or an immunologically reactive portion thereof) to compete with the test sample analyte (an FGFR-L polypeptide) for binding with a limited 20 amount of anti-FGFR-L antibody. The amount of an FGFR-L polypeptide in the test sample is inversely proportional to the amount of standard that becomes bound to the antibodies. To facilitate determining the amount of standard that becomes bound, the antibodies typically are insolubilized before or after the competition, so that the standard and analyte that are bound to the antibodies may conveniently be separated 25 from the standard and analyte which remain unbound.

Sandwich assays typically involve the use of two antibodies, each capable of binding to a different immunogenic portion, or epitope, of the protein to be detected and/or quantitated. In a sandwich assay, the test sample analyte is typically bound by a first antibody which is immobilized on a solid support, and thereafter a second 30 antibody binds to the analyte, thus forming an insoluble three-part complex. *See, e.g.*, U.S. Patent No. 4,376,110. The second antibody may itself be labeled with a detectable moiety (direct sandwich assays) or may be measured using an anti-immunoglobulin antibody that is labeled with a detectable moiety (indirect sandwich

assays). For example, one type of sandwich assay is an enzyme-linked immunosorbent assay (ELISA), in which case the detectable moiety is an enzyme.

The selective binding agents, including anti-FGFR-L antibodies, are also useful for *in vivo* imaging. An antibody labeled with a detectable moiety may be administered to an animal, preferably into the bloodstream, and the presence and location of the labeled antibody in the host assayed. The antibody may be labeled with any moiety that is detectable in an animal, whether by nuclear magnetic resonance, radiology, or other detection means known in the art.

Selective binding agents of the invention, including antibodies, may be used as therapeutics. These therapeutic agents are generally agonists or antagonists, in that they either enhance or reduce, respectively, at least one of the biological activities of an FGFR-L polypeptide. In one embodiment, antagonist antibodies of the invention are antibodies or binding fragments thereof which are capable of specifically binding to an FGFR-L polypeptide and which are capable of inhibiting or eliminating the functional activity of an FGFR-L polypeptide *in vivo* or *in vitro*. In preferred embodiments, the selective binding agent, *e.g.*, an antagonist antibody, will inhibit the functional activity of an FGFR-L polypeptide by at least about 50%, and preferably by at least about 80%. In another embodiment, the selective binding agent may be an anti-FGFR-L polypeptide antibody that is capable of interacting with an FGFR-L polypeptide binding partner (a ligand or receptor) thereby inhibiting or eliminating FGFR-L polypeptide activity *in vitro* or *in vivo*. Selective binding agents, including agonist and antagonist anti-FGFR-L polypeptide antibodies, are identified by screening assays that are well known in the art.

The invention also relates to a kit comprising FGFR-L selective binding agents (such as antibodies) and other reagents useful for detecting FGFR-L polypeptide levels in biological samples. Such reagents may include a detectable label, blocking serum, positive and negative control samples, and detection reagents.

Microarrays

It will be appreciated that DNA microarray technology can be utilized in accordance with the present invention. DNA microarrays are miniature, high-density arrays of nucleic acids positioned on a solid support, such as glass. Each cell or element within the array contains numerous copies of a single nucleic acid species

that acts as a target for hybridization with a complementary nucleic acid sequence (e.g., mRNA). In expression profiling using DNA microarray technology, mRNA is first extracted from a cell or tissue sample and then converted enzymatically to fluorescently labeled cDNA. This material is hybridized to the microarray and unbound cDNA is removed by washing. The expression of discrete genes represented on the array is then visualized by quantitating the amount of labeled cDNA that is specifically bound to each target nucleic acid molecule. In this way, the expression of thousands of genes can be quantitated in a high throughput, parallel manner from a single sample of biological material.

This high throughput expression profiling has a broad range of applications with respect to the FGFR-L molecules of the invention, including, but not limited to: the identification and validation of FGFR-L disease-related genes as targets for therapeutics; molecular toxicology of related FGFR-L molecules and inhibitors thereof; stratification of populations and generation of surrogate markers for clinical trials; and enhancing related FGFR-L polypeptide small molecule drug discovery by aiding in the identification of selective compounds in high throughput screens.

Chemical Derivatives

Chemically modified derivatives of FGFR-L polypeptides may be prepared by one skilled in the art, given the disclosures described herein. FGFR-L polypeptide derivatives are modified in a manner that is different – either in the type or location of the molecules naturally attached to the polypeptide. Derivatives may include molecules formed by the deletion of one or more naturally-attached chemical groups. The polypeptide comprising the amino acid sequence of either SEQ ID NO: 2 or SEQ ID NO: 5, or other FGFR-L polypeptide, may be modified by the covalent attachment of one or more polymers. For example, the polymer selected is typically water-soluble so that the protein to which it is attached does not precipitate in an aqueous environment, such as a physiological environment. Included within the scope of suitable polymers is a mixture of polymers. Preferably, for therapeutic use of the end-product preparation, the polymer will be pharmaceutically acceptable.

The polymers each may be of any molecular weight and may be branched or unbranched. The polymers each typically have an average molecular weight of between about 2 kDa to about 100 kDa (the term “about” indicating that in

preparations of a water-soluble polymer, some molecules will weigh more, some less, than the stated molecular weight). The average molecular weight of each polymer is preferably between about 5 kDa and about 50 kDa, more preferably between about 12 kDa and about 40 kDa and most preferably between about 20 kDa and about 35 kDa.

5 Suitable water-soluble polymers or mixtures thereof include, but are not limited to, N-linked or O-linked carbohydrates, sugars, phosphates, polyethylene glycol (PEG) (including the forms of PEG that have been used to derivatize proteins, including mono-(C₁-C₁₀), alkoxy-, or aryloxy-polyethylene glycol), monomethoxy-polyethylene glycol, dextran (such as low molecular weight dextran of, for example,
10 about 6 kD), cellulose, or other carbohydrate based polymers, poly-(N-vinyl pyrrolidone) polyethylene glycol, propylene glycol homopolymers, polypropylene oxide/ethylene oxide co-polymers, polyoxyethylated polyols (e.g., glycerol), and polyvinyl alcohol. Also encompassed by the present invention are bifunctional crosslinking molecules which may be used to prepare covalently attached FGFR-L
15 polypeptide multimers.

 In general, chemical derivatization may be performed under any suitable condition used to react a protein with an activated polymer molecule. Methods for preparing chemical derivatives of polypeptides will generally comprise the steps of: (a) reacting the polypeptide with the activated polymer molecule (such as a reactive
20 ester or aldehyde derivative of the polymer molecule) under conditions whereby the polypeptide comprising the amino acid sequence of either SEQ ID NO: 2 or SEQ ID NO: 5, or other FGFR-L polypeptide, becomes attached to one or more polymer molecules, and (b) obtaining the reaction products. The optimal reaction conditions will be determined based on known parameters and the desired result. For example,
25 the larger the ratio of polymer molecules to protein, the greater the percentage of attached polymer molecule. In one embodiment, the FGFR-L polypeptide derivative may have a single polymer molecule moiety at the amino-terminus. See, e.g., U.S. Patent No. 5,234,784.

 The pegylation of a polypeptide may be specifically carried out using any of
30 the pegylation reactions known in the art. Such reactions are described, for example, in the following references: Francis *et al.*, 1992, *Focus on Growth Factors* 3:4-10; European Patent Nos. 0154316 and 0401384; and U.S. Patent No. 4,179,337. For example, pegylation may be carried out via an acylation reaction or an alkylation

reaction with a reactive polyethylene glycol molecule (or an analogous reactive water-soluble polymer) as described herein. For the acylation reactions, a selected polymer should have a single reactive ester group. For reductive alkylation, a selected polymer should have a single reactive aldehyde group. A reactive aldehyde is, for example, polyethylene glycol propionaldehyde, which is water stable, or mono C₁-C₁₀ alkoxy or aryloxy derivatives thereof (see U.S. Patent No. 5,252,714).

In another embodiment, FGFR-L polypeptides may be chemically coupled to biotin. The biotin/FGFR-L polypeptide molecules are then allowed to bind to avidin, resulting in tetravalent avidin/biotin/FGFR-L polypeptide molecules. FGFR-L polypeptides may also be covalently coupled to dinitrophenol (DNP) or trinitrophenol (TNP) and the resulting conjugates precipitated with anti-DNP or anti-TNP-IgM to form decameric conjugates with a valency of 10.

Generally, conditions that may be alleviated or modulated by the administration of the present FGFR-L polypeptide derivatives include those described herein for FGFR-L polypeptides. However, the FGFR-L polypeptide derivatives disclosed herein may have additional activities, enhanced or reduced biological activity, or other characteristics, such as increased or decreased half-life, as compared to the non-derivatized molecules.

Genetically Engineered Non-Human Animals

Additionally included within the scope of the present invention are non-human animals such as mice, rats, or other rodents; rabbits, goats, sheep, or other farm animals, in which the genes encoding native FGFR-L polypeptide have been disrupted (*i.e.*, "knocked out") such that the level of expression of FGFR-L polypeptide is significantly decreased or completely abolished. Such animals may be prepared using techniques and methods such as those described in U.S. Patent No. 5,557,032.

The present invention further includes non-human animals such as mice, rats, or other rodents; rabbits, goats, sheep, or other farm animals, in which either the native form of an FGFR-L gene for that animal or a heterologous FGFR-L gene is over-expressed by the animal, thereby creating a "transgenic" animal. Such transgenic animals may be prepared using well known methods such as those described in U.S. Patent No 5,489,743 and PCT Pub. No. WO 94/28122.

The present invention further includes non-human animals in which the promoter for one or more of the FGFR-L polypeptides of the present invention is either activated or inactivated (*e.g.*, by using homologous recombination methods) to alter the level of expression of one or more of the native FGFR-L polypeptides.

5 These non-human animals may be used for drug candidate screening. In such screening, the impact of a drug candidate on the animal may be measured. For example, drug candidates may decrease or increase the expression of the FGFR-L gene. In certain embodiments, the amount of FGFR-L polypeptide that is produced may be measured after the exposure of the animal to the drug candidate.
10 Additionally, in certain embodiments, one may detect the actual impact of the drug candidate on the animal. For example, over-expression of a particular gene may result in, or be associated with, a disease or pathological condition. In such cases, one may test a drug candidate's ability to decrease expression of the gene or its ability to prevent or inhibit a pathological condition. In other examples, the production of a
15 particular metabolic product such as a fragment of a polypeptide, may result in, or be associated with, a disease or pathological condition. In such cases, one may test a drug candidate's ability to decrease the production of such a metabolic product or its ability to prevent or inhibit a pathological condition.

20 Assaying for Other Modulators of FGFR-L Polypeptide Activity

 In some situations, it may be desirable to identify molecules that are modulators, *i.e.*, agonists or antagonists, of the activity of FGFR-L polypeptide. Natural or synthetic molecules that modulate FGFR-L polypeptide may be identified using one or more screening assays, such as those described herein. Such molecules
25 may be administered either in an *ex vivo* manner or in an *in vivo* manner by injection, or by oral delivery, implantation device, or the like.

 "Test molecule" refers to a molecule that is under evaluation for the ability to modulate (*i.e.*, increase or decrease) the activity of an FGFR-L polypeptide. Most commonly, a test molecule will interact directly with an FGFR-L polypeptide.
30 However, it is also contemplated that a test molecule may also modulate FGFR-L polypeptide activity indirectly, such as by affecting FGFR-L gene expression, or by binding to an FGFR-L polypeptide binding partner (*e.g.*, receptor or ligand). In one embodiment, a test molecule will bind to an FGFR-L polypeptide with an affinity

constant of at least about 10^{-6} M, preferably about 10^{-8} M, more preferably about 10^{-9} M, and even more preferably about 10^{-10} M.

Methods for identifying compounds that interact with FGFR-L polypeptides are encompassed by the present invention. In certain embodiments, an FGFR-L polypeptide is incubated with a test molecule under conditions that permit the interaction of the test molecule with an FGFR-L polypeptide, and the extent of the interaction is measured. The test molecule can be screened in a substantially purified form or in a crude mixture.

In certain embodiments, an FGFR-L polypeptide agonist or antagonist may be a protein, peptide, carbohydrate, lipid, or small molecular weight molecule that interacts with FGFR-L polypeptide to regulate its activity. Molecules which regulate FGFR-L polypeptide expression include nucleic acids which are complementary to nucleic acids encoding an FGFR-L polypeptide, or are complementary to nucleic acids sequences which direct or control the expression of FGFR-L polypeptide, and which act as anti-sense regulators of expression.

Once a test molecule has been identified as interacting with an FGFR-L polypeptide, the molecule may be further evaluated for its ability to increase or decrease FGFR-L polypeptide activity. The measurement of the interaction of a test molecule with FGFR-L polypeptide may be carried out in several formats, including cell-based binding assays, membrane binding assays, solution-phase assays, and immunoassays. In general, a test molecule is incubated with an FGFR-L polypeptide for a specified period of time, and FGFR-L polypeptide activity is determined by one or more assays for measuring biological activity.

The interaction of test molecules with FGFR-L polypeptides may also be assayed directly using polyclonal or monoclonal antibodies in an immunoassay. Alternatively, modified forms of FGFR-L polypeptides containing epitope tags as described herein may be used in solution and immunoassays.

In the event that FGFR-L polypeptides display biological activity through an interaction with a binding partner (e.g., a receptor or a ligand), a variety of *in vitro* assays may be used to measure the binding of an FGFR-L polypeptide to the corresponding binding partner (such as a selective binding agent, receptor, or ligand). These assays may be used to screen test molecules for their ability to increase or decrease the rate and/or the extent of binding of an FGFR-L polypeptide to its binding

partner. In one assay, an FGFR-L polypeptide is immobilized in the wells of a microtiter plate. Radiolabeled FGFR-L polypeptide binding partner (for example, iodinated FGFR-L polypeptide binding partner) and a test molecule can then be added either one at a time (in either order) or simultaneously to the wells. After incubation, the wells can be washed and counted for radioactivity, using a scintillation counter, to determine the extent to which the binding partner bound to the FGFR-L polypeptide. Typically, a molecule will be tested over a range of concentrations, and a series of control wells lacking one or more elements of the test assays can be used for accuracy in the evaluation of the results. An alternative to this method involves reversing the “positions” of the proteins, *i.e.*, immobilizing FGFR-L polypeptide binding partner to the microtiter plate wells, incubating with the test molecule and radiolabeled FGFR-L polypeptide, and determining the extent of FGFR-L polypeptide binding. *See, e.g., Current Protocols in Molecular Biology*, chap. 18 (Ausubel *et al.*, eds., Green Publishers Inc. and Wiley and Sons 1995).

As an alternative to radiolabeling, an FGFR-L polypeptide or its binding partner may be conjugated to biotin, and the presence of biotinylated protein can then be detected using streptavidin linked to an enzyme, such as horse radish peroxidase (HRP) or alkaline phosphatase (AP), which can be detected colorometrically, or by fluorescent tagging of streptavidin. An antibody directed to an FGFR-L polypeptide or to an FGFR-L polypeptide binding partner, and which is conjugated to biotin, may also be used for purposes of detection following incubation of the complex with enzyme-linked streptavidin linked to AP or HRP.

A FGFR-L polypeptide or an FGFR-L polypeptide binding partner can also be immobilized by attachment to agarose beads, acrylic beads, or other types of such inert solid phase substrates. The substrate-protein complex can be placed in a solution containing the complementary protein and the test compound. After incubation, the beads can be precipitated by centrifugation, and the amount of binding between an FGFR-L polypeptide and its binding partner can be assessed using the methods described herein. Alternatively, the substrate-protein complex can be immobilized in a column with the test molecule and complementary protein passing through the column. The formation of a complex between an FGFR-L polypeptide and its binding partner can then be assessed using any of the techniques described herein (*e.g.*, radiolabelling or antibody binding).

Another *in vitro* assay that is useful for identifying a test molecule which increases or decreases the formation of a complex between an FGFR-L polypeptide binding protein and an FGFR-L polypeptide binding partner is a surface plasmon resonance detector system such as the BIAcore assay system (Pharmacia, Piscataway, NJ). The BIAcore system is utilized as specified by the manufacturer. This assay essentially involves the covalent binding of either FGFR-L polypeptide or an FGFR-L polypeptide binding partner to a dextran-coated sensor chip that is located in a detector. The test compound and the other complementary protein can then be injected, either simultaneously or sequentially, into the chamber containing the sensor chip. The amount of complementary protein that binds can be assessed based on the change in molecular mass that is physically associated with the dextran-coated side of the sensor chip, with the change in molecular mass being measured by the detector system.

In some cases, it may be desirable to evaluate two or more test compounds together for their ability to increase or decrease the formation of a complex between an FGFR-L polypeptide and an FGFR-L polypeptide binding partner. In these cases, the assays set forth herein can be readily modified by adding such additional test compound(s) either simultaneously with, or subsequent to, the first test compound. The remainder of the steps in the assay are as set forth herein.

In vitro assays such as those described herein may be used advantageously to screen large numbers of compounds for an effect on the formation of a complex between an FGFR-L polypeptide and FGFR-L polypeptide binding partner. The assays may be automated to screen compounds generated in phage display, synthetic peptide, and chemical synthesis libraries.

Compounds which increase or decrease the formation of a complex between an FGFR-L polypeptide and an FGFR-L polypeptide binding partner may also be screened in cell culture using cells and cell lines expressing either FGFR-L polypeptide or FGFR-L polypeptide binding partner. Cells and cell lines may be obtained from any mammal, but preferably will be from human or other primate, canine, or rodent sources. The binding of an FGFR-L polypeptide to cells expressing FGFR-L polypeptide binding partner at the surface is evaluated in the presence or absence of test molecules, and the extent of binding may be determined by, for example, flow cytometry using a biotinylated antibody to an FGFR-L polypeptide

binding partner. Cell culture assays can be used advantageously to further evaluate compounds that score positive in protein binding assays described herein.

Cell cultures can also be used to screen the impact of a drug candidate. For example, drug candidates may decrease or increase the expression of the FGFR-L gene. In certain embodiments, the amount of FGFR-L polypeptide or an FGFR-L polypeptide fragment that is produced may be measured after exposure of the cell culture to the drug candidate. In certain embodiments, one may detect the actual impact of the drug candidate on the cell culture. For example, the over-expression of a particular gene may have a particular impact on the cell culture. In such cases, one may test a drug candidate's ability to increase or decrease the expression of the gene or its ability to prevent or inhibit a particular impact on the cell culture. In other examples, the production of a particular metabolic product such as a fragment of a polypeptide, may result in, or be associated with, a disease or pathological condition. In such cases, one may test a drug candidate's ability to decrease the production of such a metabolic product in a cell culture.

Internalizing Proteins

The *tat* protein sequence (from HIV) can be used to internalize proteins into a cell. See, e.g., Falwell *et al.*, 1994, *Proc. Natl. Acad. Sci. U.S.A.* 91:664-68. For example, an 11 amino acid sequence (Y-G-R-K-K-R-R-Q-R-R-R; SEQ ID NO: 9) of the HIV *tat* protein (termed the "protein transduction domain," or TAT PDT) has been described as mediating delivery across the cytoplasmic membrane and the nuclear membrane of a cell. See Schwarze *et al.*, 1999, *Science* 285:1569-72; and Nagahara *et al.*, 1998, *Nat. Med.* 4:1449-52. In these procedures, FITC-constructs (FITC-labeled G-G-G-G-Y-G-R-K-K-R-R-Q-R-R-R; SEQ ID NO: 10), which penetrate tissues following intraperitoneal administration, are prepared, and the binding of such constructs to cells is detected by fluorescence-activated cell sorting (FACS) analysis. Cells treated with a *tat*- β -gal fusion protein will demonstrate β -gal activity. Following injection, expression of such a construct can be detected in a number of tissues, including liver, kidney, lung, heart, and brain tissue. It is believed that such constructs undergo some degree of unfolding in order to enter the cell, and as such, may require a refolding following entry into the cell.

It will thus be appreciated that the *tat* protein sequence may be used to

internalize a desired polypeptide into a cell. For example, using the *tat* protein sequence, an FGFR-L antagonist (such as an anti-FGFR-L selective binding agent, small molecule, soluble receptor, or antisense oligonucleotide) can be administered intracellularly to inhibit the activity of an FGFR-L molecule. As used herein, the term
5 "FGFR-L molecule" refers to both FGFR-L nucleic acid molecules and FGFR-L polypeptides as defined herein. Where desired, the FGFR-L protein itself may also be internally administered to a cell using these procedures. *See also*, Straus, 1999, *Science* 285:1466-67.

10 Cell Source Identification Using FGFR-L Polypeptide

In accordance with certain embodiments of the invention, it may be useful to be able to determine the source of a certain cell type associated with an FGFR-L polypeptide. For example, it may be useful to determine the origin of a disease or pathological condition as an aid in selecting an appropriate therapy. In certain
15 embodiments, nucleic acids encoding an FGFR-L polypeptide can be used as a probe to identify cells described herein by screening the nucleic acids of the cells with such a probe. In other embodiments, one may use anti-FGFR-L polypeptide antibodies to test for the presence of FGFR-L polypeptide in cells, and thus, determine if such cells are of the types described herein.

20 FGFR-L Polypeptide Compositions and Administration

Therapeutic compositions are within the scope of the present invention. Such FGFR-L polypeptide pharmaceutical compositions may comprise a therapeutically effective amount of an FGFR-L polypeptide or an FGFR-L nucleic acid molecule in
25 admixture with a pharmaceutically or physiologically acceptable formulation agent selected for suitability with the mode of administration. Pharmaceutical compositions may comprise a therapeutically effective amount of one or more FGFR-L polypeptide selective binding agents in admixture with a pharmaceutically or physiologically acceptable formulation agent selected for suitability with the mode of administration.

30 Acceptable formulation materials preferably are nontoxic to recipients at the dosages and concentrations employed.

The pharmaceutical composition may contain formulation materials for modifying, maintaining, or preserving, for example, the pH, osmolarity, viscosity,

clarity, color, isotonicity, odor, sterility, stability, rate of dissolution or release, adsorption, or penetration of the composition. Suitable formulation materials include, but are not limited to, amino acids (such as glycine, glutamine, asparagine, arginine, or lysine), antimicrobials, antioxidants (such as ascorbic acid, sodium sulfite, or sodium hydrogen-sulfite), buffers (such as borate, bicarbonate, Tris-HCl, citrates, phosphates, or other organic acids), bulking agents (such as mannitol or glycine), chelating agents (such as ethylenediamine tetraacetic acid (EDTA)), complexing agents (such as caffeine, polyvinylpyrrolidone, beta-cyclodextrin, or hydroxypropyl-beta-cyclodextrin), fillers, monosaccharides, disaccharides, and other carbohydrates (such as glucose, mannose, or dextrans), proteins (such as serum albumin, gelatin, or immunoglobulins), coloring, flavoring and diluting agents, emulsifying agents, hydrophilic polymers (such as polyvinylpyrrolidone), low molecular weight polypeptides, salt-forming counterions (such as sodium), preservatives (such as benzalkonium FGFR-Loride, benzoic acid, salicylic acid, thimerosal, phenethyl alcohol, methylparaben, propylparaben, FGFR-Lorhexidine, sorbic acid, or hydrogen peroxide), solvents (such as glycerin, propylene glycol, or polyethylene glycol), sugar alcohols (such as mannitol or sorbitol), suspending agents, surfactants or wetting agents (such as pluronics; PEG; sorbitan esters; polysorbates such as polysorbate 20 or polysorbate 80; triton; tromethamine; lecithin; cholesterol or tyloxapal), stability enhancing agents (such as sucrose or sorbitol), tonicity enhancing agents (such as alkali metal halides – preferably sodium or potassium FGFR-Loride – or mannitol sorbitol), delivery vehicles, diluents, excipients and/or pharmaceutical adjuvants. *See Remington's Pharmaceutical Sciences* (18th Ed., A.R. Gennaro, ed., Mack Publishing Company 1990).

The optimal pharmaceutical composition will be determined by a skilled artisan depending upon, for example, the intended route of administration, delivery format, and desired dosage. *See, e.g., Remington's Pharmaceutical Sciences, supra.* Such compositions may influence the physical state, stability, rate of *in vivo* release, and rate of *in vivo* clearance of the FGFR-L molecule.

The primary vehicle or carrier in a pharmaceutical composition may be either aqueous or non-aqueous in nature. For example, a suitable vehicle or carrier for injection may be water, physiological saline solution, or artificial cerebrospinal fluid, possibly supplemented with other materials common in compositions for parenteral

administration. Neutral buffered saline or saline mixed with serum albumin are further exemplary vehicles. Other exemplary pharmaceutical compositions comprise Tris buffer of about pH 7.0-8.5, or acetate buffer of about pH 4.0-5.5, which may further include sorbitol or a suitable substitute. In one embodiment of the present invention, FGFR-L polypeptide compositions may be prepared for storage by mixing the selected composition having the desired degree of purity with optional formulation agents (*Remington's Pharmaceutical Sciences, supra*) in the form of a lyophilized cake or an aqueous solution. Further, the FGFR-L polypeptide product may be formulated as a lyophilizate using appropriate excipients such as sucrose.

10 The FGFR-L polypeptide pharmaceutical compositions can be selected for parenteral delivery. Alternatively, the compositions may be selected for inhalation or for delivery through the digestive tract, such as orally. The preparation of such pharmaceutically acceptable compositions is within the skill of the art.

15 The formulation components are present in concentrations that are acceptable to the site of administration. For example, buffers are used to maintain the composition at physiological pH or at a slightly lower pH, typically within a pH range of from about 5 to about 8.

20 When parenteral administration is contemplated, the therapeutic compositions for use in this invention may be in the form of a pyrogen-free, parenterally acceptable, aqueous solution comprising the desired FGFR-L molecule in a pharmaceutically acceptable vehicle. A particularly suitable vehicle for parenteral injection is sterile distilled water in which an FGFR-L molecule is formulated as a sterile, isotonic solution, properly preserved. Yet another preparation can involve the formulation of the desired molecule with an agent, such as injectable microspheres, bio-erodible particles, polymeric compounds (such as polylactic acid or polyglycolic acid), beads, or liposomes, that provides for the controlled or sustained release of the product which may then be delivered via a depot injection. Hyaluronic acid may also be used, and this may have the effect of promoting sustained duration in the circulation. Other suitable means for the introduction of the desired molecule include implantable drug delivery devices.

30 In one embodiment, a pharmaceutical composition may be formulated for inhalation. For example, FGFR-L polypeptide may be formulated as a dry powder for inhalation. FGFR-L polypeptide or nucleic acid molecule inhalation solutions may

also be formulated with a propellant for aerosol delivery. In yet another embodiment, solutions may be nebulized. Pulmonary administration is further described in PCT Pub. No. WO 94/20069, which describes the pulmonary delivery of chemically modified proteins.

5 It is also contemplated that certain formulations may be administered orally. In one embodiment of the present invention, FGFR-L polypeptides that are administered in this fashion can be formulated with or without those carriers customarily used in the compounding of solid dosage forms such as tablets and capsules. For example, a capsule may be designed to release the active portion of the
10 formulation at the point in the gastrointestinal tract when bioavailability is maximized and pre-systemic degradation is minimized. Additional agents can be included to facilitate absorption of the FGFR-L polypeptide. Diluents, flavorings, low melting point waxes, vegetable oils, lubricants, suspending agents, tablet disintegrating agents, and binders may also be employed.

15 Another pharmaceutical composition may involve an effective quantity of FGFR-L polypeptides in a mixture with non-toxic excipients that are suitable for the manufacture of tablets. By dissolving the tablets in sterile water, or another appropriate vehicle, solutions can be prepared in unit-dose form. Suitable excipients include, but are not limited to, inert diluents, such as calcium carbonate, sodium
20 carbonate or bicarbonate, lactose, or calcium phosphate; or binding agents, such as starch, gelatin, or acacia; or lubricating agents such as magnesium stearate, stearic acid, or talc.

 Additional FGFR-L polypeptide pharmaceutical compositions will be evident to those skilled in the art, including formulations involving FGFR-L polypeptides in
25 sustained- or controlled-delivery formulations. Techniques for formulating a variety of other sustained- or controlled-delivery means, such as liposome carriers, bio-erodible microparticles or porous beads and depot injections, are also known to those skilled in the art. *See, e.g.*, PCT/US93/00829, which describes the controlled release of porous polymeric microparticles for the delivery of pharmaceutical compositions.

30 Additional examples of sustained-release preparations include semipermeable polymer matrices in the form of shaped articles, *e.g.* films, or microcapsules. Sustained release matrices may include polyesters, hydrogels, polylactides (U.S. Patent No. 3,773,919 and European Patent No. 058481), copolymers of L-glutamic

acid and gamma ethyl-L-glutamate (Sidman *et al.*, 1983, *Biopolymers* 22:547-56), poly(2-hydroxyethyl-methacrylate) (Langer *et al.*, 1981, *J. Biomed. Mater. Res.* 15:167-277 and Langer, 1982, *Chem. Tech.* 12:98-105), ethylene vinyl acetate (Langer *et al.*, *supra*) or poly-D(-)-3-hydroxybutyric acid (European Patent No. 133988). Sustained-release compositions may also include liposomes, which can be prepared by any of several methods known in the art. See, *e.g.*, Eppstein *et al.*, 1985, *Proc. Natl. Acad. Sci. USA* 82:3688-92; and European Patent Nos. 036676, 088046, and 143949.

The FGFR-L pharmaceutical composition to be used for *in vivo* administration typically must be sterile. This may be accomplished by filtration through sterile filtration membranes. Where the composition is lyophilized, sterilization using this method may be conducted either prior to, or following, lyophilization and reconstitution. The composition for parenteral administration may be stored in lyophilized form or in a solution. In addition, parenteral compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

Once the pharmaceutical composition has been formulated, it may be stored in sterile vials as a solution, suspension, gel, emulsion, solid, or as a dehydrated or lyophilized powder. Such formulations may be stored either in a ready-to-use form or in a form (*e.g.*, lyophilized) requiring reconstitution prior to administration.

In a specific embodiment, the present invention is directed to kits for producing a single-dose administration unit. The kits may each contain both a first container having a dried protein and a second container having an aqueous formulation. Also included within the scope of this invention are kits containing single and multi-chambered pre-filled syringes (*e.g.*, liquid syringes and lyosyringes).

The effective amount of an FGFR-L pharmaceutical composition to be employed therapeutically will depend, for example, upon the therapeutic context and objectives. One skilled in the art will appreciate that the appropriate dosage levels for treatment will thus vary depending, in part, upon the molecule delivered, the indication for which the FGFR-L molecule is being used, the route of administration, and the size (body weight, body surface, or organ size) and condition (the age and general health) of the patient. Accordingly, the clinician may titer the dosage and modify the route of administration to obtain the optimal therapeutic effect. A typical

dosage may range from about 0.1 $\mu\text{g/kg}$ to up to about 100 mg/kg or more, depending on the factors mentioned above. In other embodiments, the dosage may range from 0.1 $\mu\text{g/kg}$ up to about 100 mg/kg; or 1 $\mu\text{g/kg}$ up to about 100 mg/kg; or 5 $\mu\text{g/kg}$ up to about 100 mg/kg.

5 The frequency of dosing will depend upon the pharmacokinetic parameters of the FGFR-L molecule in the formulation being used. Typically, a clinician will administer the composition until a dosage is reached that achieves the desired effect. The composition may therefore be administered as a single dose, as two or more doses (which may or may not contain the same amount of the desired molecule) over
10 time, or as a continuous infusion via an implantation device or catheter. Further refinement of the appropriate dosage is routinely made by those of ordinary skill in the art and is within the ambit of tasks routinely performed by them. Appropriate dosages may be ascertained through use of appropriate dose-response data.

 The route of administration of the pharmaceutical composition is in accord
15 with known methods, *e.g.*, orally; through injection by intravenous, intraperitoneal, intracerebral (intraparenchymal), intracerebroventricular, intramuscular, intraocular, intraarterial, intraportal, or intralesional routes; by sustained release systems; or by implantation devices. Where desired, the compositions may be administered by bolus injection or continuously by infusion, or by implantation device.

20 Alternatively or additionally, the composition may be administered locally via implantation of a membrane, sponge, or other appropriate material onto which the desired molecule has been absorbed or encapsulated. Where an implantation device is used, the device may be implanted into any suitable tissue or organ, and delivery of the desired molecule may be via diffusion, timed-release bolus, or continuous
25 administration.

 In some cases, it may be desirable to use FGFR-L polypeptide pharmaceutical compositions in an *ex vivo* manner. In such instances, cells, tissues, or organs that have been removed from the patient are exposed to FGFR-L polypeptide pharmaceutical compositions after which the cells, tissues, or organs are subsequently
30 implanted back into the patient.

 In other cases, an FGFR-L polypeptide can be delivered by implanting certain cells that have been genetically engineered, using methods such as those described herein, to express and secrete the FGFR-L polypeptide. Such cells may be animal or

human cells, and may be autologous, heterologous, or xenogeneic. Optionally, the cells may be immortalized. In order to decrease the chance of an immunological response, the cells may be encapsulated to avoid infiltration of surrounding tissues. The encapsulation materials are typically biocompatible, semi-permeable polymeric enclosures or membranes that allow the release of the protein product(s) but prevent the destruction of the cells by the patient's immune system or by other detrimental factors from the surrounding tissues.

As discussed herein, it may be desirable to treat isolated cell populations (such as stem cells, lymphocytes, red blood cells, chondrocytes, neurons, and the like) with one or more FGFR-L polypeptides. This can be accomplished by exposing the isolated cells to the polypeptide directly, where it is in a form that is permeable to the cell membrane.

Additional embodiments of the present invention relate to cells and methods (e.g., homologous recombination and/or other recombinant production methods) for both the *in vitro* production of therapeutic polypeptides and for the production and delivery of therapeutic polypeptides by gene therapy or cell therapy. Homologous and other recombination methods may be used to modify a cell that contains a normally transcriptionally-silent FGFR-L gene, or an under-expressed gene, and thereby produce a cell which expresses therapeutically efficacious amounts of FGFR-L polypeptides.

Homologous recombination is a technique originally developed for targeting genes to induce or correct mutations in transcriptionally active genes. Kucherlapati, 1989, *Prog. in Nucl. Acid Res. & Mol. Biol.* 36:301. The basic technique was developed as a method for introducing specific mutations into specific regions of the mammalian genome (Thomas *et al.*, 1986, *Cell* 44:419-28; Thomas and Capecchi, 1987, *Cell* 51:503-12; Doetschman *et al.*, 1988, *Proc. Natl. Acad. Sci. U.S.A.* 85:8583-87) or to correct specific mutations within defective genes (Doetschman *et al.*, 1987, *Nature* 330:576-78). Exemplary homologous recombination techniques are described in U.S. Patent No. 5,272,071; European Patent Nos. 9193051 and 505500; PCT/US90/07642, and PCT Pub No. WO 91/09955).

Through homologous recombination, the DNA sequence to be inserted into the genome can be directed to a specific region of the gene of interest by attaching it to targeting DNA. The targeting DNA is a nucleotide sequence that is complementary

(homologous) to a region of the genomic DNA. Small pieces of targeting DNA that are complementary to a specific region of the genome are put in contact with the parental strand during the DNA replication process. It is a general property of DNA that has been inserted into a cell to hybridize, and therefore, recombine with other
5 pieces of endogenous DNA through shared homologous regions. If this complementary strand is attached to an oligonucleotide that contains a mutation or a different sequence or an additional nucleotide, it too is incorporated into the newly synthesized strand as a result of the recombination. As a result of the proofreading function, it is possible for the new sequence of DNA to serve as the template. Thus,
10 the transferred DNA is incorporated into the genome.

Attached to these pieces of targeting DNA are regions of DNA that may interact with or control the expression of an FGFR-L polypeptide, *e.g.*, flanking sequences. For example, a promoter/enhancer element, a suppressor, or an exogenous transcription modulatory element is inserted in the genome of the intended host cell in
15 proximity and orientation sufficient to influence the transcription of DNA encoding the desired FGFR-L polypeptide. The control element controls a portion of the DNA present in the host cell genome. Thus, the expression of the desired FGFR-L polypeptide may be achieved not by transfection of DNA that encodes the FGFR-L gene itself, but rather by the use of targeting DNA (containing regions of homology
20 with the endogenous gene of interest) coupled with DNA regulatory segments that provide the endogenous gene sequence with recognizable signals for transcription of an FGFR-L gene.

In an exemplary method, the expression of a desired targeted gene in a cell (*i.e.*, a desired endogenous cellular gene) is altered via homologous recombination
25 into the cellular genome at a preselected site, by the introduction of DNA which includes at least a regulatory sequence, an exon, and a splice donor site. These components are introduced into the chromosomal (genomic) DNA in such a manner that this, in effect, results in the production of a new transcription unit (in which the regulatory sequence, the exon, and the splice donor site present in the DNA construct
30 are operatively linked to the endogenous gene). As a result of the introduction of these components into the chromosomal DNA, the expression of the desired endogenous gene is altered.

Altered gene expression, as described herein, encompasses activating (or causing to be expressed) a gene which is normally silent (unexpressed) in the cell as obtained, as well as increasing the expression of a gene which is not expressed at physiologically significant levels in the cell as obtained. The embodiments further encompass changing the pattern of regulation or induction such that it is different from the pattern of regulation or induction that occurs in the cell as obtained, and reducing (including eliminating) the expression of a gene which is expressed in the cell as obtained.

One method by which homologous recombination can be used to increase, or cause, FGFR-L polypeptide production from a cell's endogenous FGFR-L gene involves first using homologous recombination to place a recombination sequence from a site-specific recombination system (*e.g.*, Cre/loxP, FLP/FRT) (Sauer, 1994, *Curr. Opin. Biotechnol.*, 5:521-27; Sauer, 1993, *Methods Enzymol.*, 225:890-900) upstream of (*i.e.*, 5' to) the cell's endogenous genomic FGFR-L polypeptide coding region. A plasmid containing a recombination site homologous to the site that was placed just upstream of the genomic FGFR-L polypeptide coding region is introduced into the modified cell line along with the appropriate recombinase enzyme. This recombinase causes the plasmid to integrate, via the plasmid's recombination site, into the recombination site located just upstream of the genomic FGFR-L polypeptide coding region in the cell line (Baubonis and Sauer, 1993, *Nucleic Acids Res.* 21:2025-29; O'Gorman *et al.*, 1991, *Science* 251:1351-55). Any flanking sequences known to increase transcription (*e.g.*, enhancer/promoter, intron, translational enhancer), if properly positioned in this plasmid, would integrate in such a manner as to create a new or modified transcriptional unit resulting in *de novo* or increased FGFR-L polypeptide production from the cell's endogenous FGFR-L gene.

A further method to use the cell line in which the site specific recombination sequence had been placed just upstream of the cell's endogenous genomic FGFR-L polypeptide coding region is to use homologous recombination to introduce a second recombination site elsewhere in the cell line's genome. The appropriate recombinase enzyme is then introduced into the two-recombination-site cell line, causing a recombination event (deletion, inversion, and translocation) (Sauer, 1994, *Curr. Opin. Biotechnol.*, 5:521-27; Sauer, 1993, *Methods Enzymol.*, 225:890-900) that would

create a new or modified transcriptional unit resulting in *de novo* or increased FGFR-L polypeptide production from the cell's endogenous FGFR-L gene.

5 An additional approach for increasing, or causing, the expression of FGFR-L polypeptide from a cell's endogenous FGFR-L gene involves increasing, or causing, the expression of a gene or genes (*e.g.*, transcription factors) and/or decreasing the expression of a gene or genes (*e.g.*, transcriptional repressors) in a manner which results in *de novo* or increased FGFR-L polypeptide production from the cell's endogenous FGFR-L gene. This method includes the introduction of a non-naturally occurring polypeptide (*e.g.*, a polypeptide comprising a site specific DNA binding domain fused to a transcriptional factor domain) into the cell such that *de novo* or increased FGFR-L polypeptide production from the cell's endogenous FGFR-L gene results.

10 The present invention further relates to DNA constructs useful in the method of altering expression of a target gene. In certain embodiments, the exemplary DNA constructs comprise: (a) one or more targeting sequences, (b) a regulatory sequence, (c) an exon, and (d) an unpaired splice-donor site. The targeting sequence in the DNA construct directs the integration of elements (a) - (d) into a target gene in a cell such that the elements (b) - (d) are operatively linked to sequences of the endogenous target gene. In another embodiment, the DNA constructs comprise: (a) one or more targeting sequences, (b) a regulatory sequence, (c) an exon, (d) a splice-donor site, (e) an intron, and (f) a splice-acceptor site, wherein the targeting sequence directs the integration of elements (a) - (f) such that the elements of (b) - (f) are operatively linked to the endogenous gene. The targeting sequence is homologous to the preselected site in the cellular chromosomal DNA with which homologous recombination is to occur. In the construct, the exon is generally 3' of the regulatory sequence and the splice-donor site is 3' of the exon.

25 If the sequence of a particular gene is known, such as the nucleic acid sequence of FGFR-L polypeptide presented herein, a piece of DNA that is complementary to a selected region of the gene can be synthesized or otherwise obtained, such as by appropriate restriction of the native DNA at specific recognition sites bounding the region of interest. This piece serves as a targeting sequence upon insertion into the cell and will hybridize to its homologous region within the genome. If this hybridization occurs during DNA replication, this piece of DNA, and any

additional sequence attached thereto, will act as an Okazaki fragment and will be incorporated into the newly synthesized daughter strand of DNA. The present invention, therefore, includes nucleotides encoding an FGFR-L polypeptide, which nucleotides may be used as targeting sequences.

5 FGFR-L polypeptide cell therapy, *e.g.*, the implantation of cells producing FGFR-L polypeptides, is also contemplated. This embodiment involves implanting cells capable of synthesizing and secreting a biologically active form of FGFR-L polypeptide. Such FGFR-L polypeptide-producing cells can be cells that are natural
10 producers of FGFR-L polypeptides or may be recombinant cells whose ability to produce FGFR-L polypeptides has been augmented by transformation with a gene encoding the desired FGFR-L polypeptide or with a gene augmenting the expression of FGFR-L polypeptide. Such a modification may be accomplished by means of a vector suitable for delivering the gene as well as promoting its expression and secretion. In order to minimize a potential immunological reaction in patients being
15 administered an FGFR-L polypeptide, as may occur with the administration of a polypeptide of a foreign species, it is preferred that the natural cells producing FGFR-L polypeptide be of human origin and produce human FGFR-L polypeptide. Likewise, it is preferred that the recombinant cells producing FGFR-L polypeptide be transformed with an expression vector containing a gene encoding a human FGFR-L
20 polypeptide.

 Implanted cells may be encapsulated to avoid the infiltration of surrounding tissue. Human or non-human animal cells may be implanted in patients in biocompatible, semipermeable polymeric enclosures or membranes that allow the release of FGFR-L polypeptide, but that prevent the destruction of the cells by the
25 patient's immune system or by other detrimental factors from the surrounding tissue. Alternatively, the patient's own cells, transformed to produce FGFR-L polypeptides *ex vivo*, may be implanted directly into the patient without such encapsulation.

 Techniques for the encapsulation of living cells are known in the art, and the preparation of the encapsulated cells and their implantation in patients may be
30 routinely accomplished. For example, Baetge *et al.* (PCT Pub. No. WO 95/05452 and PCT/US94/09299) describe membrane capsules containing genetically engineered cells for the effective delivery of biologically active molecules. The capsules are biocompatible and are easily retrievable. The capsules encapsulate cells transfected

with recombinant DNA molecules comprising DNA sequences coding for biologically active molecules operatively linked to promoters that are not subject to down-regulation *in vivo* upon implantation into a mammalian host. The devices provide for the delivery of the molecules from living cells to specific sites within a recipient. In addition, *see* U.S. Patent Nos. 4,892,538; 5,011,472; and 5,106,627. A system for encapsulating living cells is described in PCT Pub. No. WO 91/10425 (Aebischer *et al.*). *See also*, PCT Pub. No. WO 91/10470 (Aebischer *et al.*); Winn *et al.*, 1991, *Exper. Neurol.* 113:322-29; Aebischer *et al.*, 1991, *Exper. Neurol.* 111:269-75; and Tresco *et al.*, 1992, *ASAIO* 38:17-23.

In vivo and *in vitro* gene therapy delivery of FGFR-L polypeptides is also envisioned. One example of a gene therapy technique is to use the FGFR-L gene (either genomic DNA, cDNA, and/or synthetic DNA) encoding an FGFR-L polypeptide which may be operably linked to a constitutive or inducible promoter to form a "gene therapy DNA construct." The promoter may be homologous or heterologous to the endogenous FGFR-L gene, provided that it is active in the cell or tissue type into which the construct will be inserted. Other components of the gene therapy DNA construct may optionally include DNA molecules designed for site-specific integration (*e.g.*, endogenous sequences useful for homologous recombination), tissue-specific promoters, enhancers or silencers, DNA molecules capable of providing a selective advantage over the parent cell, DNA molecules useful as labels to identify transformed cells, negative selection systems, cell specific binding agents (as, for example, for cell targeting), cell-specific internalization factors, transcription factors enhancing expression from a vector, and factors enabling vector production.

A gene therapy DNA construct can then be introduced into cells (either *ex vivo* or *in vivo*) using viral or non-viral vectors. One means for introducing the gene therapy DNA construct is by means of viral vectors as described herein. Certain vectors, such as retroviral vectors, will deliver the DNA construct to the chromosomal DNA of the cells, and the gene can integrate into the chromosomal DNA. Other vectors will function as episomes, and the gene therapy DNA construct will remain in the cytoplasm.

In yet other embodiments, regulatory elements can be included for the controlled expression of the FGFR-L gene in the target cell. Such elements are turned

on in response to an appropriate effector. In this way, a therapeutic polypeptide can be expressed when desired. One conventional control means involves the use of small molecule dimerizers or rapalogs to dimerize chimeric proteins which contain a small molecule-binding domain and a domain capable of initiating a biological process, such as a DNA-binding protein or transcriptional activation protein (*see* PCT Pub. Nos. WO 96/41865, WO 97/31898, and WO 97/31899). The dimerization of the proteins can be used to initiate transcription of the transgene.

An alternative regulation technology uses a method of storing proteins expressed from the gene of interest inside the cell as an aggregate or cluster. The gene of interest is expressed as a fusion protein that includes a conditional aggregation domain that results in the retention of the aggregated protein in the endoplasmic reticulum. The stored proteins are stable and inactive inside the cell. The proteins can be released, however, by administering a drug (*e.g.*, small molecule ligand) that removes the conditional aggregation domain and thereby specifically breaks apart the aggregates or clusters so that the proteins may be secreted from the cell. *See* Aridor *et al.*, 2000, *Science* 287:816-17 and Rivera *et al.*, 2000, *Science* 287:826-30.

Other suitable control means or gene switches include, but are not limited to, the systems described herein. Mifepristone (RU486) is used as a progesterone antagonist. The binding of a modified progesterone receptor ligand-binding domain to the progesterone antagonist activates transcription by forming a dimer of two transcription factors that then pass into the nucleus to bind DNA. The ligand-binding domain is modified to eliminate the ability of the receptor to bind to the natural ligand. The modified steroid hormone receptor system is further described in U.S. Patent No. 5,364,791 and PCT Pub. Nos. WO 96/40911 and WO 97/10337.

Yet another control system uses ecdysone (a fruit fly steroid hormone) which binds to and activates an ecdysone receptor (cytoplasmic receptor). The receptor then translocates to the nucleus to bind a specific DNA response element (promoter from ecdysone-responsive gene). The ecdysone receptor includes a transactivation domain, DNA-binding domain, and ligand-binding domain to initiate transcription. The ecdysone system is further described in U.S. Patent No. 5,514,578 and PCT Pub. Nos. WO 97/38117, WO 96/37609, and WO 93/03162.

Another control means uses a positive tetracycline-controllable transactivator. This system involves a mutated tet repressor protein DNA-binding domain (mutated tet R-4 amino acid changes which resulted in a reverse tetracycline-regulated transactivator protein, *i.e.*, it binds to a tet operator in the presence of tetracycline) linked to a polypeptide which activates transcription. Such systems are described in U.S. Patent Nos. 5,464,758, 5,650,298, and 5,654,168.

Additional expression control systems and nucleic acid constructs are described in U.S. Patent Nos. 5,741,679 and 5,834,186, to Innovir Laboratories Inc.

In vivo gene therapy may be accomplished by introducing the gene encoding FGFR-L polypeptide into cells via local injection of an FGFR-L nucleic acid molecule or by other appropriate viral or non-viral delivery vectors. Hefti 1994, *Neurobiology* 25:1418-35. For example, a nucleic acid molecule encoding an FGFR-L polypeptide may be contained in an adeno-associated virus (AAV) vector for delivery to the targeted cells (*see, e.g.*, Johnson, PCT Pub. No. WO 95/34670; PCT App. No. PCT/US95/07178). The recombinant AAV genome typically contains AAV inverted terminal repeats flanking a DNA sequence encoding an FGFR-L polypeptide operably linked to functional promoter and polyadenylation sequences.

Alternative suitable viral vectors include, but are not limited to, retrovirus, adenovirus, herpes simplex virus, lentivirus, hepatitis virus, parvovirus, papovavirus, poxvirus, alphavirus, coronavirus, rhabdovirus, paramyxovirus, and papilloma virus vectors. U.S. Patent No. 5,672,344 describes an *in vivo* viral-mediated gene transfer system involving a recombinant neurotrophic HSV-1 vector. U.S. Patent No. 5,399,346 provides examples of a process for providing a patient with a therapeutic protein by the delivery of human cells which have been treated *in vitro* to insert a DNA segment encoding a therapeutic protein. Additional methods and materials for the practice of gene therapy techniques are described in U.S. Patent Nos. 5,631,236 (involving adenoviral vectors), 5,672,510 (involving retroviral vectors), 5,635,399 (involving retroviral vectors expressing cytokines).

Nonviral delivery methods include, but are not limited to, liposome-mediated transfer, naked DNA delivery (direct injection), receptor-mediated transfer (ligand-DNA complex), electroporation, calcium phosphate precipitation, and microparticle bombardment (*e.g.*, gene gun). Gene therapy materials and methods may also include inducible promoters, tissue-specific enhancer-promoters, DNA sequences designed for

site-specific integration, DNA sequences capable of providing a selective advantage over the parent cell, labels to identify transformed cells, negative selection systems and expression control systems (safety measures), cell-specific binding agents (for cell targeting), cell-specific internalization factors, and transcription factors to enhance
5 expression by a vector as well as methods of vector manufacture. Such additional methods and materials for the practice of gene therapy techniques are described in U.S. Patent Nos. 4,970,154 (involving electroporation techniques), 5,679,559 (describing a lipoprotein-containing system for gene delivery), 5,676,954 (involving liposome carriers), 5,593,875 (describing methods for calcium phosphate
10 transfection), and 4,945,050 (describing a process wherein biologically active particles are propelled at cells at a speed whereby the particles penetrate the surface of the cells and become incorporated into the interior of the cells), and PCT Pub. No. WO 96/40958 (involving nuclear ligands).

It is also contemplated that FGFR-L gene therapy or cell therapy can further
15 include the delivery of one or more additional polypeptide(s) in the same or a different cell(s). Such cells may be separately introduced into the patient, or the cells may be contained in a single implantable device, such as the encapsulating membrane described above, or the cells may be separately modified by means of viral vectors.

A means to increase endogenous FGFR-L polypeptide expression in a cell via
20 gene therapy is to insert one or more enhancer elements into the FGFR-L polypeptide promoter, where the enhancer elements can serve to increase transcriptional activity of the FGFR-L gene. The enhancer elements used will be selected based on the tissue in which one desires to activate the gene – enhancer elements known to confer promoter activation in that tissue will be selected. For example, if a gene encoding an
25 FGFR-L polypeptide is to be “turned on” in T-cells, the *lck* promoter enhancer element may be used. Here, the functional portion of the transcriptional element to be added may be inserted into a fragment of DNA containing the FGFR-L polypeptide promoter (and optionally, inserted into a vector and/or 5’ and/or 3’ flanking sequences) using standard cloning techniques. This construct, known as a
30 “homologous recombination construct,” can then be introduced into the desired cells either *ex vivo* or *in vivo*.

Gene therapy also can be used to decrease FGFR-L polypeptide expression by modifying the nucleotide sequence of the endogenous promoter. Such modification is

typically accomplished via homologous recombination methods. For example, a DNA molecule containing all or a portion of the promoter of the FGFR-L gene selected for inactivation can be engineered to remove and/or replace pieces of the promoter that regulate transcription. For example, the TATA box and/or the binding site of a transcriptional activator of the promoter may be deleted using standard molecular biology techniques; such deletion can inhibit promoter activity thereby repressing the transcription of the corresponding FGFR-L gene. The deletion of the TATA box or the transcription activator binding site in the promoter may be accomplished by generating a DNA construct comprising all or the relevant portion of the FGFR-L polypeptide promoter (from the same or a related species as the FGFR-L gene to be regulated) in which one or more of the TATA box and/or transcriptional activator binding site nucleotides are mutated via substitution, deletion and/or insertion of one or more nucleotides. As a result, the TATA box and/or activator binding site has decreased activity or is rendered completely inactive. This construct, which also will typically contain at least about 500 bases of DNA that correspond to the native (endogenous) 5' and 3' DNA sequences adjacent to the promoter segment that has been modified, may be introduced into the appropriate cells (either *ex vivo* or *in vivo*) either directly or via a viral vector as described herein. Typically, the integration of the construct into the genomic DNA of the cells will be via homologous recombination, where the 5' and 3' DNA sequences in the promoter construct can serve to help integrate the modified promoter region via hybridization to the endogenous chromosomal DNA.

Therapeutic Uses

FGFR-L nucleic acid molecules, polypeptides, and agonists and antagonists thereof can be used to treat, diagnose, ameliorate, or prevent a number of diseases, disorders, or conditions, including those recited herein.

FGFR-L polypeptide agonists and antagonists include those molecules which regulate FGFR-L polypeptide activity and either increase or decrease at least one activity of the mature form of the FGFR-L polypeptide. Agonists or antagonists may be co-factors, such as a protein, peptide, carbohydrate, lipid, or small molecular weight molecule, which interact with FGFR-L polypeptide and thereby regulate its activity. Potential polypeptide agonists or antagonists include antibodies that react

with either soluble or membrane-bound forms of FGFR-L polypeptides that comprise part or all of the extracellular domains of the said proteins. Molecules that regulate FGFR-L polypeptide expression typically include nucleic acids encoding FGFR-L polypeptide that can act as anti-sense regulators of expression.

5 The extra-cellular domain of FGFR-L polypeptide was found to share sequence identity with the Fibroblast Growth Factor (FGF) Receptor family of genes, FGFR-L nucleic acid molecules, polypeptides, and agonists and antagonists thereof (including, but not limited to, anti-FGFR-L selective binding agents) may be useful in the identification of novel growth factors.

10 The sequence identity between FGFR-L polypeptide and the FGF Receptor family also suggests that FGFR-L polypeptides may play a role in mitogenesis in fibroblasts, endothelial cells, and epithelial cells. Such epithelial cells include pancreatic ductal cells, which have been shown to differentiate in response to an FGF to form insulin producing beta islet cells. In addition to a 3-4 kb FGFR-L transcript,
15 pancreas also expresses a 6 kb transcript which may encode a FGFR-L polypeptide variant. This potential FGFR-L polypeptide variant may have activities that differ from those of the FGFR-L transcript. Accordingly, FGFR-L nucleic acid molecules, polypeptides, and agonists and antagonists thereof may be useful in tissue repair, wound healing, the modulation of angiogenesis, and the diagnosis and treatment of
20 diabetes.

 In several tumor cell lines, the extra-cellular domain of FGFR-L polypeptide is shed into the culture medium. This suggests that the FGFR-L polypeptide extra-cellular domain may play a role in the growth and/or differentiation of tumor cells. Accordingly, FGFR-L nucleic acid molecules, polypeptides, and agonists and
25 antagonists thereof may be useful in the diagnosis and treatment of cancer.

 The FGFR-L gene was found to be up-regulated in bone marrow stromal cell lines that support the maintenance of hematopoietic stem cells. Accordingly, FGFR-L nucleic acid molecules and polypeptides may be useful for *ex vivo* expansion of stem cells, gene therapy protocols, or treatment of hematopoietic disorders.

30 The FGFR-L gene was also found to be up-regulated under conditions of osteoclastogenesis. Accordingly, FGFR-L nucleic acid molecules, polypeptides, and agonists and antagonists thereof may be useful in the diagnosis and treatment of bone

disorders including, but not limited to, osteoporosis, osteopetrosis, osteogenesis imperfecta, Paget's disease, periodontal disease, and hypercalcemia.

5 FGFR-L polypeptide expression was detected in kidney. Accordingly, FGFR-L nucleic acid molecules, polypeptides, and agonists and antagonists thereof may be useful for the diagnosis and/or treatment of diseases involving the kidney. Examples of such diseases include, but are not limited to, acute and chronic glomerulonephritis. Other diseases associated with the kidney are encompassed within the scope of this invention.

10 The FGFR-L gene is most abundantly expressed in adipose tissue as determined by *in situ* hybridization. Based on this expression pattern, FGFR-L polypeptides may play a role in adipogenesis or in adipocyte function, including but not limited to, energy balance control and lipolysis. Accordingly, FGFR-L nucleic acid molecules, polypeptides, and agonists and antagonists thereof may be useful for achieving dietary weight loss, weight gain, or treating cachexia.

15 Agonists or antagonists of FGFR-L polypeptide function may be used (simultaneously or sequentially) in combination with one or more cytokines, growth factors, antibiotics, anti-inflammatories, and/or chemotherapeutic agents as is appropriate for the condition being treated.

20 Other diseases caused by or mediated by undesirable levels of FGFR-L polypeptides are encompassed within the scope of the invention. Undesirable levels include excessive levels of FGFR-L polypeptides and sub-normal levels of FGFR-L polypeptides.

Uses of FGFR-L Nucleic Acids and Polypeptides

25 Nucleic acid molecules of the invention (including those that do not themselves encode biologically active polypeptides) may be used to map the locations of the FGFR-L gene and related genes on chromosomes. Mapping may be done by techniques known in the art, such as PCR amplification and *in situ* hybridization.

30 FGFR-L nucleic acid molecules (including those that do not themselves encode biologically active polypeptides), may be useful as hybridization probes in diagnostic assays to test, either qualitatively or quantitatively, for the presence of an FGFR-L nucleic acid molecule in mammalian tissue or bodily fluid samples.

Other methods may also be employed where it is desirable to inhibit the

activity of one or more FGFR-L polypeptides. Such inhibition may be effected by nucleic acid molecules that are complementary to and hybridize to expression control sequences (triple helix formation) or to FGFR-L mRNA. For example, antisense DNA or RNA molecules, which have a sequence that is complementary to at least a portion of an FGFR-L gene can be introduced into the cell. Anti-sense probes may be designed by available techniques using the sequence of the FGFR-L gene disclosed herein. Typically, each such antisense molecule will be complementary to the start site (5' end) of each selected FGFR-L gene. When the antisense molecule then hybridizes to the corresponding FGFR-L mRNA, translation of this mRNA is prevented or reduced. Anti-sense inhibitors provide information relating to the decrease or absence of an FGFR-L polypeptide in a cell or organism.

Alternatively, gene therapy may be employed to create a dominant-negative inhibitor of one or more FGFR-L polypeptides. In this situation, the DNA encoding a mutant polypeptide of each selected FGFR-L polypeptide can be prepared and introduced into the cells of a patient using either viral or non-viral methods as described herein. Each such mutant is typically designed to compete with endogenous polypeptide in its biological role.

In addition, an FGFR-L polypeptide, whether biologically active or not, may be used as an immunogen, that is, the polypeptide contains at least one epitope to which antibodies may be raised. Selective binding agents that bind to an FGFR-L polypeptide (as described herein) may be used for *in vivo* and *in vitro* diagnostic purposes, including, but not limited to, use in labeled form to detect the presence of FGFR-L polypeptide in a body fluid or cell sample. The antibodies may also be used to prevent, treat, or diagnose a number of diseases and disorders, including those recited herein. The antibodies may bind to an FGFR-L polypeptide so as to diminish or block at least one activity characteristic of an FGFR-L polypeptide, or may bind to a polypeptide to increase at least one activity characteristic of an FGFR-L polypeptide (including by increasing the pharmacokinetics of the FGFR-L polypeptide).

FGFR-L polypeptides can be used to clone FGFR-L polypeptide ligands using an "expression cloning" strategy. Radiolabeled (¹²⁵Iodine) FGFR-L polypeptide or "affinity/activity-tagged" FGFR-L polypeptide (such as an Fc fusion or an alkaline phosphatase fusion) can be used in binding assays to identify a cell type, cell line, or tissue that expresses FGFR-L polypeptide ligands. RNA isolated from such cells or

tissues can then be converted to cDNA, cloned into a mammalian expression vector, and transfected into mammalian cells (e.g., COS or 293) to create an expression library. Radiolabeled or tagged FGFR-L polypeptide can then be used as an affinity reagent to identify and isolate the subset of cells in this library expressing FGFR-L polypeptide ligands. DNA is then isolated from these cells and transfected into mammalian cells to create a secondary expression library in which the fraction of cells expressing FGFR-L polypeptide ligands would be many-fold higher than in the original library. This enrichment process can be repeated iteratively until a single recombinant clone containing an FGFR-L polypeptide ligand is isolated. Isolation of FGFR-L polypeptide ligands is useful for identifying or developing novel agonists and antagonists of the FGFR-L polypeptide signaling pathway. Such agonists and antagonists include FGFR-L polypeptide ligands, anti-FGFR-L polypeptide ligand antibodies, small molecules, or antisense oligonucleotides.

The murine and human FGFR-L nucleic acids of the present invention are also useful tools for isolating the corresponding chromosomal FGFR-L polypeptide genes. For example, mouse chromosomal DNA containing FGFR-L sequences can be used to construct knockout mice, thereby permitting an examination of the *in vivo* role for FGFR-L polypeptide. The human FGFR-L genomic DNA can be used to identify heritable tissue-degenerating diseases.

A deposit of cDNA encoding human FGFR-L polypeptide, having Accession No. _____, was made with the American Type Culture Collection, 10801 University Boulevard, Manassas, VA 20110-2209 on January 31, 2000.

The following examples are intended for illustration purposes only, and should not be construed as limiting the scope of the invention in any way.

Example 1: Cloning of the Murine FGFR-L Polypeptide Gene

Generally, materials and methods as described in Sambrook *et al. supra* were used to clone and analyze the gene encoding rat FGFR-L polypeptide.

Sequences encoding the murine FGFR-L polypeptide were isolated from a mouse cDNA library derived from a mixture of two hematopoietic stem cell supportive bone marrow stromal cell lines (F4 and F10). Murine bone marrow stromal cell lines D3, F4, and F10 were obtained from Dr. R. Ploemacher (Erasmus University, Rotterdam, The Netherlands) and cultured at 32°C and 5% CO₂ in IMDM

supplemented with 10% fetal bovine serum, 5% horse serum, 2 mM glutamine, 0.1 mM β -mercaptoethanol and 1 μ M hydrocortisone (Na Succinate salt). A mouse bone marrow stromal cDNA library was prepared by isolating RNA from F4 and F10 cells using the Trizol method (LTI). Poly-A RNA was purified using oligo-dT magnetic
5 beads (Dynal) and equal amounts of poly-A RNA (1.5 ug each of F4 and F10 RNA) were mixed. An oligo-dT primed full-length cDNA library was constructed from the F4/F10 RNA mixture using the Superscript Plasmid System for cDNA Synthesis and Plasmid Cloning (LTI).

The mouse bone marrow stromal cDNA library, containing 6×10^6
10 transformants, was plated, and 3.4×10^4 colonies were selected and transferred in parallel into 96-well plates and spotted onto filters. The filters were then probed with 32 P-dCTP-labeled first strand cDNA generated from poly-A mRNA isolated from the bone marrow stromal D3 cell line. Of the 3.4×10^4 colonies spotted onto filters, 11,232 failed to hybridize with the D3 probe. Plasmid was isolated from these non-
15 hybridizing colonies and the 5' end of their cDNA inserts sequenced.

One clone (smsf2-00017-f4), showing homology with various members of the FGF receptor family, was identified in the sequence analysis. A full-length clone (smsf2-00017-f4-41.6) was obtained by screening a Southern blot of 56 mouse bone marrow stromal cDNA pools – each pool comprising 1×10^4 clones from the mouse
20 bone marrow stromal cDNA library. The pool possessing the longest insert was subsequently plated and rescreened.

Sequence analysis of the full-length cDNA for murine FGFR-L polypeptide indicated that the gene encodes a type I transmembrane protein (Figure 1B, predicted transmembrane domain: L-P-W-P-V-V-I-G-I-P-A-G-A-V-F-I-L-G-T-V-L-L-W-L-C;
25 SEQ ID NO: 12). The murine FGFR-L polypeptide gene comprises a 1587 bp open reading frame encoding a protein of 529 amino acids and possessing a potential signal peptide at its amino terminus (Figure 1A, predicted signal peptide: M-T-R-S-P-A-L-L-L-L-L-G-A-L-P-S-A-E-A; SEQ ID NO: 11). Figures 1A-1C illustrate the nucleotide sequence of the murine FGFR-L nucleic acid sequence and the deduced
30 amino acid sequence of murine FGFR-L polypeptide. A murine extracellular domain-Fc fusion protein has an apparent Molecular Weight, as determined by SDS-PAGE, of approximately 55kD.

While the extracellular domain of FGFR-L polypeptide is most closely related to the FGF receptor family, the protein's cytoplasmic domain does not contain a kinase domain or other recognizable domain. Figures 2A-2B illustrate the amino acid sequence alignment of murine FGFR-L polypeptide and the known protein for which FGFR-L polypeptide shares the closest homology, Iberian ribbed newt (*Pleurodeles waltlii*) FGF receptor-4. Computer analysis using the BLAST program, also indicated that murine FGFR-L polypeptide was closely related to a single, 486 bp human EST (GenBank accession number AI245701) isolated from human kidney. A 379 bp stretch of this EST showed an 87% identity with FGFR-L polypeptide, suggesting the existence of a human ortholog.

Example 2: Cloning of the Human FGFR-L Polypeptide Gene

Generally, materials and methods as described in Sambrook *et al. supra* are used to clone and analyze the gene encoding human FGFR-L polypeptide.

A human spleen and mixed tissue cDNA library was prepared as follows. Total RNA was extracted from human tissues using Trizol extraction procedures (Gibco-BRL, Rockville, MD) and poly-A⁺ RNA was selected from this total RNA using Dynabeads (Dynal, Oslo, Norway) according to the manufacturer's recommended protocol. Random primed or oligo(dT) primed cDNA was synthesized from this poly-A⁺ RNA using the the Superscript Plasmid System for cDNA Synthesis and Plasmid Cloning kit (Gibco-BRL, Rockville, MD) according to the manufacturer's recommended protocol. The resulting cDNA was digested with appropriate restriction enzymes and cloned into the pSPORT 1 vector. Ligation products were transformed into *E. coli* using standard techniques known in the art and transformants were selected on bacterial media plates containing an appropriate antibiotic. The cDNA library consisted of all, or a subset, of these transformants.

Plasmid DNA isolated from pools of 1×10^4 colonies was used as a template in PCR amplifications performed with the primers 5'-C-G-C-T-G-A-C-C-A-T-G-T-G-G-A-C-C-A-A-G-G-A-T-G-3' (SEQ ID NO: 13) and 5'-C-T-T-G-A-C-C-C-C-A-G-A-A-G-G-A-G-C-T-G-T-C-G-G-3' (SEQ ID NO: 14). The PCR primers were designed on the basis of the human EST sequence AI245701 described in Example 1. Several pools yielded a 234 bp fragment which was subcloned and determined to have a nucleic acid sequence corresponding to positions 208-441 of the human FGFR-L

nucleic acid sequence shown in Figure 3A.

Plasmid pools that yielded the 234 bp PCR product in the amplification reactions above were then plated for colony hybridization analysis. Plated colonies were screened using the 234 bp PCR fragment generated above as a probe following radiolabeling with the Rediprime II random prime labeling kit (Amersham, Piscataway, NJ). A 1333 bp cDNA insert was determined to have a sequence corresponding to positions 118-1450 of the human FGFR-L nucleic acid sequence shown in Figures 3A-3C. Assembly of this 1333 bp sequence and the 234 bp sequence of AI245701 yielded the human FGFR-L nucleic acid sequence shown in Figures 3A-3C.

Baker *et al.* (PCT Pub. No. WO 99/63088) teach a polypeptide sequence of 504 amino acids (SEQ ID NO: 15) which they call PRO943 that shares sequence identity with human FGFR-L polypeptide. Ruben and Young (PCT Pub. No. WO 00/24756) teach a nucleic acid sequence of 3112 bp (SEQ ID NO: 16) encoding a polypeptide of 504 amino acids (SEQ ID NO: 17) which they call Fibroblast Growth Factor Receptor-5 (FGFR5) that shares sequence identity with human FGFR-L polypeptide. Finally, Wiedemann and Trueb, 2000, *Genomics* 69:275-79, teach a nucleic acid sequence of 3080 bp (SEQ ID NO: 18) encoding a polypeptide 504 amino acids (SEQ ID NO: 19) which they call Fibroblast Growth Factor Receptor-Like Protein 1 (FGFRL1) that shares sequence identity with human FGFR-L polypeptide.

Example 3: FGFR-L mRNA Expression

The expression of FGFR-L mRNA was examined by Northern blot analysis. Multiple murine and human tissue northern blots (Clontech) were probed with a ³²P-dCTP labeled, 234 bp PCR fragment corresponding to a portion of the human FGFR-L gene (see Example 2). Additional blots containing RNA isolated from a variety of cell lines were also screened with this probe.

Northern blots were prehybridized for 2 hours at 42°C in 5X SSC, 35% deionized formamide, 0.05% (w/v) sodium pyrophosphate, 20 mM sodium phosphate pH 6.8, 5 mM EDTA, 5X Denhardt's solution, 0.2% SDS, and 94 µg/mL denatured salmon sperm DNA, and then were hybridized at 42°C overnight in fresh prehybridization buffer containing approximately 1 ng/mL of the labeled probe.

Following hybridization, the filters were washed once in prehybridization buffer for 5 minutes at room temperature, once for 5 minutes at room temperature in 2X SSC and 0.1% SDS, and then twice for 20 minutes at 42°C in 2X SSC and 0.1% SDS. The blots were then exposed to autoradiography.

5 Analysis of the Northern blots (Figures 5-7) indicated that a single transcript having a molecular mass of 2.9 kb was highly expressed in murine liver, kidney, F4 and F10 bone marrow stromal cells, NIH-3T3 cells, and ST2 bone marrow stromal cells (following exposure with vitamin D3 and dexamethasone). The detection of a single transcript in the positive samples suggests that, in these tissues, there is no
10 obvious splice variant encoding a longer cytoplasmic domain that could contain a kinase domain or other recognizable domain. Weak expression of the transcript was detected in murine heart, brain, lung, skeletal muscle, testis, F10 bone marrow stromal cells, and ST2 cells (prior to exposure with Vitamin D3 and Dexamethasone).

 Northern blot analysis (Figures 8-10) also indicated that a transcript having a
15 molecular mass of 3.2 kb was highly expressed among human tissues and that a transcript having a molecular mass of 6.0 kb was expressed in human pancreas. The existence of the 6.0 kb transcript suggests that a functionally distinct FGFR-L protein variant may exist. Lower expression of the 3.2 kb transcript was seen in liver, kidney, heart, skeletal muscle, brain, and the cell lines HeLa, K562, SW480, Molt4, and Raji.

20 Northern blot analysis (Figure 11) also indicated that single transcript could be detected in the following cell lines: 266-6 (mouse acinar pancreatic tumor), AR42J (rat pancreas tumor, exocrine), CaPan I (human pancreatic adenocarcinoma), HIG-82 (rabbit synoviocyte), OHS4 (human osteoblast), SW 1353 (human chondrosarcoma, humerus), SW 872 (human liposarcoma), K562 (old, *i.e.*, later passage; chronic
25 myelogenous leukemia; later passage), K562 (new, *i.e.*, earlier passage), Jurkat (human T cell line), and F4 (murine bone marrow derived stromal cell line). Probes derived from human and murine FGFR-L cDNA were also capable of detecting FGFR-L mRNA in human and murine adipose tissue, respectively (Figures 12A-12B).

30 The expression of FGFR-L mRNA was also examined in RNase protection assays (Figure 13). A signal was detected in most of the tissues that were examined, with the strongest signal being detected in brown adipose tissue, white adipose tissue, and testis.

The expression of FGFR-L mRNA was localized by *in situ* hybridization, using standard techniques. The highest levels of FGFR-L mRNA were found in both white and brown adipose tissue; FGFR-L mRNA expression was detected in a perirenal adipose depot adjacent to the adrenal gland and kidney (Figure 14). In the digestive tissues, signal corresponding to FGFR-L mRNA was found in the small intestine (duodenum and ileum), but not in the large intestine (Figure 15). Specifically, FGFR-L mRNA was found to be expressed at the base of the crypts, most likely the Paneth cells. FGFR-L mRNA expression was also detected in the trachea (a signal was detected over the perichondral cells adjacent to the ring of hyaline cartilage surrounding the trachea) and in the uterus (a strong signal was detected over the epithelial cells lining the uterine lumen; Figure 16). The high expression levels of FGFR-L polypeptide detected in hyaline cartilage suggests possible clinical utility for FGFR-L polypeptide in the modulation of osteoarthritis since fibro-cartilage, characteristically present in osteoarthritic joints, resembles hyaline cartilage. Lower levels of FGFR-L mRNA expression were also detected in the articular cartilage at the knee joint and in the spleen – over the red pulp (hematopoietic) as opposed to the white pulp (lymphocytes; Figure 16). Lower levels of expression were detected in ovary, testis, and small intestine. The high level of FGFR-L mRNA expression in adipose tissue warrents caution in the interpretation of the Northern blot data as a result of the contamination of some tissues with adipose tissue. This may be true in particular for the high level of expression observed in human pancreas.

The expression levels of murine FGFR-L mRNA detected in three bone marrow stromal cell lines (*i.e.*, D3, F4, and F10) correlate with the ability of these cell lines to support hematopoietic stem cells. The highest expression of murine FGFR-L mRNA was detected in the stromal cell lines providing the best support (F4 and F10; Figure 7), while much lower levels were seen in the cell line incapable of supporting hematopoietic stem cells (D3). Murine FGFR-L mRNA was also upregulated in osteoblastic ST2 cells under conditions of osteoclastogenesis (*i.e.*, in response to vitamin D3 and dexamethasone; Figure 17), a process which is known to be inhibited by bFGF (Jimi *et al.*, 1996, *J. Cell. Physiol.* 168:395-402).

Furthermore, expression analysis showed that the level of mRNA expression for murine FGFR-L polypeptide increased during fetal development with the lowest

expression being detected at the earliest time point analyzed (*i.e.*, day 7) and the highest expression being detected at the latest time point analyzed (*i.e.*, day 17) (Figure 5).

5 Finally, proteomic analysis showed that a peptide with a sequence identical to that of murine and human FGFR-L polypeptide was secreted from K562 cells and SV40 transformed AG2804 cells (human fibroblast). In this approach, protein mixtures were isolated from culture media that was conditioned by any one of a variety of cell lines and subjected to Mass Spectrometric analysis to identify the presence of individual peptide sequences. Proteomic analysis also showed N-linked
10 glycosylation at residues 231 (Asp) and 293 (Asp) in the human FGFR-L polypeptide.

Example 4: Production of Anti-FGFR-L Polypeptide Antibodies

FGFR-L polypeptide antibodies were obtained by immunizing rabbits with a polypeptide corresponding to a portion of the extracellular domain (ECD) of murine
15 FGFR-L polypeptide (Des7-FGFR-L/ECD; SEQ ID NO: 20; comprising residues 28-368 of FGFR-L polypeptide). An FGFR-L polypeptide-Fc fusion construct was also prepared using residues 1-366 of the extracellular domain (SEQ ID NO: 21 and SEQ ID NO: 22). Suitable procedures for generating antibodies were used (*see, e.g.*, Hudson and Bay, *Practical Immunology* (2nd ed., Blackwell Scientific Publications)).

20 FGFR-L polypeptide antiserum was used in Western blot analysis of SDS-PAGE separated *E.coli*-derived Des7-FGFR-L/ECD and CHO-derived FGFR-L/ECD-Fc proteins. Immunoreactive bands of 95-100 kD and of 40-45 kD were detected in the CHO- and *E. coli*-derived samples, respectively (Figure 18). A immunoreactive band of 60-70 kD was also detected in murine adipose tissue (Figure
25 19), 266-6 cells (mouse acinar pancreatic tumor), AR42J cells (rat pancreas exocrine tumor), MRC5 cells (human diploid lung fibroblasts), OHS4 cells (human osteoblast), SW1353 cells (human chondrosarcoma), and K562 cells (chronic myelogenous sarcoma) following immunoprecipitation and Western blot analysis of cell lysates and conditioned media collected from these tissues and cell lines. An additional band of
30 100-120 kD was detected in adipose tissue, OHS4 cells and K562 cells. The crude antiserum could also be used to immunoprecipitate both proteins. Using the crude antiserum in FACS analysis, FGFR-L polypeptide cell surface staining was detected on F4 bone marrow stromal cells (shown to express high levels of FGFR-L RNA), but

not on D3 bone marrow stromal cells (shown to express low levels of FGFR-L RNA; Figure 20A-20B).

Example 5: *In Vitro* Characterization of FGFR-L Polypeptides

5 Constructs encoding full-length FGFR-L polypeptide or the extracellular domain of FGFR-L polypeptide have proven to be poorly tolerated by transfected cells (e.g., CHO, 329 HEK, and stromal cell lines), as measured by the lower number and decreased growth rate of stable transfectants versus untransfected cells (Figures 21A-21D) or cells transfected with a construct encoding an FGFR-L polypeptide
10 point mutant. When *E. coli*-derived Des7-FGFR-L/ECD was added to bone marrow stromal cell cultures, FGF-mediated, but not serum-mediated growth, was inhibited (Figure 22). The observation that soluble FGFR-L/ECD inhibits growth induced by FGF proteins, but has less of an inhibitory effect on growth induced by PDGF or serum, suggests that FGFR-L polypeptide may interact, alone or with a co-receptor,
15 with a natural ligand that has homology to the FGF family. Surprisingly, this effect was not observed when CHO-derived FGFR-L/ECD-Fc fusion protein was used in place of *E. coli*-derived Des7-FGFR-L/ECD (Figure 23). Similar results were obtained for FGF and VEGF-mediated growth of human vascular endothelial cells (HUVEC). The difference in activity between the CHO-derived and *E. coli*-derived
20 FGFR-L polypeptides may be due to the amino acid sequence differences at their C- and/or N-terminus.

Example 6: *In Vivo* Characterization of FGFR-L Polypeptides

 Murine bone marrow cells, transduced with a retroviral vector carrying a
25 bicistronic message encoding full-length murine FGFR-L and the neomycin resistance gene (or the neomycin resistance gene alone) were used to transplant 10 lethally irradiated recipients as described previously (Yan *et al.*, 1999, *Exp. Hematol.* 27:1409-17).

 In five of the mice (randomly selected for evaluation), the overexpression of
30 FGFR-L polypeptide over a four month period caused a 15% decrease in total body weight, a 14% decrease in serum cholesterol and 35% decrease in serum triglyceride levels. However, three weeks later the five remaining mice were weighed and bled, and the similar changes were not observed.

In vitro characterization (Example 5) of FGFR-L polypeptide indicated that the protein is poorly tolerated by a variety of cell types such that selection against FGFR-L polypeptide expression is not unexpected. As the retroviral construct described above carries both the FGFR-L gene and the neomycin resistance gene on one bicistronic message, selection against FGFR-L polypeptide expression would result in a corresponding selection against neo expression as well, either by downregulation of transcription or by removal of the transduced cells.

Northern blot analysis of peripheral blood mononuclear cell (PBMN) RNA from two FGFR-L/neo-transduced mice exhibiting the phenotype and two neo-transduced control mice indicated that the control mice show abundant expression of neo transcripts of the expected size and the FGFR-L/neo-transduced mice do not (Figure 24). This suggests that FGFR-L polypeptide expression is actively selected against and precedes the ultimate disappearance of a transient phenotype.

While the present invention has been described in terms of the preferred embodiments, it is understood that variations and modifications will occur to those skilled in the art. Therefore, it is intended that the appended claims cover all such equivalent variations that come within the scope of the invention as claimed.

WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:
 - 5 (a) the nucleotide sequence as set forth in either SEQ ID NO: 1 or SEQ ID NO: 4;
 - (b) the nucleotide sequence of the DNA insert in ATCC Deposit No. _____;
 - 10 (c) a nucleotide sequence encoding the polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;
 - (d) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a) - (c); and
 - (e) a nucleotide sequence complementary to any of (a) - (c).
- 15 2. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:
 - (a) a nucleotide sequence encoding a polypeptide which is at least about 70 percent identical to the polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5, wherein the encoded polypeptide has an activity of the polypeptide set forth in
20 either SEQ ID NO: 2 or SEQ ID NO: 5;
 - (b) a nucleotide sequence encoding an allelic variant or splice variant of the nucleotide sequence as set forth in either SEQ ID NO: 1 or SEQ ID NO: 4, the nucleotide sequence of the DNA insert in ATCC Deposit No. _____, or (a);
 - 25 (c) a region of the nucleotide sequence of either SEQ ID NO: 1 or SEQ ID NO: 4, the DNA insert in ATCC Deposit No. _____, (a), or (b) encoding a polypeptide fragment of at least about 25 amino acid residues, wherein the polypeptide fragment has an activity of the encoded polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5, or is antigenic;
 - 30 (d) a region of the nucleotide sequence of either SEQ ID NO: 1 or SEQ ID NO: 4, the DNA insert in ATCC Deposit No. _____, or any of (a) - (c) comprising a fragment of at least about 16 nucleotides;
 - (e) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a) - (d); and

(f) a nucleotide sequence complementary to any of (a) - (d).

3. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

5 (a) a nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one conservative amino acid substitution, wherein the encoded polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

10 (b) a nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one amino acid insertion, wherein the encoded polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

15 (c) a nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one amino acid deletion, wherein the encoded polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

20 (d) a nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 which has a C- and/or N- terminal truncation, wherein the encoded polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

25 (e) a nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one modification selected from the group consisting of amino acid substitutions, amino acid insertions, amino acid deletions, C-terminal truncation, and N-terminal truncation, wherein the encoded polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

(f) a nucleotide sequence of any of (a) - (e) comprising a fragment of at least about 16 nucleotides;

30 (g) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a) - (f); and

(h) a nucleotide sequence complementary to any of (a) - (e).

4. A vector comprising the nucleic acid molecule of any of Claims 1, 2, or 3.

5. A host cell comprising the vector of Claim 4.

5

6. The host cell of Claim 5 that is a eukaryotic cell.

7. The host cell of Claim 5 that is a prokaryotic cell.

10

8. A process of producing an FGFR-L polypeptide comprising culturing the host cell of Claim 5 under suitable conditions to express the polypeptide, and optionally isolating the polypeptide from the culture.

9. A polypeptide produced by the process of Claim 8.

15

10. The process of Claim 8, wherein the nucleic acid molecule comprises promoter DNA other than the promoter DNA for the native FGFR-L polypeptide operatively linked to the DNA encoding the FGFR-L polypeptide.

20

11. The isolated nucleic acid molecule according to Claim 2, wherein the percent identity is determined using a computer program selected from the group consisting of GAP, BLASTN, FASTA, BLASTA, BLASTX, BestFit, and the Smith-Waterman algorithm.

25

12. A process for determining whether a compound inhibits FGFR-L polypeptide activity or FGFR-L polypeptide production comprising exposing a cell according to any of Claims 5, 6, or 7 to the compound and measuring FGFR-L polypeptide activity or FGFR-L polypeptide production in said cell.

30

13. An isolated polypeptide comprising the amino acid sequence selected from the group consisting of:

(a) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5; and

(b) the amino acid sequence encoded by the DNA insert in ATCC Deposit No. _____.

5 14. An isolated polypeptide comprising the amino acid sequence selected from the group consisting of:

(a) the amino acid sequence as set forth in SEQ ID NO: 3 or SEQ ID NO: 6, optionally further comprising an amino-terminal methionine;

(b) an amino acid sequence for an ortholog of either SEQ ID NO: 2 or SEQ ID NO: 5;

10 (c) an amino acid sequence which is at least about 70 percent identical to the amino acid sequence of either SEQ ID NO: 2 or SEQ ID NO: 5, wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

(d) a fragment of the amino acid sequence set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 comprising at least about 25 amino acid residues, wherein the fragment has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5, or is antigenic; and

(e) an amino acid sequence for an allelic variant or splice variant of the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5, the amino acid sequence encoded by the DNA insert in ATCC Deposit No. _____, or any of (a) - (c)

15 15. An isolated polypeptide comprising the amino acid sequence selected from the group consisting of:

(a) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one conservative amino acid substitution, wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

(b) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one amino acid insertion, wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

30 (c) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one amino acid deletion, wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

(d) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 which has a C- and/or N- terminal truncation, wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5; and

(e) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one modification selected from the group consisting of amino acid substitutions, amino acid insertions, amino acid deletions, C-terminal truncation, and N-terminal truncation, wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5.

16. An isolated polypeptide encoded by the nucleic acid molecule of any of Claims 1, 2, or 3, wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5.

17. The isolated polypeptide according to Claim 14, wherein the percent identity is determined using a computer program selected from the group consisting of GAP, BLASTP, FASTA, BLASTA, BLASTX, BestFit, and the Smith-Waterman algorithm.

18. A selective binding agent or fragment thereof which specifically binds the polypeptide of any of Claims 13, 14, or 15.

19. The selective binding agent or fragment thereof of Claim 18 that specifically binds the polypeptide comprising the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5, or a fragment thereof.

20. The selective binding agent of Claim 18 that is an antibody or fragment thereof.

21. The selective binding agent of Claim 18 that is a humanized antibody.

22. The selective binding agent of Claim 18 that is a human antibody or fragment thereof.

23. The selective binding agent of Claim 18 that is a polyclonal antibody or fragment thereof.

5 24. The selective binding agent Claim 18 that is a monoclonal antibody or fragment thereof.

25. The selective binding agent of Claim 18 that is a chimeric antibody or fragment thereof.

10 26. The selective binding agent of Claim 18 that is a CDR-grafted antibody or fragment thereof.

27. The selective binding agent of Claim 18 that is an antiidiotypic antibody or fragment thereof.

15

28. The selective binding agent of Claim 18 that is a variable region fragment.

20 29. The variable region fragment of Claim 28 that is a Fab or a Fab' fragment.

30. A selective binding agent or fragment thereof comprising at least one complementarity determining region with specificity for a polypeptide having the amino acid sequence of either SEQ ID NO: 2 or SEQ ID NO: 5.

25

31. The selective binding agent of Claim 18 that is bound to a detectable label.

30 32. The selective binding agent of Claim 18 that antagonizes FGFR-L polypeptide biological activity.

33. A method for treating, preventing, or ameliorating an FGFR-L polypeptide-related disease, condition, or disorder comprising administering to a patient an effective amount of a selective binding agent according to Claim 18.

5 34. A selective binding agent produced by immunizing an animal with a polypeptide comprising an amino acid sequence of either SEQ ID NO: 2 or SEQ ID NO: 5.

10 35. A hybridoma which produces a selective binding agent which is capable of binding a polypeptide according to any of Claims 1, 2, or 3.

36. A method of detecting or quantitating the amount of FGFR-L polypeptide using the selective binding agent or fragment of Claim 18.

15 37. A composition comprising the polypeptide of any of Claims 13, 14, or 15, and a pharmaceutically acceptable formulation agent.

20 38. The composition of Claim 37, wherein the pharmaceutically acceptable formulation agent is a carrier, adjuvant, solubilizer, stabilizer, or anti-oxidant.

39. The composition of Claim 38 wherein the polypeptide comprises the amino acid sequence as set forth in SEQ ID NO: 3 or SEQ ID NO: 6.

25 40. A polypeptide comprising a derivative of the polypeptide of any of Claims 13, 14, or 15.

41. The polypeptide of Claim 40 that is covalently modified with a water-soluble polymer.

30 42. The polypeptide of Claim 41, wherein the water-soluble polymer is selected from the group consisting of polyethylene glycol, monomethoxy-polyethylene glycol, dextran, cellulose, poly-(N-vinyl pyrrolidone) polyethylene

glycol, propylene glycol homopolymers, polypropylene oxide/ethylene oxide copolymers, polyoxyethylated polyols, and polyvinyl alcohol.

43. A composition comprising a nucleic acid molecule of any of Claims 1,
5 2, or 3 and a pharmaceutically acceptable formulation agent.

44. The composition of Claim 43, wherein said nucleic acid molecule is contained in a viral vector.

10 45. A viral vector comprising a nucleic acid molecule of any of Claims 1, 2, or 3.

46. A fusion polypeptide comprising the polypeptide of any of Claims 13, 14, or 15 fused to a heterologous amino acid sequence.

15 47. The fusion polypeptide of Claim 46, wherein the heterologous amino acid sequence is an IgG constant domain or fragment thereof.

48. A method for treating, preventing, or ameliorating a medical condition
20 comprising administering to a patient the polypeptide of any of Claims 13, 14, or 15, or the polypeptide encoded by the nucleic acid of any of Claims 1, 2, or 3.

49. The method of Claim 48 wherein the medical condition being treated, prevented, or ameliorated is a hematopoietic disorder, osteoporosis, osteopetrosis,
25 osteogenesis imperfecta, Paget's disease, periodontal disease, hypercalcemia, acute glomerulonephritis, chronic glomerulonephritis, cancer, diabetes, obesity, or cachexia.

50. A method of diagnosing a pathological condition or a susceptibility to
30 a pathological condition in a subject comprising:

(a) determining the presence or amount of expression of the polypeptide of any of Claims 13, 14, or 15, or the polypeptide encoded by the nucleic acid molecule of any of Claims 1, 2, or 3 in a sample; and

(b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or amount of expression of the polypeptide.

5 51. A device, comprising:
 (a) a membrane suitable for implantation; and
 (b) cells encapsulated within said membrane, wherein said cells secrete a protein of any of Claims 13, 14, or 15; and

10 said membrane is permeable to said protein and impermeable to materials detrimental to said cells.

 52. A method of identifying a compound which binds to an FGFR-L polypeptide comprising:

 (a) contacting the polypeptide of any of Claims 13, 14, or 15 with a
15 compound; and

 (b) determining the extent of binding of the FGFR-L polypeptide to the compound.

 53. The method of Claim 52, further comprising determining the activity
20 of the polypeptide when bound to the compound.

 54. A method of modulating levels of a polypeptide in an animal comprising administering to the animal the nucleic acid molecule of any of Claims 1, 2, or 3.

25 55. A transgenic non-human mammal comprising the nucleic acid molecule of any of Claims 1, 2, or 3.

 56. A process for determining whether a compound inhibits FGFR-L
30 polypeptide activity or FGFR-L polypeptide production comprising exposing a transgenic mammal according to Claim 55 to the compound, and measuring FGFR-L polypeptide activity or FGFR-L polypeptide production in said mammal.

gacctggggtc	ttgcggggcct	gagccctgag	tggcgctccag	tccagctccc	agtgaccgcg	60
ccccctgcttc	agggtccgacc	ggcgag	atg acg cgg agc ccc gcg ctg ctg ctg	113		
			<u>Met Thr Arg Ser Pro Ala Leu Leu Leu</u>			
			1 5			
ctg cta ttg ggg gcc ctc ccg tcg gct gag gcg gcg cga gga ccc cca	161					
<u>Leu Leu Leu Gly Ala Leu Pro Ser Ala Glu Ala Ala Arg Gly Pro Pro</u>						
10 15 20 25						
aga atg gca gac aaa gtg gtc cca cgg cag gtg gcc cgc ctg ggc cgc	209					
<u>Arg Met Ala Asp Lys Val Val Pro Arg Gln Val Ala Arg Leu Gly Arg</u>						
30 35 40						
act gtg cgg cta cag tgc cca gtg gag ggg gac cca cca ccg ttg acc	257					
<u>Thr Val Arg Leu Gln Cys Pro Val Glu Gly Asp Pro Pro Pro Leu Thr</u>						
45 50 55						
atg tgg acc aaa gat ggc cgc aca atc cac agt ggc tgg agc cgc ttc	305					
<u>Met Trp Thr Lys Asp Gly Arg Thr Ile His Ser Gly Trp Ser Arg Phe</u>						
60 65 70						
cgt gtg ctg ccc cag ggt ctg aag gtg aag gag gtg gag gcc gag gat	353					
<u>Arg Val Leu Pro Gln Gly Leu Lys Val Lys Glu Val Glu Ala Glu Asp</u>						
75 80 85						
gcc ggt gtt tat gtg tgc aag gcc acc aat ggc ttt ggc agc ctc agc	401					
<u>Ala Gly Val Tyr Val Cys Lys Ala Thr Asn Gly Phe Gly Ser Leu Ser</u>						
90 95 100 105						
gtc aac tac act ctc atc atc atg gat gat att agt cca ggg aag gag	449					
<u>Val Asn Tyr Thr Leu Ile Ile Met Asp Asp Ile Ser Pro Gly Lys Glu</u>						
110 115 120						
agc cct ggg cca ggt ggt tct tcg ggg ggc cag gag gac cca gcc agc	497					
<u>Ser Pro Gly Pro Gly Gly Ser Ser Gly Gly Gln Glu Asp Pro Ala Ser</u>						
125 130 135						
cag cag tgg gca cgg cct cgc ttc aca cag ccc tcc aag atg agg cgc	545					
<u>Gln Gln Trp Ala Arg Pro Arg Phe Thr Gln Pro Ser Lys Met Arg Arg</u>						
140 145 150						
cga gtg att gca cgg cct gtg ggt agc tct gtg cgg ctc aag tgt gtg	593					
<u>Arg Val Ile Ala Arg Pro Val Gly Ser Ser Val Arg Leu Lys Cys Val</u>						
155 160 165						
gcc agt ggg cac cca cgg cca gac atc atg tgg atg aag gat gac cag	641					
<u>Ala Ser Gly His Pro Arg Pro Asp Ile Met Trp Met Lys Asp Asp Gln</u>						
170 175 180 185						
acc ttg acg cat cta gag gct agt gaa cac aga aag aag aag tgg aca	689					
<u>Thr Leu Thr His Leu Glu Ala Ser Glu His Arg Lys Lys Lys Trp Thr</u>						
190 195 200						

FIG. 1B

ctg agc ttg aag aac ctg aag cct gaa gac agt ggc aag tac acg tgc	737
Leu Ser Leu Lys Asn Leu Lys Pro Glu Asp Ser Gly Lys Tyr Thr Cys	
205 210 215	
cggt gta tct aac aag gcc ggt gcc atc aac gcc acc tac aaa gtg gat	785
Arg Val Ser Asn Lys Ala Gly Ala Ile Asn Ala Thr Tyr Lys Val Asp	
220 225 230	
gta atc cag cgg act cgt tcc aag cct gtg ctc aca ggg aca cac cct	833
Val Ile Gln Arg Thr Arg Ser Lys Pro Val Leu Thr Gly Thr His Pro	
235 240 245	
gtg aac aca acg gtg gac ttc ggt ggg aca acg tcc ttc cag tgc aag	881
Val Asn Thr Thr Val Asp Phe Gly Gly Thr Thr Ser Phe Gln Cys Lys	
250 255 260 265	
gtg cgc agt gac gtg aag cct gtg atc cag tgg ctg aag cgg gtg gag	929
Val Arg Ser Asp Val Lys Pro Val Ile Gln Trp Leu Lys Arg Val Glu	
270 275 280	
tac ggc tcc gag gga cgc cac aac tcc acc att gat gtg ggt ggc cag	977
Tyr Gly Ser Glu Gly Arg His Asn Ser Thr Ile Asp Val Gly Gly Gln	
285 290 295	
aag ttt gtg gtg ttg ccc acg ggt gat gtg tgg tca cgg cct gat ggc	1025
Lys Phe Val Val Leu Pro Thr Gly Asp Val Trp Ser Arg Pro Asp Gly	
300 305 310	
tcc tac ctc aac aag ctg ctc atc tct cgg gcc cgc cag gat gat gct	1073
Ser Tyr Leu Asn Lys Leu Leu Ile Ser Arg Ala Arg Gln Asp Asp Ala	
315 320 325	
ggc atg tac atc tgc cta ggt gca aat acc atg ggc tac agt ttc cgt	1121
Gly Met Tyr Ile Cys Leu Gly Ala Asn Thr Met Gly Tyr Ser Phe Arg	
330 335 340 345	
agc gcc ttc ctc act gta tta cca gac ccc aaa cct cca ggg cct cct	1169
Ser Ala Phe Leu Thr Val Leu Pro Asp Pro Lys Pro Pro Gly Pro Pro	
350 355 360	
atg gct tct tca tcg tca tcc aca agc ctg cca tgg cct gtg gtg atc	1217
Met Ala Ser Ser Ser Ser Ser Thr Ser Leu Pro Trp Pro <u>Val Val Ile</u>	
365 370 375	
ggc atc cca gct ggt gct gtc ttc atc cta ggc act gtg ctg ctc tgg	1265
<u>Gly Ile Pro Ala Gly Ala Val Phe Ile Leu Gly Thr Val Leu Leu Trp</u>	
380 385 390	
ctt tgc cag acc aag aag aag cca tgt gcc cca gca tct aca ctt cct	1313
<u>Leu Cys</u> Gln Thr Lys Lys Lys Pro Cys Ala Pro Ala Ser Thr Leu Pro	
395 400 405	

FIG. 1C

gtg cct ggg cat cgt ccc cca ggg aca tcc cga gaa cgc agt ggt gac 1361
 Val Pro Gly His Arg Pro Pro Gly Thr Ser Arg Glu Arg Ser Gly Asp
 410 415 420 425

aag gac ctg ccc tca ttg gct gtg ggc ata tgt gag gag cat gga tcc 1409
 Lys Asp Leu Pro Ser Leu Ala Val Gly Ile Cys Glu Glu His Gly Ser
 430 435 440

gcc atg gcc ccc cag cac atc ctg gcc tct ggc tca act gct ggc ccc 1457
 Ala Met Ala Pro Gln His Ile Leu Ala Ser Gly Ser Thr Ala Gly Pro
 445 450 455

aag ctg tac ccc aag cta tac aca gat gtg cac aca cac aca cat aca 1505
 Lys Leu Tyr Pro Lys Leu Tyr Thr Asp Val His Thr His Thr His Thr
 460 465 470

cac acc tgc act cac acg ctc tca tgt gga ggg caa ggt tca tca aca 1553
 His Thr Cys Thr His Thr Leu Ser Cys Gly Gly Gln Gly Ser Ser Thr
 475 480 485

cca gca tgt cca cta tca gtg cta aat aca gcg aat ctc caa gca ctg 1601
 Pro Ala Cys Pro Leu Ser Val Leu Asn Thr Ala Asn Leu Gln Ala Leu
 490 495 500 505

tgt cct gag gta ggc ata tgg ggg cca agg caa cag gtt ggg aga att 1649
 Cys Pro Glu Val Gly Ile Trp Gly Pro Arg Gln Gln Val Gly Arg Ile
 510 515 520

gag aac aat gga gga aga gta tct tag ggtgccttat ggtggacact 1696
 Glu Asn Asn Gly Gly Arg Val Ser
 525 530

cacaaacttg gccatataga tgtatgtact accagatgaa cagccagcca gattcacaca 1756

cgcacatggt taaacgtgta aacgtgtgca caactgcaca cacaacctga gaaaccttca 1816

ggaggatttg tgggtgtgact ttgcagtgcac atgtagcgat ggctagttag aggaatctcc 1876

ctcatgtctt agtgggtcatg gccacttccc caccctgccc catctgtggt cctgcctggc 1936

cttgggtgtgc ttccgtgtgc cctgggtatc aggagcctat catcaacctg actgggggtga 1996

gcagtgcagc catgcctgga ggtttgagcc accctcccct tgctagagag aagggcctca 2056

atatttatat ttaagaaatg aaataatatt aataataatg taaggagggc tgggacacag 2116

ggactctggc ctccctggg gcctgggacc tgctggcct tgtgggttaca ttgggtaccc 2176

tcaactgtcca tggctgcctg gtctctgtaa ttttatatag agtttgagct gaagcctcgt 2236

atatttaatt tattttgtta aacaagaaaa aaaaaaaaaa a 2277

FIG. 2A

gcggccgcga cccaggtcc ggacaggccg ag	atg acg ccg agc ccc ctg ttg	53
	Met Thr Pro Ser Pro Leu Leu	
	1 5	
ctg ctc ctg ctg ccg ccg ctg ctg ctg ggg gcc ttc cca ccg gcc gcc	101	
Leu Leu Leu Leu Pro Pro Leu Leu Leu Gly Ala Phe Pro Pro Ala Ala		
10 15 20		
gcc gcc cga ggc ccc cca aag atg gcg gac aag gtg gtc cca cgg cag	149	
Ala Ala Arg Gly Pro Pro Lys Met Ala Asp Lys Val Val Pro Arg Gln		
25 30 35		
gtg gcc cgg ctg ggc cgc act gtg cgg ctg cag tgc cca gtg gag ggg	197	
Val Ala Arg Leu Gly Arg Thr Val Arg Leu Gln Cys Pro Val Glu Gly		
40 45 50 55		
gac ccg ccg ccg ctg acc atg tgg acc aag gat ggc cgc acc atc cac	245	
Asp Pro Pro Pro Leu Thr Met Trp Thr Lys Asp Gly Arg Thr Ile His		
60 65 70		
agc ggc tgg agc cgc ttc cgc gtg ctg ccg cag ggg ctg aag gtg aag	293	
Ser Gly Trp Ser Arg Phe Arg Val Leu Pro Gln Gly Leu Lys Val Lys		
75 80 85		
cag gtg gag cgg gag gat gcc ggc gtg tac gtg tgc aag gcc acc aac	341	
Gln Val Glu Arg Glu Asp Ala Gly Val Tyr Val Cys Lys Ala Thr Asn		
90 95 100		
ggc ttc ggc agc ctg agc gtc aac tac acc ctc gtc gtg ctg gat gac	389	
Gly Phe Gly Ser Leu Ser Val Asn Tyr Thr Leu Val Val Leu Asp Asp		
105 110 115		
att agc cca ggg aag gag agc ctg ggg ccc gac agc tcc tct ggg ggt	437	
Ile Ser Pro Gly Lys Glu Ser Leu Gly Pro Asp Ser Ser Ser Gly Gly		
120 125 130 135		
caa gag gac ccc gcc agc cag cag tgg gca cga ccg cgc ttc aca cag	485	
Gln Glu Asp Pro Ala Ser Gln Gln Trp Ala Arg Pro Arg Phe Thr Gln		
140 145 150		
ccc tcc aag atg agg cgc cgg gtg atc gca cgg ccc gtg ggt agc tcc	533	
Pro Ser Lys Met Arg Arg Arg Val Ile Ala Arg Pro Val Gly Ser Ser		
155 160 165		
gtg cgg ctc aag tgc gtg gcc agc ggg cac cct cgg ccc gac atc acg	581	
Val Arg Leu Lys Cys Val Ala Ser Gly His Pro Arg Pro Asp Ile Thr		
170 175 180		
tgg atg aag gac gac cag gcc ttg acg cgc cca gag gcc gct gag ccc	629	
Trp Met Lys Asp Asp Gln Ala Leu Thr Arg Pro Glu Ala Ala Glu Pro		
185 190 195		

FIG. 2B

agg aag aag aag tgg aca ctg agc ctg aag aac ctg cgg ccg gag gac	677
Arg Lys Lys Lys Trp Thr Leu Ser Leu Lys Asn Leu Arg Pro Glu Asp	
200 205 210 215	
agc ggc aaa tac acc tgc cgc gtg tgc aac cgc gcg ggc gcc atc aac	725
Ser Gly Lys Tyr Thr Cys Arg Val Ser Asn Arg Ala Gly Ala Ile Asn	
220 225 230	
gcc acc tac aag gtg gat gtg atc cag cgg acc cgt tcc aag ccc gtg	773
Ala Thr Tyr Lys Val Asp Val Ile Gln Arg Thr Arg Ser Lys Pro Val	
235 240 245	
ctc aca ggc acg cac ccc gtg aac acg acg gtg gac ttc ggg ggg acc	821
Leu Thr Gly Thr His Pro Val Asn Thr Thr Val Asp Phe Gly Gly Thr	
250 255 260	
acg tcc ttc cag tgc aag gtg cgc agc gac gtg aag ccg gtg atc cag	869
Thr Ser Phe Gln Cys Lys Val Arg Ser Asp Val Lys Pro Val Ile Gln	
265 270 275	
tgg ctg aag cgc gtg gag tac ggc gct gag ggc cgc cac aac tcc acc	917
Trp Leu Lys Arg Val Glu Tyr Gly Ala Glu Gly Arg His Asn Ser Thr	
280 285 290 295	
atc gat gtg ggc ggc cag aag ttt gtg gtg ctg ccc acg ggt gac gtg	965
Ile Asp Val Gly Gly Gln Lys Phe Val Val Leu Pro Thr Gly Asp Val	
300 305 310	
tgg tgc cgg ccc gac ggc tcc tac ctc aat aag ctg ctc atc acc cgt	1013
Trp Ser Arg Pro Asp Gly Ser Tyr Leu Asn Lys Leu Leu Ile Thr Arg	
315 320 325	
gcc cgc cag gac gat gcg ggc atg tac atc tgc ctt ggc gcc aac acc	1061
Ala Arg Gln Asp Asp Ala Gly Met Tyr Ile Cys Leu Gly Ala Asn Thr	
330 335 340	
atg ggc tac agc ttc cgc agc gcc ttc ctc acc gtg ctg cca gac cca	1109
Met Gly Tyr Ser Phe Arg Ser Ala Phe Leu Thr Val Leu Pro Asp Pro	
345 350 355	
aaa ccg cca ggg cca cct gtg gcc tcc tgc tcc tgc gcc act agc ctg	1157
Lys Pro Pro Gly Pro Pro Val Ala Ser Ser Ser Ser Ala Thr Ser Leu	
360 365 370 375	
ccg tgg ccc gtg gtc atc ggc atc cca gcc ggc gct gtc ttc atc ctg	1205
Pro Trp Pro Val Val Ile Gly Ile Pro Ala Gly Ala Val Phe Ile Leu	
380 385 390	
ggc acc ctg ctc ctg tgg ctt tgc cag gcc cag aag aag ccg tgc acc	1253
Gly Thr Leu Leu Leu Trp Leu Cys Gln Ala Gln Lys Lys Pro Cys Thr	
395 400 405	

FIG. 2C

ccc gcg cct gcc cct ccc ctg cct ggg cac cgc ccg ccg ggg acg gcc	1301
Pro Ala Pro Ala Pro Pro Leu Pro Gly His Arg Pro Pro Gly Thr Ala	
410 415 420	
cgc gac cgc agc gga gac aag gac ctt ccc tcg ttg gcc gcc ctc agc	1349
Arg Asp Arg Ser Gly Asp Lys Asp Leu Pro Ser Leu Ala Ala Leu Ser	
425 430 435	
gct ggc cct ggt gtg ggg ctg tgt gag gag cat ggg tct ccg gca gcc	1397
Ala Gly Pro Gly Val Gly Leu Cys Glu Glu His Gly Ser Pro Ala Ala	
440 445 450 455	
ccc cag cac tta ctg ggc cca ggc cca gtt gct ggc cct aag ttg tac	1445
Pro Gln His Leu Leu Gly Pro Gly Pro Val Ala Gly Pro Lys Leu Tyr	
460 465 470	
ccc ta	1450
Pro	

FIG. 3A

muFGFR-L	1MTRSPALLLLLLLGALPSAEAARGPP.....RMADKV	31
		. : :	
neFGFR-4	1	MGVQKDSRDIRWNRTTRPLALLLCGLLAFSALSCARTLPEGRKANLAELV	50
muFGFR-L	32	VPRQVARL...GRTVRLQCPVEGDPPPLTMWTKDGRTIHSGWSRFRVLPQ	78
		: . . : . .	
neFGFR-4	51	SEEEHFLLDPGNALRLFCDT.NQTTIVNWYTESTRLQHGG..RIRLTDT	97
muFGFR-L	79	GLKVKEVEAEDAGVYVCKATNGFGSLSVNYTLIIMDDISPG...KESPGP	125
		.: : . . . : ::: :. :.	
neFGFR-4	98	VLEIADVITYEDSGLYLC.VVPGTGHILRNFTISVVDSLASGDDDDDEDHGR	146
muFGFR-L	126	GGSSG..GQEDPASQQWARPRFTQPSKMRRRVIA RPVGSSVRLKCVASGH	173
		. :: : .. : ::. .. : : .. .	
neFGFR-4	147	EDSAGDMGEDPPYSTSYRAPFWSQPQRMDDKKLYAVPAGNTVKFRCPSAGN	196
muFGFR-L	174	PRPDIMWMKDDQTL...THLEASEHRKKKWTLSLKNLKPEDSGKYTCRVS	220
		:: . . : . . :...	
neFGFR-4	197	PTPGIRWLKNGREFGGEHRIGGIRLRHQHWSLVMESVVPSPDRGNYTCLVE	246
muFGFR-L	221	NKAGAINATYKVDVIQRTSRKPVLGTGTHPVNTTVDFGGTTSFQCKVRSDV	270
	 :: . : :	
neFGFR-4	247	NKFGSISYSYLLDVLEERSPHRPILQAGLPANTTAMLGSDVQFFCKVYSDA	296
muFGFR-L	271	KPVIQWLKRVEYGSEGRHNSTIDVGGQKFV.VLPTGDVWSRPDGSYLNKL	319
		. : . : . . .	
neFGFR-4	297	QPHIQWLKHIEV.....NGSRYGPDGVPFVQVLKTADI.....NSSEVEVL	337
muFGFR-L	320	LISRARQDDAGMYICLGANTMGYSFRSAFLTVPDPKPPGPPMASSSSST	369
		: : .. :. . : .	
neFGFR-4	338	YLHNVSFEDAGEYTCLAGNSIGLSYQSAWLTVLPEEDFAKEAEGPETRYT	387
muFGFR-L	370	SLPWPVVIGIPAGAVFILGTVLLWLCQTKKKPCAPASTLPVPGHRPPGTS	419
		:: : : :. . .	
neFGFR-4	388	D....IIIYTSGSLALLMAAVIVVLCRMQLP.....PTKTHLEPATV	425
muFGFR-L	420	RERSGDKDLPSLAVGICEEHGSAMAPQHILASGSTAGPKLYPKLYTDVHT	469
		. : : . .	
neFGFR-4	426	HKLSRFPLMRQFSLESSSSGKSSTSLVRVTRLSSSCTPMLPGVLEFDLPL	475

FIG. 3B

muFGFR-L	470	HTHTHTCTHTLSCGGQGSSTPACPLSVLNTANLQALCPEVGIWGPRQQVG	519
		: :	
neFGFR-4	476	DSKWEFPRERLVLGKPLGEGCFGQVVRAEAYGINKDQPDKAI.TVAIKIV	524
muFGFR-L	520	RIENNGGRVS*.....	530
		: . .	
neFGFR-4	525	KDKGTDKELSDLISEMELMKLMGKHKNIINLLGVCTQDGPLYMIVEYASK	574

FIG. 4

1 MTPSPLLLLLLLPPLLLGAFPPAAAARGPPKMADKVVPRQVARLGRTVRLQ 50
|| || | | | | | : | | | | | | | | | | | |
1 MTRSPALLL...LLL GALPSAEAARGPPRMADKVVPRQVARLGRTVRLQ. 46

51 CPVEGDPPPLTMWTKDGRTIHSGWSRFRVLPOGLKV KQVEREDAGVYVCK 100
| | | | | | | | | | | | | | | | | | : | | | | | | | |
47 CPVEGDPPPLTMWTKDGRTIHSGWSRFRVLPOGLKV KEVEAEADAGVYVCK 96

101 ATNGFGSLSVNYTLVVLDDISPGKESLGPDSSSSGGQEDPASQQWARPRFT 150
| | | | | | | | | | : : | | | | | | | | | | | | | | | |
97 ATNGFGSLSVNYTLIIMDDISPGKESPGPGGSSGGQEDPASQQWARPRFT 146

151 QPSKMRRRV IARPVGSSVRLKCVASGHPRPDITWMKDDQALTRPEAAEPR 200
| | | | | | | | | | | | | | | | | | | | | | | | . | |
147 QPSKMRRRV IARPVGSSVRLKCVASGHPRPDIMWMKDDQTLTHLEASEHR 196

201 KKKWTLSLKNLRPEDSGKYTCRVSNRAGAINATYKVDVIQRTRSKPVL TG 250
| | | | | | | | : | | | | | | | | : | | | | | | | | | | | |
197 KKKWTLSLKNLKPEDSGKYTCRVSNKAGAINATYKVDVIQRTRSKPVL TG 246

251 THPVNTTVDFGGTTSFQCKVRSDVKPVIQWLKRVEYGAEGRHNSTIDVGG 300
| | | | | | | | | | | | | | | | | | | | | | . | | | | | | | |
247 THPVNTTVDFGGTTSFQCKVRSDVKPVIQWLKRVEYGSEGRHNSTIDVGG 296

301 QKFVVLPTGDVWSRPDGSYLNKLLITRARQDDAGMYICLGANTMGYSFRS 350
| | | | | | | | | | | | | | | | | | . | | | | | | | | | | | |
297 QKFVVLPTGDVWSRPDGSYLNKLLISRARQDDAGMYICLGANTMGYSFRS 346

351 AFLTVLPDPKPPGPPVASSSSATS L PWPV VIGIPAGAVFILGTLLLWLCQ 400
| | | | | | | | | | . | | | | . | | | | | | | | | | | | | | | | |
347 AFLTVLPDPKPPGPPMASSSSSTS L PWPV VIGIPAGAVFILGTVLLWLCQ 396

401 AQKKPCTPAPAPPLPGHRPPGTARD RSGDKDLPSLAALSAGPGVGLCEEH 450
. | | | | | | . | | | | | | . : | | | | | | | | | | : | | | |
397 TKKKPCAPASTLPVP GHRPPGTSRERSGDKDLPSLA.....VGICEEH 439

451 GSPAAPQHLLGPGPVAGPKLYPKLYTDIHTHTHTSHSTHSHVEGKVHQHI 500
| | | | : | | | | | | | | | | : | | | | | . | | .
440 GSAMAPQHILASGSTAGPKLYPKLYTDVHTHTHTHTCTHTLSCGGQGSST 489

501 HYQC* 504

490 PACPLSVLNTANLQALCPEVGIWGPRQQVGRIENNGGRVS* 529

FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15




FIG. 16




FIG. 17




FIG. 18




FIG. 19




FIG. 20A




FIG. 20B




FIG. 21A




FIG. 21B




FIG. 21C




FIG. 21D




FIG. 22




FIG. 23





FIG. 24



SEQUENCE LISTING

<110> Saris, Christiaan M.
 Sharon, Mu X.
 Xia, Min
 Boone, Thomas Charles
 Covey, Todd

<120> Fibroblast Growth Factor Receptor-Like Molecules and
 Uses Thereof

<130> 99-513-A

<140>

<141>

<150> 60/191,379

<151> 2000-03-22

<160> 22

<170> PatentIn Ver. 2.0

<210> 1

<211> 2277

<212> DNA

<213> Mus musculus

<220>

<221> CDS

<222> (87)..(1673)

<220>

<221> sig_peptide

<222> (87)..(146)

<220>

<221> misc_feature

<222> (1208)..(1271)

<223> predicted transmembrane domain

<400> 1

gacctggggtc ttgcgggcct gagccctgag tggcgtccag tccagctccc agtgaccgcg 60

cccctgcttc aggtccgacc ggcgag atg acg cgg agc ccc gcg ctg ctg ctg 113

Met Thr Arg Ser Pro Ala Leu Leu Leu

1

5

ctg cta ttg ggg gcc ctc ccg tcg gct gag gcg gcg cga gga ccc cca 161

Leu Leu Leu Gly Ala Leu Pro Ser Ala Glu Ala Ala Arg Gly Pro Pro

10

15

20

25

aga atg gca gac aaa gtg gtc cca cgg cag gtg gcc cgc ctg ggc cgc 209

Arg Met Ala Asp Lys Val Val Pro Arg Gln Val Ala Arg Leu Gly Arg

30

35

40

act gtg cgg cta cag tgc cca gtg gag ggg gac cca cca ccg ttg acc 257

Thr Val Arg Leu Gln Cys Pro Val Gly Asp Pro Pro Pro Leu Thr

45

50

55

atg tgg acc aaa gat ggc cgc aca atc cac agt ggc tgg agc cgc ttc	305
Met Trp Thr Lys Asp Gly Arg Thr Ile His Ser Gly Trp Ser Arg Phe	
60 65 70	
cgt gtg ctg ccc cag ggt ctg aag gtg aag gag gtg gag gcc gag gat	353
Arg Val Leu Pro Gln Gly Leu Lys Val Lys Glu Val Glu Ala Glu Asp	
75 80 85	
gcc ggt gtt tat gtg tgc aag gcc acc aat ggc ttt ggc agc ctc agc	401
Ala Gly Val Tyr Val Cys Lys Ala Thr Asn Gly Phe Gly Ser Leu Ser	
90 95 100 105	
gtc aac tac act ctc atc atc atg gat gat att agt cca ggg aag gag	449
Val Asn Tyr Thr Leu Ile Ile Met Asp Asp Ile Ser Pro Gly Lys Glu	
110 115 120	
agc cct ggg cca ggt ggt tct tcg ggg ggc cag gag gac cca gcc agc	497
Ser Pro Gly Pro Gly Gly Ser Ser Gly Gly Gln Glu Asp Pro Ala Ser	
125 130 135	
cag cag tgg gca cgg cct cgc ttc aca cag ccc tcc aag atg agg cgc	545
Gln Gln Trp Ala Arg Pro Arg Phe Thr Gln Pro Ser Lys Met Arg Arg	
140 145 150	
cga gtg att gca cgg cct gtg ggt agc tct gtg cgg ctc aag tgt gtg	593
Arg Val Ile Ala Arg Pro Val Gly Ser Ser Val Arg Leu Lys Cys Val	
155 160 165	
gcc agt ggg cac cca cgg cca gac atc atg tgg atg aag gat gac cag	641
Ala Ser Gly His Pro Arg Pro Asp Ile Met Trp Met Lys Asp Asp Gln	
170 175 180 185	
acc ttg acg cat cta gag gct agt gaa cac aga aag aag aag tgg aca	689
Thr Leu Thr His Leu Glu Ala Ser Glu His Arg Lys Lys Lys Trp Thr	
190 195 200	
ctg agc ttg aag aac ctg aag cct gaa gac agt ggc aag tac acg tgc	737
Leu Ser Leu Lys Asn Leu Lys Pro Glu Asp Ser Gly Lys Tyr Thr Cys	
205 210 215	
cgt gta tct aac aag gcc ggt gcc atc aac gcc acc tac aaa gtg gat	785
Arg Val Ser Asn Lys Ala Gly Ala Ile Asn Ala Thr Tyr Lys Val Asp	
220 225 230	
gta atc cag cgg act cgt tcc aag cct gtg ctc aca ggg aca cac cct	833
Val Ile Gln Arg Thr Arg Ser Lys Pro Val Leu Thr Gly Thr His Pro	
235 240 245	
gtg aac aca acg gtg gac ttc ggt ggg aca acg tcc ttc cag tgc aag	881
Val Asn Thr Thr Val Asp Phe Gly Gly Thr Thr Ser Phe Gln Cys Lys	
250 255 260 265	
gtg cgc agt gac gtg aag cct gtg atc cag tgg ctg aag cgg gtg gag	929
Val Arg Ser Asp Val Lys Pro Val Ile Gln Trp Leu Lys Arg Val Glu	
270 275 280	
tac ggc tcc gag gga cgc cac aac tcc acc att gat gtg ggt ggc cag	977
Tyr Gly Ser Glu Gly Arg His Asn Ser Thr Ile Asp Val Gly Gly Gln	

285	290	295	
aag ttt gtg gtg ttg ccc acg ggt gat gtg tgg tca cgg cct gat ggc			1025
Lys Phe Val Val Leu Pro Thr Gly Asp Val Trp Ser Arg Pro Asp Gly			
300	305	310	
tcc tac ctc aac aag ctg ctc atc tct cgg gcc cgc cag gat gat gct			1073
Ser Tyr Leu Asn Lys Leu Leu Ile Ser Arg Ala Arg Gln Asp Asp Ala			
315	320	325	
ggc atg tac atc tgc cta ggt gca aat acc atg ggc tac agt ttc cgt			1121
Gly Met Tyr Ile Cys Leu Gly Ala Asn Thr Met Gly Tyr Ser Phe Arg			
330	335	340	345
agc gcc ttc ctc act gta tta cca gac ccc aaa cct cca ggg cct cct			1169
Ser Ala Phe Leu Thr Val Leu Pro Asp Pro Lys Pro Pro Gly Pro Pro			
350	355	360	
atg gct tct tca tcg tca tcc aca agc ctg cca tgg cct_gtg gtg atc			1217
Met Ala Ser Ser Ser Ser Ser Thr Ser Leu Pro Trp Pro Val Val Ile			
365	370	375	
ggc atc cca gct ggt gct gtc ttc atc cta ggc act gtg ctg ctc tgg			1265
Gly Ile Pro Ala Gly Ala Val Phe Ile Leu Gly Thr Val Leu Leu Trp			
380	385	390	
ctt tgc cag acc aag aag aag cca tgt gcc cca gca tct aca ctt cct			1313
Leu Cys Gln Thr Lys Lys Lys Pro Cys Ala Pro Ala Ser Thr Leu Pro			
395	400	405	
gtg cct ggg cat cgt ccc cca ggg aca tcc cga gaa cgc agt ggt gac			1361
Val Pro Gly His Arg Pro Pro Gly Thr Ser Arg Glu Arg Ser Gly Asp			
410	415	420	425
aag gac ctg ccc tca ttg gct gtg ggc ata tgt gag gag cat gga tcc			1409
Lys Asp Leu Pro Ser Leu Ala Val Gly Ile Cys Glu Glu His Gly Ser			
430	435	440	
gcc atg gcc ccc cag cac atc ctg gcc tct ggc tca act gct ggc ccc			1457
Ala Met Ala Pro Gln His Ile Leu Ala Ser Gly Ser Thr Ala Gly Pro			
445	450	455	
aag ctg tac ccc aag cta tac aca gat gtg cac aca cac aca cat aca			1505
Lys Leu Tyr Pro Lys Leu Tyr Thr Asp Val His Thr His Thr His Thr			
460	465	470	
cac acc tgc act cac acg ctc tca tgt gga ggg caa ggt tca tca aca			1553
His Thr Cys Thr His Thr Leu Ser Cys Gly Gly Gln Gly Ser Ser Thr			
475	480	485	
cca gca tgt cca cta tca gtg cta aat aca gcg aat ctc caa gca ctg			1601
Pro Ala Cys Pro Leu Ser Val Leu Asn Thr Ala Asn Leu Gln Ala Leu			
490	495	500	505
tgt cct gag gta ggc ata tgg ggg cca agg caa cag gtt ggg aga att			1649
Cys Pro Glu Val Gly Ile Trp Gly Pro Arg Gln Gln Val Gly Arg Ile			
510	515	520	
gag aac aat gga gga aga gta tct taggggtgcct tatgggtggac actcacaac			1703

Glu Asn Asn Gly Gly Arg Val Ser
525

ttggccatat agatgtatgt actaccagat gaacagccag ccagattcac acacgcacat 1763
gtttaaacgt gtaaactgt gcacaactgc acacacaacc tgagaaacct tcaggaggat 1823
ttgtggtgtg actttgcagt gacatgtagc gatggctagt tgaaggaatc tccctcatgt 1883
cttagtggtc atggccactt cccacccct gcccatctgt gttcctgcct ggccttggtg 1943
tgcttccgtg tgccctgggt atcaggagcc tatcatcaac ctgactgggg tgagcagtgc 2003
agccatgcct ggaggtttga gccaccctcc ccttgctaga gagaagggcc tcaatattta 2063
tatttaagaa atgaaataat attaataata atgtaaggag ggctgggaca cagggactct 2123
ggccttccct ggggcctggg acctgcctgg ccttggtggtt acattgggta ccctcactgt 2183
ccatggctgc ctggtctctg taattttata tagagtttga gctgaagcct cgtatattta 2243
atttattttg ttaaacaaga aaaaaaaaaa aaaa 2277

<210> 2

<211> 529

<212> PRT

<213> Mus musculus

<400> 2

Met	Thr	Arg	Ser	Pro	Ala	Leu	Leu	Leu	Leu	Leu	Gly	Ala	Leu	Pro		
1				5				10					15			
Ser	Ala	Glu	Ala	Ala	Arg	Gly	Pro	Pro	Arg	Met	Ala	Asp	Lys	Val	Val	
			20					25					30			
Pro	Arg	Gln	Val	Ala	Arg	Leu	Gly	Arg	Thr	Val	Arg	Leu	Gln	Cys	Pro	
		35					40					45				
Val	Glu	Gly	Asp	Pro	Pro	Pro	Leu	Thr	Met	Trp	Thr	Lys	Asp	Gly	Arg	
	50					55					60					
Thr	Ile	His	Ser	Gly	Trp	Ser	Arg	Phe	Arg	Val	Leu	Pro	Gln	Gly	Leu	
65					70					75					80	
Lys	Val	Lys	Glu	Val	Glu	Ala	Glu	Asp	Ala	Gly	Val	Tyr	Val	Cys	Lys	
			85						90					95		
Ala	Thr	Asn	Gly	Phe	Gly	Ser	Leu	Ser	Val	Asn	Tyr	Thr	Leu	Ile	Ile	
			100					105					110			
Met	Asp	Asp	Ile	Ser	Pro	Gly	Lys	Glu	Ser	Pro	Gly	Pro	Gly	Gly	Ser	
	115						120					125				
Ser	Gly	Gly	Gln	Glu	Asp	Pro	Ala	Ser	Gln	Gln	Trp	Ala	Arg	Pro	Arg	
	130					135					140					
Phe	Thr	Gln	Pro	Ser	Lys	Met	Arg	Arg	Arg	Val	Ile	Ala	Arg	Pro	Val	
145					150					155					160	

Gly Ser Ser Val Arg Leu Lys Cys Val Ala Ser Gly His Pro Arg Pro
 165 170 175
 Asp Ile Met Trp Met Lys Asp Asp Gln Thr Leu Thr His Leu Glu Ala
 180 185 190
 Ser Glu His Arg Lys Lys Lys Trp Thr Leu Ser Leu Lys Asn Leu Lys
 195 200 205
 Pro Glu Asp Ser Gly Lys Tyr Thr Cys Arg Val Ser Asn Lys Ala Gly
 210 215 220
 Ala Ile Asn Ala Thr Tyr Lys Val Asp Val Ile Gln Arg Thr Arg Ser
 225 230 235 240
 Lys Pro Val Leu Thr Gly Thr His Pro Val Asn Thr Thr Val Asp Phe
 245 250 255
 Gly Gly Thr Thr Ser Phe Gln Cys Lys Val Arg Ser Asp Val Lys Pro
 260 265 270
 Val Ile Gln Trp Leu Lys Arg Val Glu Tyr Gly Ser Glu Gly Arg His
 275 280 285
 Asn Ser Thr Ile Asp Val Gly Gly Gln Lys Phe Val Val Leu Pro Thr
 290 295 300
 Gly Asp Val Trp Ser Arg Pro Asp Gly Ser Tyr Leu Asn Lys Leu Leu
 305 310 315 320
 Ile Ser Arg Ala Arg Gln Asp Asp Ala Gly Met Tyr Ile Cys Leu Gly
 325 330 335
 Ala Asn Thr Met Gly Tyr Ser Phe Arg Ser Ala Phe Leu Thr Val Leu
 340 345 350
 Pro Asp Pro Lys Pro Pro Gly Pro Pro Met Ala Ser Ser Ser Ser Ser
 355 360 365
 Thr Ser Leu Pro Trp Pro Val Val Ile Gly Ile Pro Ala Gly Ala Val
 370 375 380
 Phe Ile Leu Gly Thr Val Leu Leu Trp Leu Cys Gln Thr Lys Lys Lys
 385 390 395 400
 Pro Cys Ala Pro Ala Ser Thr Leu Pro Val Pro Gly His Arg Pro Pro
 405 410 415
 Gly Thr Ser Arg Glu Arg Ser Gly Asp Lys Asp Leu Pro Ser Leu Ala
 420 425 430
 Val Gly Ile Cys Glu Glu His Gly Ser Ala Met Ala Pro Gln His Ile
 435 440 445
 Leu Ala Ser Gly Ser Thr Ala Gly Pro Lys Leu Tyr Pro Lys Leu Tyr
 450 455 460
 Thr Asp Val His Thr His Thr His Thr His Thr Cys Thr His Thr Leu

Ser

```
<220>  
<221> TRANSMEM  
<222> (355) .. (375)
```

6

Gly Lys Tyr Thr Cys Arg Val Ser Asn Lys Ala Gly Ala Ile Asn Ala
 195 200 205
 Thr Tyr Lys Val Asp Val Ile Gln Arg Thr Arg Ser Lys Pro Val Leu
 210 215 220
 Thr Gly Thr His Pro Val Asn Thr Thr Val Asp Phe Gly Gly Thr Thr
 225 230 235 240
 Ser Phe Gln Cys Lys Val Arg Ser Asp Val Lys Pro Val Ile Gln Trp
 245 250 255
 Leu Lys Arg Val Glu Tyr Gly Ser Glu Gly Arg His Asn Ser Thr Ile
 260 265 270
 Asp Val Gly Gly Gln Lys Phe Val Val Leu Pro Thr Gly Asp Val Trp
 275 280 285
 Ser Arg Pro Asp Gly Ser Tyr Leu Asn Lys Leu Leu Ile Ser Arg Ala
 290 295 300
 Arg Gln Asp Asp Ala Gly Met Tyr Ile Cys Leu Gly Ala Asn Thr Met
 305 310 315 320
 Gly Tyr Ser Phe Arg Ser Ala Phe Leu Thr Val Leu Pro Asp Pro Lys
 325 330 335
 Pro Pro Gly Pro Pro Met Ala Ser Ser Ser Ser Thr Ser Leu Pro
 340 345 350
 Trp Pro Val Val Ile Gly Ile Pro Ala Gly Ala Val Phe Ile Leu Gly
 355 360 365
 Thr Val Leu Leu Trp Leu Cys Gln Thr Lys Lys Lys Pro Cys Ala Pro
 370 375 380
 Ala Ser Thr Leu Pro Val Pro Gly His Arg Pro Pro Gly Thr Ser Arg
 385 390 395 400
 Glu Arg Ser Gly Asp Lys Asp Leu Pro Ser Leu Ala Val Gly Ile Cys
 405 410 415
 Glu Glu His Gly Ser Ala Met Ala Pro Gln His Ile Leu Ala Ser Gly
 420 425 430
 Ser Thr Ala Gly Pro Lys Leu Tyr Pro Lys Leu Tyr Thr Asp Val His
 435 440 445
 Thr His Thr His Thr His Thr Cys Thr His Thr Leu Ser Cys Gly Gly
 450 455 460
 Gln Gly Ser Ser Thr Pro Ala Cys Pro Leu Ser Val Leu Asn Thr Ala
 465 470 475 480
 Asn Leu Gln Ala Leu Cys Pro Glu Val Gly Ile Trp Gly Pro Arg Gln
 485 490 495
 Gln Val Gly Arg Ile Glu Asn Asn Gly Gly Arg Val Ser

500

505

```
<210> 4
<211> 1450
<212> DNA
<213> Homo sapiens
```

```
<220>
<221> CDS
<222> (33) .. (1448)
```

```
<220>
<221> sig_peptide
<222> (33) .. (104)
```

```
<220>  
<221> misc_feature  
<222> (1167) .. (1229)
```

<400> 4																
gcggccgcga	ccccaggtcc	ggacaggccg	ag	atg	acg	ccg	agc	ccc	ctg	ttg						53
				Met	Thr	Pro	Ser	Pro	Leu	Leu						
					1			5								
ctg	ctc	ctg	ctg	ccg	ccg	ctg	ctg	ctg	ggg	gcc	ttc	cca	ccg	gcc	gcc	101
Leu	Leu	Leu	Leu	Pro	Pro	Leu	Leu	Leu	Gly	Ala	Phe	Pro	Pro	Ala	Ala	
		10					15					20				
gcc	gcc	cga	ggc	ccc	cca	aag	atg	gcg	gac	aag	gtg	gtc	cca	cgg	cag	149
Ala	Ala	Arg	Gly	Pro	Pro	Lys	Met	Ala	Asp	Lys	Val	Val	Pro	Arg	Gln	
	25					30					35					
gtg	gcc	cgg	ctg	ggc	cgc	act	gtg	cgg	ctg	cag	tgc	cca	gtg	gag	ggg	197
Val	Ala	Arg	Leu	Gly	Arg	Thr	Val	Arg	Leu	Gln	Cys	Pro	Val	Glu	Gly	
40					45					50					55	
gac	ccg	ccg	ccg	ctg	acc	atg	tgg	acc	aag	gat	ggc	cgc	acc	atc	cac	245
Asp	Pro	Pro	Pro	Leu	Thr	Met	Trp	Thr	Lys	Asp	Gly	Arg	Thr	Ile	His	
				60					65					70		
agc	ggc	tgg	agc	cgc	ttc	cgc	gtg	ctg	ccg	cag	ggg	ctg	aag	gtg	aag	293
Ser	Gly	Trp	Ser	Arg	Phe	Arg	Val	Leu	Pro	Gln	Gly	Leu	Lys	Val	Lys	
			75				80						85			
cag	gtg	gag	cgg	gag	gat	gcc	ggc	gtg	tac	gtg	tgc	aag	gcc	acc	aac	341
Gln	Val	Glu	Arg	Glu	Asp	Ala	Gly	Val	Tyr	Val	Cys	Lys	Ala	Thr	Asn	
		90					95					100				
ggc	ttc	ggc	agc	ctg	agc	gtc	aac	tac	acc	ctc	gtc	gtg	ctg	gat	gac	389
Gly	Phe	Gly	Ser	Leu	Ser	Val	Asn	Tyr	Thr	Leu	Val	Val	Leu	Asp	Asp	
	105					110					115					
att	agc	cca	ggg	aag	gag	agc	ctg	ggg	ccc	gac	agc	tcc	tct	ggg	ggt	437
Ile	Ser	Pro	Gly	Lys	Glu	Ser	Leu	Gly	Pro	Asp	Ser	Ser	Ser	Gly	Gly	
120					125					130					135	
caa	gag	gac	ccc	gcc	agc	cag	cag	tgg	gca	cga	ccg	cgc	ttc	aca	cag	485
Gln	Glu	Asp	Pro	Ala	Ser	Gln	Gln	Trp	Ala	Arg	Pro	Arg	Phe	Thr	Gln	

140								145					150					
ccc	tcc	aag	atg	agg	cgc	cgg	gtg	atc	gca	cgg	ccc	gtg	ggt	agc	tcc	533		
Pro	Ser	Lys	Met	Arg	Arg	Arg	Val	Ile	Ala	Arg	Pro	Val	Gly	Ser	Ser			
			155					160					165					
gtg	cgg	ctc	aag	tgc	gtg	gcc	agc	ggg	cac	cct	cgg	ccc	gac	atc	acg	581		
Val	Arg	Leu	Lys	Cys	Val	Ala	Ser	Gly	His	Pro	Arg	Pro	Asp	Ile	Thr			
			170					175					180					
tgg	atg	aag	gac	gac	cag	gcc	ttg	acg	cgc	cca	gag	gcc	gct	gag	ccc	629		
Trp	Met	Lys	Asp	Asp	Gln	Ala	Leu	Thr	Arg	Pro	Glu	Ala	Ala	Glu	Pro			
			185					190					195					
agg	aag	aag	aag	tgg	aca	ctg	agc	ctg	aag	aac	ctg	cgg	ccg	gag	gac	677		
Arg	Lys	Lys	Lys	Trp	Thr	Leu	Ser	Leu	Lys	Asn	Leu	Arg	Pro	Glu	Asp			
						205					210				215			
agc	ggc	aaa	tac	acc	tgc	cgc	gtg	tcg	aac	cgc	gcg	ggc	gcc	atc	aac	725		
Ser	Gly	Lys	Tyr	Thr	Cys	Arg	Val	Ser	Asn	Arg	Ala	Gly	Ala	Ile	Asn			
				220					225						230			
gcc	acc	tac	aag	gtg	gat	gtg	atc	cag	cgg	acc	cgt	tcc	aag	ccc	gtg	773		
Ala	Thr	Tyr	Lys	Val	Asp	Val	Ile	Gln	Arg	Thr	Arg	Ser	Lys	Pro	Val			
			235					240					245					
ctc	aca	ggc	acg	cac	ccc	gtg	aac	acg	acg	gtg	gac	ttc	ggg	ggg	acc	821		
Leu	Thr	Gly	Thr	His	Pro	Val	Asn	Thr	Thr	Val	Asp	Phe	Gly	Gly	Thr			
			250					255				260						
acg	tcc	ttc	cag	tgc	aag	gtg	cgc	agc	gac	gtg	aag	ccg	gtg	atc	cag	869		
Thr	Ser	Phe	Gln	Cys	Lys	Val	Arg	Ser	Asp	Val	Lys	Pro	Val	Ile	Gln			
			265					270				275						
tgg	ctg	aag	cgc	gtg	gag	tac	ggc	gct	gag	ggc	cgc	cac	aac	tcc	acc	917		
Trp	Leu	Lys	Arg	Val	Glu	Tyr	Gly	Ala	Glu	Gly	Arg	His	Asn	Ser	Thr			
					285					290					295			
atc	gat	gtg	ggc	ggc	cag	aag	ttt	gtg	gtg	ctg	ccc	acg	ggt	gac	gtg	965		
Ile	Asp	Val	Gly	Gly	Gln	Lys	Phe	Val	Val	Leu	Pro	Thr	Gly	Asp	Val			
				300						305					310			
tgg	tcg	cgg	ccc	gac	ggc	tcc	tac	ctc	aat	aag	ctg	ctc	atc	acc	cgt	1013		
Trp	Ser	Arg	Pro	Asp	Gly	Ser	Tyr	Leu	Asn	Lys	Leu	Leu	Ile	Thr	Arg			
			315					320					325					
gcc	cgc	cag	gac	gat	gcg	ggc	atg	tac	atc	tgc	ctt	ggc	gcc	aac	acc	1061		
Ala	Arg	Gln	Asp	Asp	Ala	Gly	Met	Tyr	Ile	Cys	Leu	Gly	Ala	Asn	Thr			
			330					335					340					
atg	ggc	tac	agc	ttc	cgc	agc	gcc	ttc	ctc	acc	gtg	ctg	cca	gac	cca	1109		
Met	Gly	Tyr	Ser	Phe	Arg	Ser	Ala	Phe	Leu	Thr	Val	Leu	Pro	Asp	Pro			
			345					350					355					
aaa	ccg	cca	ggg	cca	cct	gtg	gcc	tcc	tcg	tcc	tcg	gcc	act	agc	ctg	1157		
Lys	Pro	Pro	Gly	Pro	Pro	Val	Ala	Ser	Ser	Ser	Ser	Ala	Thr	Ser	Leu			
					365					370					375			
ccg	tgg	ccc	gtg	gtc	atc	ggc	atc	cca	gcc	ggc	gct	gtc	ttc	atc	ctg	1205		

Pro Trp Pro Val Val Ile Gly Ile Pro Ala Gly Ala Val Phe Ile Leu
 380 385 390

ggc acc ctg ctc ctg tgg ctt tgc cag gcc cag aag aag ccg tgc acc 1253
 Gly Thr Leu Leu Leu Trp Leu Cys Gln Ala Gln Lys Lys Pro Cys Thr
 395 400 405

ccc gcg cct gcc cct ccc ctg cct ggg cac cgc ccg ccg ggg acg gcc 1301
 Pro Ala Pro Ala Pro Pro Leu Pro Gly His Arg Pro Pro Gly Thr Ala
 410 415 420

cgc gac cgc agc gga gac aag gac ctt ccc tcg ttg gcc gcc ctc agc 1349
 Arg Asp Arg Ser Gly Asp Lys Asp Leu Pro Ser Leu Ala Ala Leu Ser
 425 430 435

gct ggc cct ggt gtg ggg ctg tgt gag gag cat ggg tct ccg gca gcc 1397
 Ala Gly Pro Gly Val Gly Leu Cys Glu Glu His Gly Ser Pro Ala Ala
 440 445 450 455

ccc cag cac tta ctg ggc cca ggc cca gtt gct ggc cct aag ttg tac 1445
 Pro Gln His Leu Leu Gly Pro Gly Pro Val Ala Gly Pro Lys Leu Tyr
 460 465 470

ccc ta 1450
 Pro

<210> 5
 <211> 472
 <212> PRT
 <213> Homo sapiens

<400> 5
 Met Thr Pro Ser Pro Leu Leu Leu Leu Leu Leu Pro Pro Leu Leu Leu
 1 5 10 15

Gly Ala Phe Pro Pro Ala Ala Ala Ala Arg Gly Pro Pro Lys Met Ala
 20 25 30

Asp Lys Val Val Pro Arg Gln Val Ala Arg Leu Gly Arg Thr Val Arg
 35 40 45

Leu Gln Cys Pro Val Glu Gly Asp Pro Pro Pro Leu Thr Met Trp Thr
 50 55 60

Lys Asp Gly Arg Thr Ile His Ser Gly Trp Ser Arg Phe Arg Val Leu
 65 70 75 80

Pro Gln Gly Leu Lys Val Lys Gln Val Glu Arg Glu Asp Ala Gly Val
 85 90 95

Tyr Val Cys Lys Ala Thr Asn Gly Phe Gly Ser Leu Ser Val Asn Tyr
 100 105 110

Thr Leu Val Val Leu Asp Asp Ile Ser Pro Gly Lys Glu Ser Leu Gly
 115 120 125

Pro Asp Ser Ser Ser Gly Gly Gln Glu Asp Pro Ala Ser Gln Gln Trp
 130 135 140

Ala Arg Pro Arg Phe Thr Gln Pro Ser Lys Met Arg Arg Arg Val Ile
 145 150 155 160
 Ala Arg Pro Val Gly Ser Ser Val Arg Leu Lys Cys Val Ala Ser Gly
 165 170 175
 His Pro Arg Pro Asp Ile Thr Trp Met Lys Asp Asp Gln Ala Leu Thr
 180 185 190
 Arg Pro Glu Ala Ala Glu Pro Arg Lys Lys Lys Trp Thr Leu Ser Leu
 195 200 205
 Lys Asn Leu Arg Pro Glu Asp Ser Gly Lys Tyr Thr Cys Arg Val Ser
 210 215 220
 Asn Arg Ala Gly Ala Ile Asn Ala Thr Tyr Lys Val Asp Val Ile Gln
 225 230 235 240
 Arg Thr Arg Ser Lys Pro Val Leu Thr Gly Thr His Pro Val Asn Thr
 245 250 255
 Thr Val Asp Phe Gly Gly Thr Thr Ser Phe Gln Cys Lys Val Arg Ser
 260 265 270
 Asp Val Lys Pro Val Ile Gln Trp Leu Lys Arg Val Glu Tyr Gly Ala
 275 280 285
 Glu Gly Arg His Asn Ser Thr Ile Asp Val Gly Gly Gln Lys Phe Val
 290 295 300
 Val Leu Pro Thr Gly Asp Val Trp Ser Arg Pro Asp Gly Ser Tyr Leu
 305 310 315 320
 Asn Lys Leu Leu Ile Thr Arg Ala Arg Gln Asp Asp Ala Gly Met Tyr
 325 330 335
 Ile Cys Leu Gly Ala Asn Thr Met Gly Tyr Ser Phe Arg Ser Ala Phe
 340 345 350
 Leu Thr Val Leu Pro Asp Pro Lys Pro Pro Gly Pro Pro Val Ala Ser
 355 360 365
 Ser Ser Ser Ala Thr Ser Leu Pro Trp Pro Val Val Ile Gly Ile Pro
 370 375 380
 Ala Gly Ala Val Phe Ile Leu Gly Thr Leu Leu Leu Trp Leu Cys Gln
 385 390 395 400
 Ala Gln Lys Lys Pro Cys Thr Pro Ala Pro Ala Pro Pro Leu Pro Gly
 405 410 415
 His Arg Pro Pro Gly Thr Ala Arg Asp Arg Ser Gly Asp Lys Asp Leu
 420 425 430
 Pro Ser Leu Ala Ala Leu Ser Ala Gly Pro Gly Val Gly Leu Cys Glu
 435 440 445
 Glu His Gly Ser Pro Ala Ala Pro Gln His Leu Leu Gly Pro Gly Pro

450 455 460
 Val Ala Gly Pro Lys Leu Tyr Pro
 465 470

 <210> 6
 <211> 448
 <212> PRT
 <213> Homo sapiens

 <220>
 <221> TRANSMEM
 <222> (355)..(375)

 <400> 6
 Ala Arg Gly Pro Pro Lys Met Ala Asp Lys Val Val Pro Arg Gln Val
 1 5 10 15
 Ala Arg Leu Gly Arg Thr Val Arg Leu Gln Cys Pro Val Glu Gly Asp
 20 25 30
 Pro Pro Pro Leu Thr Met Trp Thr Lys Asp Gly Arg Thr Ile His Ser
 35 40 45
 Gly Trp Ser Arg Phe Arg Val Leu Pro Gln Gly Leu Lys Val Lys Gln
 50 55 60
 Val Glu Arg Glu Asp Ala Gly Val Tyr Val Cys Lys Ala Thr Asn Gly
 65 70 75 80
 Phe Gly Ser Leu Ser Val Asn Tyr Thr Leu Val Val Leu Asp Asp Ile
 85 90 95
 Ser Pro Gly Lys Glu Ser Leu Gly Pro Asp Ser Ser Ser Gly Gly Gln
 100 105 110
 Glu Asp Pro Ala Ser Gln Gln Trp Ala Arg Pro Arg Phe Thr Gln Pro
 115 120 125
 Ser Lys Met Arg Arg Arg Val Ile Ala Arg Pro Val Gly Ser Ser Val
 130 135 140
 Arg Leu Lys Cys Val Ala Ser Gly His Pro Arg Pro Asp Ile Thr Trp
 145 150 155 160
 Met Lys Asp Asp Gln Ala Leu Thr Arg Pro Glu Ala Ala Glu Pro Arg
 165 170 175
 Lys Lys Lys Trp Thr Leu Ser Leu Lys Asn Leu Arg Pro Glu Asp Ser
 180 185 190
 Gly Lys Tyr Thr Cys Arg Val Ser Asn Arg Ala Gly Ala Ile Asn Ala
 195 200 205
 Thr Tyr Lys Val Asp Val Ile Gln Arg Thr Arg Ser Lys Pro Val Leu
 210 215 220
 Thr Gly Thr His Pro Val Asn Thr Thr Val Asp Phe Gly Gly Thr Thr

225					230					235					240		
Ser	Phe	Gln	Cys	Lys	Val	Arg	Ser	Asp	Val	Lys	Pro	Val	Ile	Gln	Trp		
				245					250					255			
Leu	Lys	Arg	Val	Glu	Tyr	Gly	Ala	Glu	Gly	Arg	His	Asn	Ser	Thr	Ile		
				260					265					270			
Asp	Val	Gly	Gly	Gln	Lys	Phe	Val	Val	Leu	Pro	Thr	Gly	Asp	Val	Trp		
				275					280					285			
Ser	Arg	Pro	Asp	Gly	Ser	Tyr	Leu	Asn	Lys	Leu	Leu	Ile	Thr	Arg	Ala		
				290					295					300			
Arg	Gln	Asp	Asp	Ala	Gly	Met	Tyr	Ile	Cys	Leu	Gly	Ala	Asn	Thr	Met		
305					310					315					320		
Gly	Tyr	Ser	Phe	Arg	Ser	Ala	Phe	Leu	Thr	Val	Leu	Pro	Asp	Pro	Lys		
				325					330					335			
Pro	Pro	Gly	Pro	Pro	Val	Ala	Ser	Ser	Ser	Ser	Ala	Thr	Ser	Leu	Pro		
				340					345					350			
Trp	Pro	Val	Val	Ile	Gly	Ile	Pro	Ala	Gly	Ala	Val	Phe	Ile	Leu	Gly		
				355					360					365			
Thr	Leu	Leu	Leu	Trp	Leu	Cys	Gln	Ala	Gln	Lys	Lys	Pro	Cys	Thr	Pro		
				370					375					380			
Ala	Pro	Ala	Pro	Pro	Leu	Pro	Gly	His	Arg	Pro	Pro	Gly	Thr	Ala	Arg		
385					390					395					400		
Asp	Arg	Ser	Gly	Asp	Lys	Asp	Leu	Pro	Ser	Leu	Ala	Ala	Leu	Ser	Ala		
				405					410					415			
Gly	Pro	Gly	Val	Gly	Leu	Cys	Glu	Glu	His	Gly	Ser	Pro	Ala	Ala	Pro		
				420					425					430			
Gln	His	Leu	Leu	Gly	Pro	Gly	Pro	Val	Ala	Gly	Pro	Lys	Leu	Tyr	Pro		
				435					440					445			

```
<210> 7
<211> 574
<212> PRT
<213> Pleurodeles waltlii
```

```
<400> 7
Met Gly Val Gln Lys Asp Ser Arg Asp Ile Arg Trp Asn Arg Thr Thr
  1                               5                10                15

Arg Pro Leu Ala Leu Leu Leu Cys Gly Leu Leu Ala Phe Ser Ala Leu
          20                          25                      30

Ser Cys Ala Arg Thr Leu Pro Glu Gly Arg Lys Ala Asn Leu Ala Glu
          35                          40                      45

Leu Val Ser Glu Glu Glu Glu His Phe Leu Leu Asp Pro Gly Asn Ala
  50                          55                60
```


Leu Arg Leu Phe Cys Asp Thr Asn Gln Thr Thr Ile Val Asn Trp Tyr
 65 70 75 80
 Thr Glu Ser Thr Arg Leu Gln His Gly Gly Arg Ile Arg Leu Thr Asp
 85 90 95
 Thr Val Leu Glu Ile Ala Asp Val Thr Tyr Glu Asp Ser Gly Leu Tyr
 100 105 110
 Leu Cys Val Val Pro Gly Thr Gly His Ile Leu Arg Asn Phe Thr Ile
 115 120 125
 Ser Val Val Asp Ser Leu Ala Ser Gly Asp Asp Asp Asp Glu Asp His
 130 135 140
 Gly Arg Glu Asp Ser Ala Gly Asp Met Gly Glu Asp Pro Pro Tyr Ser
 145 150 155 160
 Thr Ser Tyr Arg Ala Pro Phe Trp Ser Gln Pro Gln Arg Met Asp Lys
 165 170 175
 Lys Leu Tyr Ala Val Pro Ala Gly Asn Thr Val Lys Phe Arg Cys Pro
 180 185 190
 Ser Ala Gly Asn Pro Thr Pro Gly Ile Arg Trp Leu Lys Asn Gly Arg
 195 200 205
 Glu Phe Gly Gly Glu His Arg Ile Gly Gly Ile Arg Leu Arg His Gln
 210 215 220
 His Trp Ser Leu Val Met Glu Ser Val Val Pro Ser Asp Arg Gly Asn
 225 230 235 240
 Tyr Thr Cys Leu Val Glu Asn Lys Phe Gly Ser Ile Ser Tyr Ser Tyr
 245 250 255
 Leu Leu Asp Val Leu Glu Arg Ser Pro His Arg Pro Ile Leu Gln Ala
 260 265 270
 Gly Leu Pro Ala Asn Thr Thr Ala Met Leu Gly Ser Asp Val Gln Phe
 275 280 285
 Phe Cys Lys Val Tyr Ser Asp Ala Gln Pro His Ile Gln Trp Leu Lys
 290 295 300
 His Ile Glu Val Asn Gly Ser Arg Tyr Gly Pro Asp Gly Val Pro Phe
 305 310 315 320
 Val Gln Val Leu Lys Thr Ala Asp Ile Asn Ser Ser Glu Val Glu Val
 325 330 335
 Leu Tyr Leu His Asn Val Ser Phe Glu Asp Ala Gly Glu Tyr Thr Cys
 340 345 350
 Leu Ala Gly Asn Ser Ile Gly Leu Ser Tyr Gln Ser Ala Trp Leu Thr
 355 360 365
 Val Leu Pro Glu Glu Asp Phe Ala Lys Glu Ala Glu Gly Pro Glu Thr

370 375 380
 Arg Tyr Thr Asp Ile Ile Ile Tyr Thr Ser Gly Ser Leu Ala Leu Leu
 385 390 395 400
 Met Ala Ala Val Ile Val Val Leu Cys Arg Met Gln Leu Pro Pro Thr
 405 410 415
 Lys Thr His Leu Glu Pro Ala Thr Val His Lys Leu Ser Arg Phe Pro
 420 425 430
 Leu Met Arg Gln Phe Ser Leu Glu Ser Ser Ser Ser Gly Lys Ser Ser
 435 440 445
 Thr Ser Leu Val Arg Val Thr Arg Leu Ser Ser Ser Cys Thr Pro Met
 450 455 460
 Leu Pro Gly Val Leu Glu Phe Asp Leu Pro Leu Asp Ser Lys Trp Glu
 465 470 475 480
 Phe Pro Arg Glu Arg Leu Val Leu Gly Lys Pro Leu Gly Glu Gly Cys
 485 490 495
 Phe Gly Gln Val Val Arg Ala Glu Ala Tyr Gly Ile Asn Lys Asp Gln
 500 505 510
 Pro Asp Lys Ala Ile Thr Val Ala Ile Lys Ile Val Lys Asp Lys Gly
 515 520 525
 Thr Asp Lys Glu Leu Ser Asp Leu Ile Ser Glu Met Glu Leu Met Lys
 530 535 540
 Leu Met Gly Lys His Lys Asn Ile Ile Asn Leu Leu Gly Val Cys Thr
 545 550 555 560
 Gln Asp Gly Pro Leu Tyr Met Ile Val Glu Tyr Ala Ser Lys
 565 570

<210> 8

<211> 504

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: virtual human
 FGFR-L amino acid sequence comprising residues
 1-472 of SEQ ID NO: 5 and residues 473-504 of
 GenBank accession no. AJ277437

<400> 8

Met Thr Pro Ser Pro Leu Leu Leu Leu Leu Pro Pro Leu Leu Leu
 1 5 10 15
 Gly Ala Phe Pro Pro Ala Ala Ala Ala Arg Gly Pro Pro Lys Met Ala
 20 25 30
 Asp Lys Val Val Pro Arg Gln Val Ala Arg Leu Gly Arg Thr Val Arg
 35 40 45

Leu Gln Cys Pro Val Glu Gly Asp Pro Pro Pro Leu Thr Met Trp Thr
 50 55 60
 Lys Asp Gly Arg Thr Ile His Ser Gly Trp Ser Arg Phe Arg Val Leu
 65 70 75 80
 Pro Gln Gly Leu Lys Val Lys Gln Val Glu Arg Glu Asp Ala Gly Val
 85 90 95
 Tyr Val Cys Lys Ala Thr Asn Gly Phe Gly Ser Leu Ser Val Asn Tyr
 100 105 110
 Thr Leu Val Val Leu Asp Asp Ile Ser Pro Gly Lys Glu Ser Leu Gly
 115 120 125
 Pro Asp Ser Ser Ser Gly Gly Gln Glu Asp Pro Ala Ser Gln Gln Trp
 130 135 140
 Ala Arg Pro Arg Phe Thr Gln Pro Ser Lys Met Arg Arg Arg Val Ile
 145 150 155 160
 Ala Arg Pro Val Gly Ser Ser Val Arg Leu Lys Cys Val Ala Ser Gly
 165 170 175
 His Pro Arg Pro Asp Ile Thr Trp Met Lys Asp Asp Gln Ala Leu Thr
 180 185 190
 Arg Pro Glu Ala Ala Glu Pro Arg Lys Lys Lys Trp Thr Leu Ser Leu
 195 200 205
 Lys Asn Leu Arg Pro Glu Asp Ser Gly Lys Tyr Thr Cys Arg Val Ser
 210 215 220
 Asn Arg Ala Gly Ala Ile Asn Ala Thr Tyr Lys Val Asp Val Ile Gln
 225 230 235 240
 Arg Thr Arg Ser Lys Pro Val Leu Thr Gly Thr His Pro Val Asn Thr
 245 250 255
 Thr Val Asp Phe Gly Gly Thr Thr Ser Phe Gln Cys Lys Val Arg Ser
 260 265 270
 Asp Val Lys Pro Val Ile Gln Trp Leu Lys Arg Val Glu Tyr Gly Ala
 275 280 285
 Glu Gly Arg His Asn Ser Thr Ile Asp Val Gly Gly Gln Lys Phe Val
 290 295 300
 Val Leu Pro Thr Gly Asp Val Trp Ser Arg Pro Asp Gly Ser Tyr Leu
 305 310 315 320
 Asn Lys Leu Leu Ile Thr Arg Ala Arg Gln Asp Asp Ala Gly Met Tyr
 325 330 335
 Ile Cys Leu Gly Ala Asn Thr Met Gly Tyr Ser Phe Arg Ser Ala Phe
 340 345 350
 Leu Thr Val Leu Pro Asp Pro Lys Pro Pro Gly Pro Pro Val Ala Ser

355		360		365	
Ser	Ser	Ser	Ala	Thr	Ser
370		375		380	
Ala	Gly	Ala	Val	Phe	Ile
385		390		395	
Ala	Gln	Lys	Lys	Pro	Cys
		405		410	
His	Arg	Pro	Pro	Gly	Thr
		420		425	
Pro	Ser	Leu	Ala	Ala	Leu
		435		440	
Glu	His	Gly	Ser	Pro	Ala
450				455	
Val	Ala	Gly	Pro	Lys	Leu
465				470	
His	Thr	His	Thr	His	Ser
		485		490	
His	Gln	His	Ile	His	Tyr
		500			

<210> 9
 <211> 11
 <212> PRT
 <213> Human immunodeficiency virus type 1

<400> 9
 Tyr Gly Arg Lys Lys Arg Arg Gln Arg Arg Arg
 1 5 10

<210> 10
 <211> 15
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: internalizing
 domain derived from HIV tat protein

<400> 10
 Gly Gly Gly Gly Tyr Gly Arg Lys Lys Arg Arg Gln Arg Arg Arg
 1 5 10 15

<210> 11
 <211> 20
 <212> PRT
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: predicted
signal peptide of murine FGFR-L polypeptide

<400> 11

Met Thr Arg Ser Pro Ala Leu Leu Leu Leu Leu Gly Ala Leu Pro
1 5 10 15Ser Ala Glu Ala
20

<210> 12

<211> 25

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: predicted
transmembrane domain for murine FRL polypeptide

<400> 12

Leu Pro Trp Pro Val Val Ile Gly Ile Pro Ala Gly Ala Val Phe Ile
1 5 10 15Leu Gly Thr Val Leu Leu Trp Leu Cys
20 25

<210> 13

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide; PCR primer

<400> 13

cgctgaccat gtggaccaag gatg

24

<210> 14

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide; PCR primer

<400> 14

cttgacccca gaaggagctg tcgg

24

<210> 15

<211> 504

<212> PRT

<213> Homo sapiens

<400> 15
 Met Thr Pro Ser Pro Leu Leu Leu Leu Leu Leu Pro Pro Leu Leu Leu
 1 5 10 15
 Gly Ala Phe Pro Pro Ala Ala Ala Ala Arg Gly Pro Pro Lys Met Ala
 20 25 30
 Asp Lys Val Val Pro Arg Gln Val Ala Arg Leu Gly Arg Thr Val Arg
 35 40 45
 Leu Gln Cys Pro Val Glu Gly Asp Pro Pro Pro Leu Thr Met Trp Thr
 50 55 60
 Lys Asp Gly Arg Thr Ile His Ser Gly Trp Ser Arg Phe Arg Val Leu
 65 70 75 80
 Pro Gln Gly Leu Lys Val Lys Gln Val Glu Arg Glu Asp Ala Gly Val
 85 90 95
 Tyr Val Cys Lys Ala Thr Asn Gly Phe Gly Ser Leu Ser Val Asn Tyr
 100 105 110
 Thr Leu Val Val Leu Asp Asp Ile Ser Pro Gly Lys Glu Ser Leu Gly
 115 120 125
 Pro Asp Ser Ser Ser Gly Gly Gln Glu Asp Pro Ala Ser Gln Gln Trp
 130 135 140
 Ala Arg Pro Arg Phe Thr Gln Pro Ser Lys Met Arg Arg Arg Val Ile
 145 150 155 160
 Ala Arg Pro Val Gly Ser Ser Val Arg Leu Lys Cys Val Ala Ser Gly
 165 170 175
 His Pro Arg Pro Asp Ile Thr Trp Met Lys Asp Asp Gln Ala Leu Thr
 180 185 190
 Arg Pro Glu Ala Ala Glu Pro Arg Lys Lys Lys Trp Thr Leu Ser Leu
 195 200 205
 Lys Asn Leu Arg Pro Glu Asp Ser Gly Lys Tyr Thr Cys Arg Val Ser
 210 215 220
 Asn Arg Ala Gly Ala Ile Asn Ala Thr Tyr Lys Val Asp Val Ile Gln
 225 230 235 240
 Arg Thr Arg Ser Lys Pro Val Leu Thr Gly Thr His Pro Val Asn Thr
 245 250 255
 Thr Val Asp Phe Gly Gly Thr Thr Ser Phe Gln Cys Lys Val Arg Ser
 260 265 270
 Asp Val Lys Pro Val Ile Gln Trp Leu Lys Arg Val Glu Tyr Gly Ala
 275 280 285
 Glu Gly Arg His Asn Ser Thr Ile Asp Val Gly Gly Gln Lys Phe Val
 290 295 300

Val Leu Pro Thr Gly Asp Val Trp Ser Arg Pro Asp Gly Ser Tyr Leu
 305 310 315 320
 Asn Lys Leu Leu Ile Thr Arg Ala Arg Gln Asp Asp Ala Gly Met Tyr
 325 330 335
 Ile Cys Leu Gly Ala Asn Thr Met Gly Tyr Ser Phe Arg Ser Ala Phe
 340 345 350
 Leu Thr Val Leu Pro Asp Pro Lys Pro Pro Gly Pro Pro Val Ala Ser
 355 360 365
 Ser Ser Ser Ala Thr Ser Leu Pro Trp Pro Val Val Ile Gly Ile Pro
 370 375 380
 Ala Gly Ala Val Phe Ile Leu Gly Thr Leu Leu Leu Trp Leu Cys Gln
 385 390 395 400
 Ala Gln Lys Lys Pro Cys Thr Pro Ala Pro Ala Pro Pro Leu Pro Gly
 405 410 415
 His Arg Pro Pro Gly Thr Ala Arg Asp Arg Ser Gly Asp Lys Asp Leu
 420 425 430
 Pro Ser Leu Ala Ala Leu Ser Ala Gly Pro Gly Val Gly Leu Cys Glu
 435 440 445
 Glu His Gly Ser Pro Ala Ala Pro Gln His Leu Leu Gly Pro Gly Pro
 450 455 460
 Val Ala Gly Pro Lys Leu Tyr Pro Lys Leu Tyr Thr Asp Ile His Thr
 465 470 475 480
 His Thr His Thr His Ser His Thr His Ser His Val Glu Gly Lys Val
 485 490 495
 His Gln His Ile His Tyr Gln Cys
 500

<210> 16
 <211> 3112
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (25)..(1536)

<400> 16
 gaccccaggt ccggacaggc cgag atg acg ccg agc ccc ctg ttg ctg ctc 51
 Met Thr Pro Ser Pro Leu Leu Leu Leu
 1 5
 ctg ctg ccg ccg ctg ctg ctg ggg gcc ttc cca ccg gcc gcc gcc gcc 99
 Leu Leu Pro Pro Leu Leu Leu Gly Ala Phe Pro Pro Ala Ala Ala Ala
 10 15 20 25
 cga ggc ccc cca aag atg gcg gac aag gtg gtc cca ccg cag gtg gcc 147

Arg Gly Pro Pro Lys Met Ala Asp Lys Val Val Pro Arg Gln Val Ala	
30 35 40	
cgg ctg ggc cgc act gtg cgg ctg cag tgc cca gtg gag ggg gac ccg	195
Arg Leu Gly Arg Thr Val Arg Leu Gln Cys Pro Val Glu Gly Asp Pro	
45 50 55	
ccg ccg ctg acc atg tgg acc aag gat ggc cgc acc atc cac agc ggc	243
Pro Pro Leu Thr Met Trp Thr Lys Asp Gly Arg Thr Ile His Ser Gly	
60 65 70	
tgg agc cgc ttc cgc gtg ctg ccg cag ggg ctg aag gtg aag cag gtg	291
Trp Ser Arg Phe Arg Val Leu Pro Gln Gly Leu Lys Val Lys Gln Val	
75 80 85	
gag cgg gag gat gcc ggc gtg tac gtg tgc aag gcc acc aac ggc ttc	339
Glu Arg Glu Asp Ala Gly Val Tyr Val Cys Lys Ala Thr Asn Gly Phe	
90 95 100 105	
ggc agc ctt agc gtc aac tac acc ctc gtc gtg ctg gat gac att agc	387
Gly Ser Leu Ser Val Asn Tyr Thr Leu Val Val Leu Asp Asp Ile Ser	
110 115 120	
cca ggg aag gag agc ctg ggg ccc gac agc tcc tct ggg ggt caa gag	435
Pro Gly Lys Glu Ser Leu Gly Pro Asp Ser Ser Ser Gly Gly Gln Glu	
125 130 135	
gac ccc gcc agc cag cag tgg gca cga ccg cgc ttc aca cag ccc tcc	483
Asp Pro Ala Ser Gln Gln Trp Ala Arg Pro Arg Phe Thr Gln Pro Ser	
140 145 150	
aag atg agg cgc cgg gtg atc gca cgg ccc gtg ggt agc tcc gtg cgg	531
Lys Met Arg Arg Arg Val Ile Ala Arg Pro Val Gly Ser Ser Val Arg	
155 160 165	
ctc aag tgc gtg gcc agc ggg cac cct cgg ccc gac atc acg tgg atg	579
Leu Lys Cys Val Ala Ser Gly His Pro Arg Pro Asp Ile Thr Trp Met	
170 175 180 185	
aag gac gac cag gcc ttg acg cgc cca gag gcc gct gag ccc agg aag	627
Lys Asp Asp Gln Ala Leu Thr Arg Pro Glu Ala Ala Glu Pro Arg Lys	
190 195 200	
aag aag tgg aca ctg agc ctg aag aac ctg cgg ccg gag gac agc ggc	675
Lys Lys Trp Thr Leu Ser Leu Lys Asn Leu Arg Pro Glu Asp Ser Gly	
205 210 215	
aaa tac acc tgc cgc gtg tcg aac cgc gcg ggc gcc atc aac gcc acc	723
Lys Tyr Thr Cys Arg Val Ser Asn Arg Ala Gly Ala Ile Asn Ala Thr	
220 225 230	
tac aag gtg gat gtg atc cag cgg acc cgt tcc aag ccc gtg ctc aca	771
Tyr Lys Val Asp Val Ile Gln Arg Thr Arg Ser Lys Pro Val Leu Thr	
235 240 245	
ggc acg cac ccc gtg aac acg acg gtg gac ttc ggg ggg acc acg tcc	819
Gly Thr His Pro Val Asn Thr Thr Val Asp Phe Gly Gly Thr Thr Ser	
250 255 260 265	

ttc cag tgc aag gtg cgc agc gac gtg aag ccg gtg atc cag tgg ctg	867
Phe Gln Cys Lys Val Arg Ser Asp Val Lys Pro Val Ile Gln Trp Leu	
270 275 280	
aag cgc gtg gag tac ggc gcc gag ggc cgc cac aac tcc acc atc gat	915
Lys Arg Val Glu Tyr Gly Ala Glu Gly Arg His Asn Ser Thr Ile Asp	
285 290 295	
gtg ggc ggc cag aag ttt gtg gtg ctg ccc acg ggt gac gtg tgg tcg	963
Val Gly Gly Gln Lys Phe Val Val Leu Pro Thr Gly Asp Val Trp Ser	
300 305 310	
cgg ccc gac ggc tcc tac ctc aat aag ctg ctc atc acc cgt gcc cgc	1011
Arg Pro Asp Gly Ser Tyr Leu Asn Lys Leu Leu Ile Thr Arg Ala Arg	
315 320 325	
cag gac gat gcg ggc atg tac atc tgc ctt ggc gcc aac acc atg ggc	1059
Gln Asp Asp Ala Gly Met Tyr Ile Cys Leu Gly Ala Asn Thr Met Gly	
330 335 340 345	
tac agc ttc cgc agc gcc ttc ctc acc gtg ctg cca gac cca aaa ccg	1107
Tyr Ser Phe Arg Ser Ala Phe Leu Thr Val Leu Pro Asp Pro Lys Pro	
350 355 360	
caa ggg cca cct gtg gcc tcc tcg tcc tcg gcc act agc ctg ccg tgg	1155
Gln Gly Pro Pro Val Ala Ser Ser Ser Ser Ala Thr Ser Leu Pro Trp	
365 370 375	
ccc gtg gtc atc ggc atc cca gcc ggc gct gtc ttc atc ctg ggc acc	1203
Pro Val Val Ile Gly Ile Pro Ala Gly Ala Val Phe Ile Leu Gly Thr	
380 385 390	
ctg ctc ctg tgg ctt tgc cag gcc cag aag aag ccg tgc acc ccc gcg	1251
Leu Leu Leu Trp Leu Cys Gln Ala Gln Lys Lys Pro Cys Thr Pro Ala	
395 400 405	
cct gcc cct ccc ctg cct ggg cac cgc ccg ccg ggg acg gcc ctc gac	1299
Pro Ala Pro Pro Leu Pro Gly His Arg Pro Pro Gly Thr Ala Leu Asp	
410 415 420 425	
cgc agc gga gac aag gac ctt ccc tcg ttg gcc gcc ctc agc gct ggc	1347
Arg Ser Gly Asp Lys Asp Leu Pro Ser Leu Ala Ala Leu Ser Ala Gly	
430 435 440	
cct ggt gtg ggg ctg tgt gag gag cat ggg tct ccg gca gcc ccc cag	1395
Pro Gly Val Gly Leu Cys Glu Glu His Gly Ser Pro Ala Ala Pro Gln	
445 450 455	
cac tta ctg ggc cca ggc cca gtt gct ggc cct aag ttg tac ccc aaa	1443
His Leu Leu Gly Pro Gly Pro Val Ala Gly Pro Lys Leu Tyr Pro Lys	
460 465 470	
ctc tac aca gac atc cac aca cac aca cac aca cac tct cac aca cac	1491
Leu Tyr Thr Asp Ile His Thr His Thr His Thr His Ser His Thr His	
475 480 485	
tca cac gtg gag ggc aag gtc cac cag cac atc cac tat cag tgc	1536
Ser His Val Glu Gly Lys Val His Gln His Ile His Tyr Gln Cys	
490 495 500	

tagacggcac cgtatctgca gtgggcacgg gggggccggc cagacaggca gactgggagg 1596
atggaggacg gagctgcaga cgaaggcagg ggacccatgg cgaggaggaa tggccagcac 1656
cccaggcagt ctgtgtgtga ggcataagccc ctggacacac acacacagac acacacacta 1716
cctggatgca tgtatgcaca cacatgcgcg cacacgtgct ccctgaaggc acacgtacgc 1776
acacacgcac atgcacagat atgccgcctg ggcacacaga taagctgccc aaatgcacgc 1836
acacgcacag agacatgcc aacatacaa ggacatgctg cctgaacata cacacgcaca 1896
cccatgcgca gatgtgctgc ctggacacac acacacacac ggatatgctg tctggacgca 1956
cacacgtgca gatatggtat ccggacacac acgtgcacag atatgctgcc tggacacaca 2016
gataatgctg ccttgacaca cacatgcacg gatattgcct ggacacacac acacacacgc 2076
gtgcacagat atgctgtctg gacaggcaca cacatgcaga tatgctgcct ggacacacac 2136
ttccagacac acgtgcacag gcgcagatat gctgcctgga cacacgcaga tatgctgtct 2196
agtcacacac acacgcagac atgctgtccg gacacacaca cgcacgcaga gatatgctgt 2256
ccggacacac acacgcacgc agatatgctg cctggacaca cacacagata atgctgcctc 2316
aacactcaca cacgtgcaga tattgcctgg acacacacat gtgcacagat atgctgtctg 2376
gacatgcaca cacgtgcaga tatgctgtcc ggatacacac gcacgcacac atgcagatat 2436
gctgcctggg cacacacttc cggacacaca tgcacacaca ggtgcagata tgctgcctgg 2496
acacacgcag actgacgtgc ttttgggagg gtgtgccgtg aagcctgcag tacgtgtgcc 2556
gtgaggctca tagttgatga gggactttcc ctgctccacc gtcactcccc caactctgcc 2616
cgcctctgtc cccgcctcag tccccgcctc catccccgcc tctgtcccct ggccttggcg 2676
gctatTTTTg ccacctgcct tgggtgccca ggagtcccct actgctgtgg gctggggttg 2736
ggggcacagc agccccaagc ctgagaggct ggagcccatg gctagtggct catccccact 2796
gcattctccc cctgacacag agaagggggc ttggtattta tatttaagaa atgaagataa 2856
tattaataat gatggaagga agactgggtt gcagggactg tggctctctc tggggcccg 2916
gacccgcctg gtctttcagc catgctgatg accacacccc gtccaggcca gacaccacc 2976
cccacccac tgcgtggtg gcccagatc tctgtaattt tatgtagagt ttgagctgaa 3036
gccccgtata ttttaatttat tttgttaaac atgaaagtgc atcctttccc tccaaaaaaa 3096
aaaaaaaaa aaaaaa 3112

<210> 17
<211> 504
<212> PRT

<213> Homo sapiens

<400> 17

```

Met Thr Pro Ser Pro Leu Leu Leu Leu Leu Leu Pro Pro Leu Leu Leu
  1           5           10           15

Gly Ala Phe Pro Pro Ala Ala Ala Ala Arg Gly Pro Pro Lys Met Ala
          20           25           30

Asp Lys Val Val Pro Arg Gln Val Ala Arg Leu Gly Arg Thr Val Arg
          35           40           45

Leu Gln Cys Pro Val Glu Gly Asp Pro Pro Pro Leu Thr Met Trp Thr
          50           55           60

Lys Asp Gly Arg Thr Ile His Ser Gly Trp Ser Arg Phe Arg Val Leu
          65           70           75           80

Pro Gln Gly Leu Lys Val Lys Gln Val Glu Arg Glu Asp Ala Gly Val
          85           90           95

Tyr Val Cys Lys Ala Thr Asn Gly Phe Gly Ser Leu Ser Val Asn Tyr
          100          105          110

Thr Leu Val Val Leu Asp Asp Ile Ser Pro Gly Lys Glu Ser Leu Gly
          115          120          125

Pro Asp Ser Ser Ser Gly Gly Gln Glu Asp Pro Ala Ser Gln Gln Trp
          130          135          140

Ala Arg Pro Arg Phe Thr Gln Pro Ser Lys Met Arg Arg Arg Val Ile
          145          150          155          160

Ala Arg Pro Val Gly Ser Ser Val Arg Leu Lys Cys Val Ala Ser Gly
          165          170          175

His Pro Arg Pro Asp Ile Thr Trp Met Lys Asp Asp Gln Ala Leu Thr
          180          185          190

Arg Pro Glu Ala Ala Glu Pro Arg Lys Lys Lys Trp Thr Leu Ser Leu
          195          200          205

Lys Asn Leu Arg Pro Glu Asp Ser Gly Lys Tyr Thr Cys Arg Val Ser
          210          215          220

Asn Arg Ala Gly Ala Ile Asn Ala Thr Tyr Lys Val Asp Val Ile Gln
          225          230          235          240

Arg Thr Arg Ser Lys Pro Val Leu Thr Gly Thr His Pro Val Asn Thr
          245          250          255

Thr Val Asp Phe Gly Gly Thr Thr Ser Phe Gln Cys Lys Val Arg Ser
          260          265          270

Asp Val Lys Pro Val Ile Gln Trp Leu Lys Arg Val Glu Tyr Gly Ala
          275          280          285

Glu Gly Arg His Asn Ser Thr Ile Asp Val Gly Gly Gln Lys Phe Val
          290          295          300

```

Val Leu Pro Thr Gly Asp Val Trp Ser Arg Pro Asp Gly Ser Tyr Leu
 305 310 315 320
 Asn Lys Leu Leu Ile Thr Arg Ala Arg Gln Asp Asp Ala Gly Met Tyr
 325 330 335
 Ile Cys Leu Gly Ala Asn Thr Met Gly Tyr Ser Phe Arg Ser Ala Phe
 340 345 350
 Leu Thr Val Leu Pro Asp Pro Lys Pro Gln Gly Pro Pro Val Ala Ser
 355 360 365
 Ser Ser Ser Ala Thr Ser Leu Pro Trp Pro Val Val Ile Gly Ile Pro
 370 375 380
 Ala Gly Ala Val Phe Ile Leu Gly Thr Leu Leu Leu Trp Leu Cys Gln
 385 390 395 400
 Ala Gln Lys Lys Pro Cys Thr Pro Ala Pro Ala Pro Pro Leu Pro Gly
 405 410 415
 His Arg Pro Pro Gly Thr Ala Leu Asp Arg Ser Gly Asp Lys Asp Leu
 420 425 430
 Pro Ser Leu Ala Ala Leu Ser Ala Gly Pro Gly Val Gly Leu Cys Glu
 435 440 445
 Glu His Gly Ser Pro Ala Ala Pro Gln His Leu Leu Gly Pro Gly Pro
 450 455 460
 Val Ala Gly Pro Lys Leu Tyr Pro Lys Leu Tyr Thr Asp Ile His Thr
 465 470 475 480
 His Thr His Thr His Ser His Thr His Ser His Val Glu Gly Lys Val
 485 490 495
 His Gln His Ile His Tyr Gln Cys
 500

<210> 18
 <211> 3080
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (23)..(1534)

<400> 18
 ccccgaggtcc ggacaggccg ag atg acg ccg agc ccc ctg ttg ctg ctc ctg 52
 Met Thr Pro Ser Pro Leu Leu Leu Leu
 1 5 10
 ctg ccg ccg ctg ctg ctg ggg gcc ttc cca ccg gcc gcc gcc gcc cga 100
 Leu Pro Pro Leu Leu Leu Gly Ala Phe Pro Pro Ala Ala Ala Ala Arg
 15 20 25

ggc ccc cca aag atg gcg gac aag gtg gtc cca cgg cag gtg gcc cgg	148
Gly Pro Pro Lys Met Ala Asp Lys Val Val Pro Arg Gln Val Ala Arg	
30 35 40	
ctg ggc cgc act gtg cgg ctg cag tgc cca gtg gag ggg gac ccg ccg	196
Leu Gly Arg Thr Val Arg Leu Gln Cys Pro Val Glu Gly Asp Pro Pro	
45 50 55	
ccg ctg acc atg tgg acc aag gat ggc cgc acc atc cac agc ggc tgg	244
Pro Leu Thr Met Trp Thr Lys Asp Gly Arg Thr Ile His Ser Gly Trp	
60 65 70	
agc cgc ttc cgc gtg ctg ccg cag ggg ctg aag gtg aag cag gtg gag	292
Ser Arg Phe Arg Val Leu Pro Gln Gly Leu Lys Val Lys Gln Val Glu	
75 80 85 90	
cgg gag gat gcc ggc gtg tac gtg tgc aag gcc acc aac ggc ttc ggc	340
Arg Glu Asp Ala Gly Val Tyr Val Cys Lys Ala Thr Asn Gly Phe Gly	
95 100 105	
agc ctt agc gtc aac tac acc ctc gtc gtg ctg gat gac att agc cca	388
Ser Leu Ser Val Asn Tyr Thr Leu Val Val Leu Asp Asp Ile Ser Pro	
110 115 120	
ggg aag gag agc ctg ggg ccc gac agc tcc tct ggg ggt caa gag gac	436
Gly Lys Glu Ser Leu Gly Pro Asp Ser Ser Ser Gly Gly Gln Glu Asp	
125 130 135	
ccc gcc agc cag cag tgg gca cga ccg cgc ttc aca cag ccc tcc aag	484
Pro Ala Ser Gln Gln Trp Ala Arg Pro Arg Phe Thr Gln Pro Ser Lys	
140 145 150	
atg agg cgc cgg gtg atc gca cgg ccc gtg ggt agc tcc gtg cgg ctc	532
Met Arg Arg Arg Val Ile Ala Arg Pro Val Gly Ser Ser Val Arg Leu	
155 160 165 170	
aag tgc gtg gcc agc ggg cac cct cgg ccc gac atc acg tgg atg aag	580
Lys Cys Val Ala Ser Gly His Pro Arg Pro Asp Ile Thr Trp Met Lys	
175 180 185	
gac gac cag gcc ttg acg cgc cca gag gcc gct gag ccc agg aag aag	628
Asp Asp Gln Ala Leu Thr Arg Pro Glu Ala Ala Glu Pro Arg Lys Lys	
190 195 200	
aag tgg aca ctg agc ctg aag aac ctg cgg ccg gag gac agc ggc aaa	676
Lys Trp Thr Leu Ser Leu Lys Asn Leu Arg Pro Glu Asp Ser Gly Lys	
205 210 215	
tac acc tgc cgc gtg tgc aac cgc gcg ggc gcc atc aac gcc acc tac	724
Tyr Thr Cys Arg Val Ser Asn Arg Ala Gly Ala Ile Asn Ala Thr Tyr	
220 225 230	
aag gtg gat gtg atc cag cgg acc cgt tcc aag ccc gtg ctc aca ggc	772
Lys Val Asp Val Ile Gln Arg Thr Arg Ser Lys Pro Val Leu Thr Gly	
235 240 245 250	
acg cac ccc gtg aac acg acg gtg gac ttc ggg ggg acc acg tcc ttc	820
Thr His Pro Val Asn Thr Thr Val Asp Phe Gly Gly Thr Thr Ser Phe	
255 260 265	

cag tgc aag gtg cgc agc gac gtg aag ccg gtg atc cag tgg ctg aag	868
Gln Cys Lys Val Arg Ser Asp Val Lys Pro Val Ile Gln Trp Leu Lys	
270 275 280	
cgc gtg gag tac ggc gcc gag ggc cgc cac aac tcc acc atc gat gtg	916
Arg Val Glu Tyr Gly Ala Glu Gly Arg His Asn Ser Thr Ile Asp Val	
285 290 295	
ggc ggc cag aag ttt gtg gtg ctg ccc acg ggt gac gtg tgg tgc cgg	964
Gly Gly Gln Lys Phe Val Val Leu Pro Thr Gly Asp Val Trp Ser Arg	
300 305 310	
ccc gac ggc tcc tac ctc aat aag ctg ctc atc acc cgt gcc cgc cag	1012
Pro Asp Gly Ser Tyr Leu Asn Lys Leu Leu Ile Thr Arg Ala Arg Gln	
315 320 325 330	
gac gat gcg ggc atg tac atc tgc ctt ggc gcc aac acc atg ggc tac	1060
Asp Asp Ala Gly Met Tyr Ile Cys Leu Gly Ala Asn Thr Met Gly Tyr	
335 340 345	
agc ttc cgc agc gcc ttc ctc acc gtg ctg cca gac cca aaa ccg caa	1108
Ser Phe Arg Ser Ala Phe Leu Thr Val Leu Pro Asp Pro Lys Pro Gln	
350 355 360	
ggg cca cct gtg gcc tcc tgc tcc tgc gcc act agc ctg ccg tgg ccc	1156
Gly Pro Pro Val Ala Ser Ser Ser Ser Ala Thr Ser Leu Pro Trp Pro	
365 370 375	
gtg gtc atc ggc atc cca gcc ggc gct gtc ttc atc ctg ggc acc ctg	1204
Val Val Ile Gly Ile Pro Ala Gly Ala Val Phe Ile Leu Gly Thr Leu	
380 385 390	
ctc ctg tgg ctt tgc cag gcc cag aag aag ccg tgc acc ccc gcg cct	1252
Leu Leu Trp Leu Cys Gln Ala Gln Lys Lys Pro Cys Thr Pro Ala Pro	
395 400 405 410	
gcc cct ccc ctg cct ggg cac cgc ccg ccg ggg acg gcc cgc gac cgc	1300
Ala Pro Pro Leu Pro Gly His Arg Pro Pro Gly Thr Ala Arg Asp Arg	
415 420 425	
agc gga gac aag gac ctt ccc tgc ttg gcc gcc ctc agc gct ggc cct	1348
Ser Gly Asp Lys Asp Leu Pro Ser Leu Ala Ala Leu Ser Ala Gly Pro	
430 435 440	
ggc gtg ggg ctg tgt gag gag cat ggg tct ccg gca gcc ccc cag cac	1396
Gly Val Gly Leu Cys Glu Glu His Gly Ser Pro Ala Ala Pro Gln His	
445 450 455	
tta ctg ggc cca ggc cca gtt gct ggc cct aag ttg tac ccc aaa ctc	1444
Leu Leu Gly Pro Gly Pro Val Ala Gly Pro Lys Leu Tyr Pro Lys Leu	
460 465 470	
tac aca gac atc cac aca cac aca cac aca cac tct cac aca cac tca	1492
Tyr Thr Asp Ile His Thr His Thr His Thr His Ser His Thr His Ser	
475 480 485 490	
cac gtg gag ggc aag gtc cac cag cac atc cac tat cag tgc	1534
His Val Glu Gly Lys Val His Gln His Ile His Tyr Gln Cys	

495

500

tagacggcac cgtatctgca gtgggcacgg gggggccggc cagacaggca gactgggagg 1594
atggaggacg gagctgcaga cgaaggcagg ggacccatgg cgaggaggaa tggccagcac 1654
cccaggcagt ctgtgtgtga ggcatagccc ctggacacac acacacagac acacacacta 1714
cctggatgca tgtatgcaca cacatgcgcg cacacgtgct ccctgaaggc acacgtacgc 1774
acacacgcac atgcacagat atgccgcctg ggcacacaga taagctgccc aaatgcacgc 1834
acacgcacag agacatgcca gaacatacaa ggacatgctg cctgaacata cacacgcaca 1894
cccatgcgca gatgtgctgc ctggacacac acacacacac ggatatgctg tctggacgca 1954
cacacgtgca gatatggtat ccggacacac acgtgcacag atatgctgcc tggacacaca 2014
gataatgctg ccttgacaca cacatgcacg gatattgcct ggacacacac acacacacgc 2074
gtgcacagat atgctgtctg gacaggcaca cacatgcaga tatgctgcct ggacacacac 2134
ttccagacac acgtgcacag gcgcagatat gctgcctgga cacacgcaga tatgctgtct 2194
agtcacacac acacgcagac atgctgtccg gacacacaca cgcatgcaca gatatgctgt 2254
ccggacacac acacgcacgc agatatgctg cctggacaca cacacagata atgctgcctc 2314
aacactcaca cacgtgcaga tattgcctgg acacacacat gtgcacagat atgctgtctg 2374
gacatgcaca cacgtgcaga tatgctgtcc ggatacacac gcacgcacac atgcagatat 2434
gctgcctggg cacacacttc cggacacaca tgcacacaca ggtgcagata tgctgcctgg 2494
acacacgcag actgacgtgc ttttgggagg gtgtgccgtg aagcctgcag tacgtgtgcc 2554
gtgaggctca tagttgatga gggactttcc ctgctccacc gtcactcccc caactctgcc 2614
cgcctctgtc cccgcctcag tccccgcctc catccccgcc tctgtcccct ggccttggcg 2674
gctatttttg ccacctgcct tgggtgcccc ggagtcccct actgctgtgg gctgggggtg 2734
ggggcacagc agccccaagc ctgagaggct ggagcccatg gctagtggct catccccact 2794
gcattctccc cctgacacag agaagggggc ttggtattta tatttaagaa atgaagataa 2854
tattaataat gatggaagga agactgggtt gcagggactg tggctctctc tggggcccg 2914
gacccgcctg gtctttcagc catgctgatg accacacccc gtccaggcca gacaccaccc 2974
cccacccac tgctgtggtg gccccagatc tctgtaattt tatgtagagt ttgagctgaa 3034
gccccgtata ttttaatttat tttgttaaac atgaaagtgc atcctt 3080

<210> 19

<211> 504

<212> PRT

<213> Homo sapiens

<400> 19

```

Met Thr Pro Ser Pro Leu Leu Leu Leu Leu Leu Pro Pro Leu Leu Leu
  1          5          10          15

Gly Ala Phe Pro Pro Ala Ala Ala Ala Arg Gly Pro Pro Lys Met Ala
          20          25          30

Asp Lys Val Val Pro Arg Gln Val Ala Arg Leu Gly Arg Thr Val Arg
          35          40          45

Leu Gln Cys Pro Val Glu Gly Asp Pro Pro Pro Leu Thr Met Trp Thr
          50          55          60

Lys Asp Gly Arg Thr Ile His Ser Gly Trp Ser Arg Phe Arg Val Leu
          65          70          75          80

Pro Gln Gly Leu Lys Val Lys Gln Val Glu Arg Glu Asp Ala Gly Val
          85          90          95

Tyr Val Cys Lys Ala Thr Asn Gly Phe Gly Ser Leu Ser Val Asn Tyr
          100          105          110

Thr Leu Val Val Leu Asp Asp Ile Ser Pro Gly Lys Glu Ser Leu Gly
          115          120          125

Pro Asp Ser Ser Ser Gly Gly Gln Glu Asp Pro Ala Ser Gln Gln Trp
          130          135          140

Ala Arg Pro Arg Phe Thr Gln Pro Ser Lys Met Arg Arg Arg Val Ile
          145          150          155          160

Ala Arg Pro Val Gly Ser Ser Val Arg Leu Lys Cys Val Ala Ser Gly
          165          170          175

His Pro Arg Pro Asp Ile Thr Trp Met Lys Asp Asp Gln Ala Leu Thr
          180          185          190

Arg Pro Glu Ala Ala Glu Pro Arg Lys Lys Lys Trp Thr Leu Ser Leu
          195          200          205

Lys Asn Leu Arg Pro Glu Asp Ser Gly Lys Tyr Thr Cys Arg Val Ser
          210          215          220

Asn Arg Ala Gly Ala Ile Asn Ala Thr Tyr Lys Val Asp Val Ile Gln
          225          230          235          240

Arg Thr Arg Ser Lys Pro Val Leu Thr Gly Thr His Pro Val Asn Thr
          245          250          255

Thr Val Asp Phe Gly Gly Thr Thr Ser Phe Gln Cys Lys Val Arg Ser
          260          265          270

Asp Val Lys Pro Val Ile Gln Trp Leu Lys Arg Val Glu Tyr Gly Ala
          275          280          285

Glu Gly Arg His Asn Ser Thr Ile Asp Val Gly Gly Gln Lys Phe Val
          290          295          300

```

Val Leu Pro Thr Gly Asp Val Trp Ser Arg Pro Asp Gly Ser Tyr Leu
 305 310 315 320
 Asn Lys Leu Leu Ile Thr Arg Ala Arg Gln Asp Asp Ala Gly Met Tyr
 325 330 335
 Ile Cys Leu Gly Ala Asn Thr Met Gly Tyr Ser Phe Arg Ser Ala Phe
 340 345 350
 Leu Thr Val Leu Pro Asp Pro Lys Pro Gln Gly Pro Pro Val Ala Ser
 355 360 365
 Ser Ser Ser Ala Thr Ser Leu Pro Trp Pro Val Val Ile Gly Ile Pro
 370 375 380
 Ala Gly Ala Val Phe Ile Leu Gly Thr Leu Leu Leu Trp Leu Cys Gln
 385 390 395 400
 Ala Gln Lys Lys Pro Cys Thr Pro Ala Pro Ala Pro Pro Leu Pro Gly
 405 410 415
 His Arg Pro Pro Gly Thr Ala Arg Asp Arg Ser Gly Asp Lys Asp Leu
 420 425 430
 Pro Ser Leu Ala Ala Leu Ser Ala Gly Pro Gly Val Gly Leu Cys Glu
 435 440 445
 Glu His Gly Ser Pro Ala Ala Pro Gln His Leu Leu Gly Pro Gly Pro
 450 455 460
 Val Ala Gly Pro Lys Leu Tyr Pro Lys Leu Tyr Thr Asp Ile His Thr
 465 470 475 480
 His Thr His Thr His Ser His Thr His Ser His Val Glu Gly Lys Val
 485 490 495
 His Gln His Ile His Tyr Gln Cys
 500

<210> 20
 <211> 342
 <212> PRT
 <213> Mus musculus

<400> 20
 Met Ala Asp Lys Val Val Pro Arg Gln Val Ala Arg Leu Gly Arg Thr
 1 5 10 15
 Val Arg Leu Gln Cys Pro Val Glu Gly Asp Pro Pro Pro Leu Thr Met
 20 25 30
 Trp Thr Lys Asp Gly Arg Thr Ile His Ser Gly Trp Ser Arg Phe Arg
 35 40 45
 Val Leu Pro Gln Gly Leu Lys Val Lys Glu Val Glu Ala Glu Asp Ala
 50 55 60
 Gly Val Tyr Val Cys Lys Ala Thr Asn Gly Phe Gly Ser Leu Ser Val

65					70						75				80
Asn	Tyr	Thr	Leu	Ile	Ile	Met	Asp	Asp	Ile	Ser	Pro	Gly	Lys	Glu	Ser
				85					90					95	
Pro	Gly	Pro	Gly	Gly	Ser	Ser	Gly	Gly	Gln	Glu	Asp	Pro	Ala	Ser	Gln
			100					105					110		
Gln	Trp	Ala	Arg	Pro	Arg	Phe	Thr	Gln	Pro	Ser	Lys	Met	Arg	Arg	Arg
		115					120					125			
Val	Ile	Ala	Arg	Pro	Val	Gly	Ser	Ser	Val	Arg	Leu	Lys	Cys	Val	Ala
	130					135					140				
Ser	Gly	His	Pro	Arg	Pro	Asp	Ile	Met	Trp	Met	Lys	Asp	Asp	Gln	Thr
145					150					155					160
Leu	Thr	His	Leu	Glu	Ala	Ser	Glu	His	Arg	Lys	Lys	Lys	Trp	Thr	Leu
			165						170					175	
Ser	Leu	Lys	Asn	Leu	Lys	Pro	Glu	Asp	Ser	Gly	Lys	Tyr	Thr	Cys	Arg
			180					185					190		
Val	Ser	Asn	Lys	Ala	Gly	Ala	Ile	Asn	Ala	Thr	Tyr	Lys	Val	Asp	Val
		195					200					205			
Ile	Gln	Arg	Thr	Arg	Ser	Lys	Pro	Val	Leu	Thr	Gly	Thr	His	Pro	Val
	210					215					220				
Asn	Thr	Thr	Val	Asp	Phe	Gly	Gly	Thr	Thr	Ser	Phe	Gln	Cys	Lys	Val
225					230					235					240
Arg	Ser	Asp	Val	Lys	Pro	Val	Ile	Gln	Trp	Leu	Lys	Arg	Val	Glu	Tyr
			245						250					255	
Gly	Ser	Glu	Gly	Arg	His	Asn	Ser	Thr	Ile	Asp	Val	Gly	Gly	Gln	Lys
			260					265					270		
Phe	Val	Val	Leu	Pro	Thr	Gly	Asp	Val	Trp	Ser	Arg	Pro	Asp	Gly	Ser
		275					280					285			
Tyr	Leu	Asn	Lys	Leu	Leu	Ile	Ser	Arg	Ala	Arg	Gln	Asp	Asp	Ala	Gly
	290					295					300				
Met	Tyr	Ile	Cys	Leu	Gly	Ala	Asn	Thr	Met	Gly	Tyr	Ser	Phe	Arg	Ser
305					310					315					320
Ala	Phe	Leu	Thr	Val	Leu	Pro	Asp	Pro	Lys	Pro	Pro	Gly	Pro	Pro	Met
				325					330					335	
Ala	Ser	Ser	Ser	Ser	Ser	Ser									
				340											

<210> 21

<211> 1788

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: murine FGFR-L
extracellular domain-Fc fusion polypeptide

<220>

<221> CDS

<222> (1)..(1782)

<400> 21

atg acg cgg agc ccc gcg ctg ctg ctg ctg cta ttg ggg gcc ctc ccg	48
Met Thr Arg Ser Pro Ala Leu Leu Leu Leu Leu Leu Gly Ala Leu Pro	
1 5 10 15	
tcg gct gag gcg gcg cga gga ccc cca aga atg gca gac aaa gtg gtc	96
Ser Ala Glu Ala Ala Arg Gly Pro Pro Arg Met Ala Asp Lys Val Val	
20 25 30	
cca cgg cag gtg gcc cgc ctg ggc cgc act gtg cgg cta cag tgc cca	144
Pro Arg Gln Val Ala Arg Leu Gly Arg Thr Val Arg Leu Gln Cys Pro	
35 40 45	
gtg gag ggg gac cca cca ccg ttg acc atg tgg acc aaa gat ggc cgc	192
Val Glu Gly Asp Pro Pro Pro Leu Thr Met Trp Thr Lys Asp Gly Arg	
50 55 60	
aca atc cac agt ggc tgg agc cgc ttc cgt gtg ctg ccc cag ggt ctg	240
Thr Ile His Ser Gly Trp Ser Arg Phe Arg Val Leu Pro Gln Gly Leu	
65 70 75 80	
aag gtg aag gag gtg gag gcc gag gat gcc ggt gtt tat gtg tgc aag	288
Lys Val Lys Glu Val Glu Ala Glu Asp Ala Gly Val Tyr Val Cys Lys	
85 90 95	
gcc acc aat ggc ttt ggc agc ctc agc gtc aac tac act ctc atc atc	336
Ala Thr Asn Gly Phe Gly Ser Leu Ser Val Asn Tyr Thr Leu Ile Ile	
100 105 110	
atg gat gat att agt cca ggg aag gag agc cct ggg cca ggt ggt tct	384
Met Asp Asp Ile Ser Pro Gly Lys Glu Ser Pro Gly Pro Gly Gly Ser	
115 120 125	
tcg ggg ggc cag gag gac cca gcc agc cag cag tgg gca cgg cct cgc	432
Ser Gly Gly Gln Glu Asp Pro Ala Ser Gln Gln Trp Ala Arg Pro Arg	
130 135 140	
ttc aca cag ccc tcc aag atg agg cgc cga gtg att gca cgg cct gtg	480
Phe Thr Gln Pro Ser Lys Met Arg Arg Arg Val Ile Ala Arg Pro Val	
145 150 155 160	
ggt agc tct gtg cgg ctc aag tgt gtg gcc agt ggg cac cca cgg cca	528
Gly Ser Ser Val Arg Leu Lys Cys Val Ala Ser Gly His Pro Arg Pro	
165 170 175	
gac atc atg tgg atg aag gat gac cag acc ttg acg cat cta gag gct	576
Asp Ile Met Trp Met Lys Asp Asp Gln Thr Leu Thr His Leu Glu Ala	
180 185 190	
agt gaa cac aga aag aag aag tgg aca ctg agc ttg aag aac ctg aag	624
Ser Glu His Arg Lys Lys Lys Trp Thr Leu Ser Leu Lys Asn Leu Lys	

195	200	205	
cct gaa gac agt ggc aag tac acg tgc cgt gta tct aac aag gcc ggt Pro Glu Asp Ser Gly Lys Tyr Thr Cys Arg Val Ser Asn Lys Ala Gly 210 215 220			672
gcc atc aac gcc acc tac aaa gtg gat gta atc cag cgg act cgt tcc Ala Ile Asn Ala Thr Tyr Lys Val Asp Val Ile Gln Arg Thr Arg Ser 225 230 235 240			720
aag cct gtg ctc aca ggg aca cac cct gtg aac aca acg gtg gac ttc Lys Pro Val Leu Thr Gly Thr His Pro Val Asn Thr Thr Val Asp Phe 245 250 255			768
ggg ggg aca acg tcc ttc cag tgc aag gtg cgc agt gac gtg aag cct Gly Gly Thr Thr Ser Phe Gln Cys Lys Val Arg Ser Asp Val Lys Pro 260 265 270			816
gtg atc cag tgg ctg aag cgg gtg gag tac ggc tcc gag gga cgc cac Val Ile Gln Trp Leu Lys Arg Val Glu Tyr Gly Ser Glu Gly Arg His 275 280 285			864
aac tcc acc att gat gtg ggt ggc cag aag ttt gtg gtg ttg ccc acg Asn Ser Thr Ile Asp Val Gly Gly Gln Lys Phe Val Val Leu Pro Thr 290 295 300			912
ggg gat gtg tgg tca cgg cct gat ggc tcc tac ctc aac aag ctg ctc Gly Asp Val Trp Ser Arg Pro Asp Gly Ser Tyr Leu Asn Lys Leu Leu 305 310 315 320			960
atc tct cgg gcc cgc cag gat gat gct ggc atg tac atc tgc cta ggt Ile Ser Arg Ala Arg Gln Asp Asp Ala Gly Met Tyr Ile Cys Leu Gly 325 330 335			1008
gca aat acc atg ggc tac agt ttc cgt agc gcc ttc ctc act gta tta Ala Asn Thr Met Gly Tyr Ser Phe Arg Ser Ala Phe Leu Thr Val Leu 340 345 350			1056
cca gac ccc aaa cct cca ggg cct cct atg gct tct tca tgc gtc gac Pro Asp Pro Lys Pro Pro Gly Pro Pro Met Ala Ser Ser Ser Val Asp 355 360 365			1104
aaa act cac aca tgc cca ccg tgc cca gca cct gaa ctc ctg ggg gga Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly 370 375 380			1152
ccg tca gtc ttc ctc ttc ccc cca aaa ccc aag gac acc ctc atg atc Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile 385 390 395 400			1200
tcc cgg acc cct gag gtc aca tgc gtg gtg gtg gac gtg agc cac gaa Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu 405 410 415			1248
gac cct gag gtc aag ttc aac tgg tac gtg gac ggc gtg gag gtg cat Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His 420 425 430			1296
aat gcc aag aca aag ccg cgg gag gag cag tac aac agc acg tac cgt			1344

Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
 435 440 445

gtg gtc agc gtc ctc acc gtc ctg cac cag gac tgg ctg aat ggc aag 1392
 Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
 450 455 460

gag tac aag tgc aag gtc tcc aac aaa gcc ctc cca gcc ccc atc gag 1440
 Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
 465 470 475 480

aaa acc atc tcc aaa gcc aaa ggg cag ccc cga gaa cca cag gtg tac 1488
 Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
 485 490 495

acc ctg ccc cca tcc cgg gat gag ctg acc aag aac cag gtc agc ctg 1536
 Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu
 500 505 510

acc tgc ctg gtc aaa ggc ttc tat ccc agc gac atc gcc gtg gag tgg 1584
 Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
 515 520 525

gag agc aat ggg cag ccg gag aac aac tac aag acc acg cct ccc gtg 1632
 Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
 530 535 540

ctg gac tcc gac ggc tcc ttc ttc ctc tac agc aag ctc acc gtg gac 1680
 Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
 545 550 555 560

aag agc agg tgg cag cag ggg aac gtc ttc tca tgc tcc gtg atg cat 1728
 Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
 565 570 575

gag gct ctg cac aac cac tac acg cag aag agc ctc tcc ctg tct ccg 1776
 Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
 580 585 590

ggt aaa tgataa 1788
 Gly Lys

<210> 22

<211> 594

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: murine FGFR-L
 extracellular domain-Fc fusion polypeptide

<400> 22

Met Thr Arg Ser Pro Ala Leu Leu Leu Leu Leu Leu Gly Ala Leu Pro
 1 5 10 15

Ser Ala Glu Ala Ala Arg Gly Pro Pro Arg Met Ala Asp Lys Val Val
 20 25 30

Pro Arg Gln Val Ala Arg Leu Gly Arg Thr Val Arg Leu Gln Cys Pro
 35 40 45
 Val Glu Gly Asp Pro Pro Pro Leu Thr Met Trp Thr Lys Asp Gly Arg
 50 55 60
 Thr Ile His Ser Gly Trp Ser Arg Phe Arg Val Leu Pro Gln Gly Leu
 65 70 75 80
 Lys Val Lys Glu Val Glu Ala Glu Asp Ala Gly Val Tyr Val Cys Lys
 85 90 95
 Ala Thr Asn Gly Phe Gly Ser Leu Ser Val Asn Tyr Thr Leu Ile Ile
 100 105 110
 Met Asp Asp Ile Ser Pro Gly Lys Glu Ser Pro Gly Pro Gly Gly Ser
 115 120 125
 Ser Gly Gly Gln Glu Asp Pro Ala Ser Gln Gln Trp Ala Arg Pro Arg
 130 135 140
 Phe Thr Gln Pro Ser Lys Met Arg Arg Arg Val Ile Ala Arg Pro Val
 145 150 155 160
 Gly Ser Ser Val Arg Leu Lys Cys Val Ala Ser Gly His Pro Arg Pro
 165 170 175
 Asp Ile Met Trp Met Lys Asp Asp Gln Thr Leu Thr His Leu Glu Ala
 180 185 190
 Ser Glu His Arg Lys Lys Lys Trp Thr Leu Ser Leu Lys Asn Leu Lys
 195 200 205
 Pro Glu Asp Ser Gly Lys Tyr Thr Cys Arg Val Ser Asn Lys Ala Gly
 210 215 220
 Ala Ile Asn Ala Thr Tyr Lys Val Asp Val Ile Gln Arg Thr Arg Ser
 225 230 235 240
 Lys Pro Val Leu Thr Gly Thr His Pro Val Asn Thr Thr Val Asp Phe
 245 250 255
 Gly Gly Thr Thr Ser Phe Gln Cys Lys Val Arg Ser Asp Val Lys Pro
 260 265 270
 Val Ile Gln Trp Leu Lys Arg Val Glu Tyr Gly Ser Glu Gly Arg His
 275 280 285
 Asn Ser Thr Ile Asp Val Gly Gly Gln Lys Phe Val Val Leu Pro Thr
 290 295 300
 Gly Asp Val Trp Ser Arg Pro Asp Gly Ser Tyr Leu Asn Lys Leu Leu
 305 310 315 320
 Ile Ser Arg Ala Arg Gln Asp Asp Ala Gly Met Tyr Ile Cys Leu Gly
 325 330 335
 Ala Asn Thr Met Gly Tyr Ser Phe Arg Ser Ala Phe Leu Thr Val Leu
 340 345 350

Pro Asp Pro Lys Pro Pro Gly Pro Pro Met Ala Ser Ser Ser Val Asp
 355 360 365
 Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
 370 375 380
 Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
 385 390 395 400
 Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
 405 410 415
 Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
 420 425 430
 Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
 435 440 445
 Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
 450 455 460
 Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
 465 470 475 480
 Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
 485 490 495
 Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu
 500 505 510
 Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
 515 520 525
 Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
 530 535 540
 Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
 545 550 555 560
 Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
 565 570 575
 Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
 580 585 590
 Gly Lys

THIS PAGE BLANK (USPTO)